

REALIZING ERGODIC PROPERTIES IN ZERO ENTROPY SUBSHIFTS

VAN CYR AND BRYNA KRA

ABSTRACT. A subshift with linear block complexity has at most countably many ergodic measures, and we continue of the study of the relation between such complexity and the invariant measures. By constructing minimal subshifts whose block complexity is arbitrarily close to linear but has uncountably many ergodic measures, we show that this behavior fails as soon as the block complexity is superlinear. With a different construction, we show that there exists a minimal subshift with an ergodic measure whose slow entropy grows slower than any given rate tending to infinitely but faster than any other rate majorizing this one yet still growing subexponentially. These constructions lead to obstructions in using subshifts in applications to properties of the prime numbers and in finding a measurable version of the complexity gap that arises for shifts of sublinear complexity.

1. INTRODUCTION

Assume that (X, σ) is a subshift over the finite alphabet \mathcal{A} , meaning that $X \subset \mathcal{A}^{\mathbb{Z}}$ is a closed set that is invariant under the left shift $\sigma: \mathcal{A}^{\mathbb{Z}} \rightarrow \mathcal{A}^{\mathbb{Z}}$. The block complexity $p_X(n)$ of the shift is defined to be the number of words of length n which occur in any $x \in X$. Boshernitzan [1] showed that a minimal subshift with linear block complexity has only finitely many ergodic measures, where the number depends on the complexity growth. In [2], we showed that any subshift (minimal or not) with linear block complexity has at most finitely many nonatomic ergodic measures, and so at most countably many ergodic measures (with no requirement that the measures are nonatomic). In the same article, we give examples of subshifts with block complexity arbitrarily close to linear which have uncountably many nonatomic ergodic measures. Our main result is to show there is no complexity bound beyond linear on a subshift that suffices for guaranteeing there are at most countably many ergodic measures. More precisely, we show as soon as the growth is superlinear, we can have the maximal number of ergodic measures:

Theorem 1.1. *If $(p_n)_{n \in \mathbb{N}}$ is a sequence of natural numbers such that*

$$\liminf_{n \rightarrow \infty} \frac{p_n}{n} = \infty,$$

then there exists a minimal subshift X which supports uncountably many ergodic measures and is such that

$$\liminf_{n \rightarrow \infty} \frac{P_X(n)}{p_n} = 0.$$

2010 *Mathematics Subject Classification.* 37B10 (primary), 37A05, 37A35.

Key words and phrases. subshift, block complexity, slow entropy.

The second author was partially supported by NSF grant 1800544.

The distinction between countably and uncountably many ergodic measures supported by a subshift has recently received attention, as it plays a role in the deep results of Frantzikinakis and Host [4] on the complexity of the Liouville shift. More precisely, by studying the subshift naturally associated to Liouville function $\lambda(n)$ (see Section 2.4) and the number of ergodic measures it supports, they conclude that the Liouville function has superlinear complexity. Given the example we construct in Theorem 1.1, any approach to showing that the Liouville function has higher growth rates must rely on further properties of the shift beyond the cardinality of the set ergodic measures supported by subshifts whose complexity grows at a given rate. In a further development, using different methods McNamara [7] has shown that the Liouville function has at least quadratic complexity.

Katok and Thouvenot [6] and Ferenczi [3] defined the slow entropy, as a measure theoretic invariant to capture a measurable version of the (topological) complexity. They defined two growth rates, $P_T^-(n)$ and $P_T^+(n)$ of a measure preserving system (X, μ, T) and whether each these growth rates is slower \prec or faster \succ than a given growth rate is a measure theoretic invariant for the system and computable using a generating partition (see Section 4 for precise definitions). Using a different construction, we exhibit the freedom on growth rate of the slow entropy, with a minimal subshift of zero topological entropy such that the slow entropy grows slower than some (arbitrarily slowly growing) given sequence while faster than another (arbitrarily quickly within the class of subexponentially growing) given sequence:

Theorem 1.2. *Assume $(a_n)_{n \in \mathbb{N}}$ and $(b_n)_{n \in \mathbb{N}}$ are two non-decreasing sequences of positive integers such that $\lim_{n \rightarrow \infty} a_n = \infty$, $\lim_{n \rightarrow \infty} \frac{1}{n} \cdot \log(b_n) = 0$, and $a_n \leq b_n$ for all $n \in \mathbb{N}$. There exists a minimal subshift (X_∞, σ) of topological entropy zero and an ergodic measure μ supported on X_∞ such that*

$$P_\sigma^-(n) \prec (a_n) \text{ and } P_\sigma^+(n) \succ (b_n).$$

If we only consider the upper growth rate $P_T^+(n)$, related constructions are given in Katok and Thouvenot [6] and Serafin [9] of a subshift with zero topological entropy and $P_T^+(n)$ growing arbitrarily quickly (but still subexponentially growing). The restrictions given in our theorem on the sequences $(a_n)_{n \in \mathbb{N}}$ and $(b_n)_{n \in \mathbb{N}}$ are the weakest for which a result like ours could hold, in a precise sense that we explain (see Section 4). By a theorem of Ferenczi [3], it follows from the lower bound on the growth $P_\sigma^+(n) \succ (b_n)$, that the system we construct is not a Kronecker system. From the upper bound $P_\sigma^-(n) \prec (a_n)$, it follows that there is no sequence (a_n) increasing to infinity and such that the analogous bound would give $P_\sigma^-(n) \prec (c_n)$ for all sequences (c_n) increasing to infinity. This exhibits a different behavior than what happens in the topological setting. Namely, the Morse-Hedlund Theorem states that if there exists some $n \in \mathbb{N}$ such that $P_X(n) \leq n$ for some system (X, σ) , then we have that the topological complexity function $P_X(n)$ is actually bounded for all $n \in \mathbb{N}$. Our construction shows that the measurable analog of the result fails. Since $P_\sigma^+(n) \succ (b_n)$, by an approximation argument we have that the topological complexity $P_X(n) \geq b_n$ for infinitely many n . In particular, there is no subexponentially growing sequence $(b_n)_{n \in \mathbb{N}}$ such that any subshift whose word complexity exceeds b_n for infinitely many n must have positive entropy. This can be viewed as a counterexample to a dual version of the Morse-Hedlund Theorem, meaning there exists no subexponentially growing sequence that asymptotically bounds the complexity of every zero entropy subshift.

2. BACKGROUND AND NOTATION

2.1. Symbolic Dynamics. Assume that \mathcal{A} is a finite set endowed with the discrete topology, and we call \mathcal{A} the *alphabet*. The space $\mathcal{A}^{\mathbb{Z}}$, endowed with the product topology, is a compact, metrizable space. An element $x \in \mathcal{A}^{\mathbb{Z}}$ denotes a bi-infinite sequence in the alphabet \mathcal{A} , meaning that $x = (x_i)_{i \in \mathbb{Z}}$ with each $x_i \in \mathcal{A}$. The left shift $\sigma: \mathcal{A}^{\mathbb{Z}} \rightarrow \mathcal{A}^{\mathbb{Z}}$ defined by $(\sigma x)_i := x_{i+1}$ is continuous and the dynamical system $(\mathcal{A}^{\mathbb{Z}}, \sigma)$ is the *full \mathcal{A} -shift*. A *subshift* $X \in \mathcal{A}^{\mathbb{Z}}$ is the restriction of σ to any closed, σ -invariant set X .

2.2. Words and complexity. If $w = (w_0, \dots, w_{n-1}) \in \mathcal{A}^n$, the *cylinder set* $[w]$ in $\mathcal{A}^{\mathbb{Z}}$ determined by w is defined to be

$$[w] := \{x \in \mathcal{A}^{\mathbb{Z}} : x_i = w_i \text{ for all } 0 \leq i \leq n-1\}.$$

If $X \subseteq \mathcal{A}^{\mathbb{Z}}$ is a subshift, then the *language* $\mathcal{L}(X)$ of X is the set of all words $w \in \mathcal{A}^*$ such that $[w] \cap X \neq \emptyset$. For $n \in \mathbb{N}$, the set $\mathcal{L}_n(X)$ of *words of length n* in X is the set

$$\mathcal{L}_n(X) := \{w \in \mathcal{A}^n : [w] \cap X \neq \emptyset\}.$$

The *block complexity* (also known as the *word complexity function*) $p_X: \mathbb{N} \rightarrow \mathbb{N}$ of X is the function that counts the number of words of each length in X . Thus

$$p_X(n) := |\mathcal{L}_n(X)|.$$

2.3. The natural subshift associated to a language. Given a subshift (X, σ) , its language $\mathcal{L}(X)$ satisfies the properties that:

- (i) If $w \in \mathcal{L}(X)$, then every subword of w also belongs to $\mathcal{L}(X)$.
- (ii) If $w \in \mathcal{L}(X)$, then there exist nonempty words $u, v \in \mathcal{L}(X)$ such that $uwv \in \mathcal{L}(X)$.

Conversely, if \mathcal{L} is any collection of words over an alphabet \mathcal{A} satisfying condition (i), then $\mathcal{L} = \mathcal{L}(X)$ for some subshift (X, σ) . Thus the language of a subshift determines the subshift, and we can define a subshift by specifying its language. When we do so, we say that (X, σ) is the subshift defined by the language \mathcal{L} .

Moreover, given a finite list of words v_1, \dots, v_k , we can consider the collection of all bi-infinite concatenations of these words, and by taking the shifts of these bi-infinite sequences, we obtain a subshift (X, σ) . We refer to this as the *subshift defined by the words v_1, \dots, v_k* .

2.4. The natural subshift associated to a sequence. Suppose $h: \mathbb{N} \rightarrow \mathcal{A}$ is a function. Fix some $a \in \mathcal{A}$ and define

$$y := \begin{cases} y_i = h(i) & \text{if } i > 0; \\ y_i = a & \text{if } i \leq 0. \end{cases}$$

Then the set

$$Y_h := \overline{\{\sigma^n y : n \in \mathbb{Z}\}}$$

is a subshift. Note that Y_h contains at most n words of length n that are not found in contiguous blocks of the function $h: \mathbb{N} \rightarrow \mathcal{A}$. Therefore number of such words differs from $P_{Y_h}(n)$ by at most n . A word $w \in \mathcal{L}(Y)$ if and only if there are arbitrarily large $m \in \mathbb{N}$ such that $w_i = h(m+i)$ for all $0 \leq i < |w|$. The resulting shift (Y_h, σ) is *transitive*, meaning that it has a point with dense orbit under the shift (note that the point y has dense orbit).

Of particular interest are functions h that arise in number theory. For example, we can consider h to be the Liouville function $\lambda(n)$, the sequence in the alphabet $\{-1, 1\}$ with $\lambda(n) = 1$ if n has an even number of prime factors counted with multiplicity and otherwise is -1 , or take h to be the Möbius function $\mu(n)$, the sequence in the alphabet $\{-1, 0, 1\}$ with $\mu(n) = 1$ if n is square free and has an even number of prime factors, $\mu(n) = -1$ if n is square free and has an odd number of prime factors, and otherwise $\mu(n) = 0$. Then studying the language of Y_h gives insight into the number theoretical properties of h .

2.5. Invariant measures on certain subshifts. A Borel measure ν supported in X is *invariant* if $\nu(A) = \nu(\sigma^{-1}A)$ for all Borel sets $A \subseteq X$ and is *ergodic* if $\nu(A) \cdot \nu(X \setminus A) = 0$ whenever $A = \sigma^{-1}A$. It was recently shown [4] that if (Y, T) is a topological dynamical system of entropy zero and if Y supports at most countably many ergodic measures, then (Y, T) satisfies a logarithmic variant of the Möbius Disjointness Conjecture of Sarnak. Namely, under these conditions, for every $y \in Y$ and every $f \in C(Y)$ we have

$$\lim_{N \rightarrow \infty} \frac{1}{\log N} \sum_{n=1}^N \frac{f(T^n y) \mu(n)}{n} = 0$$

where μ is the Möbius function (and the same conclusion holds if μ is replaced with the Liouville function λ). This result was leveraged in [4] to give a lower bound on the growth of words in the language of the *Liouville shift*: Y_λ , showing that the Liouville shift has superlinear block growth (note again that the question of whether $P_{Y_\lambda}(n)$ grows superlinearly is equivalent to the question of whether the number of words of length n that occur in the Liouville sequence grows superlinearly). Along with the deep result on logarithmic disjointness that they prove, they rely on a bound on the number of ergodic measures that can be supported by a subshift with linear growth.

As a corollary of our result in Theorem 1.1, any proof showing that the Liouville shift has a growth rate that is faster than some explicit superlinear bound necessarily relies on deeper information from number theory, rather than only on estimates on the number of ergodic measures.

3. PROOF OF THEOREM 1.1

Suppose (p_n) is a sequence of natural numbers satisfying

$$\liminf_{n \rightarrow \infty} \frac{p_n}{n} = \infty.$$

We construct a minimal subshift X , depending on the sequence (p_n) , that supports uncountably many ergodic measures and is such that

$$\limsup_{n \rightarrow \infty} \frac{P_X(n)}{p_n} = 0.$$

We build the system inductively, by constructing words at each level and then using these words to define the language of a subshift. Throughout we use superscripts to denote the level of the construction and subscripts to denote the words constructed at this level.

Step 1: the base step of the construction. Set $\mathcal{A} = \{0, 1\}$. Define

$$w_1^1 = \underbrace{00 \cdots 00}_{{N_1 \text{ times}}1}$$

and

$$w_2^1 = \underbrace{011 \cdots 11}_{{N_1 \text{ times}}},$$

where N_1 is a large integer to be determined later. Note that the words w_1^1 and w_2^1 are distinct, as can be seen from the different frequencies of 0's and 1's. Define $X_1 \subseteq \mathcal{A}^{\mathbb{Z}}$ to be the subshift of $\mathcal{A}^{\mathbb{Z}}$ consisting of all $x \in \mathcal{A}^{\mathbb{Z}}$ that can be written as bi-infinite concatenations of the words w_1^1 and w_2^1 . Consider $n_1 = \lfloor N_1/2 \rfloor$ (note that $\lfloor \cdot \rfloor$ denotes the floor function). Then $\mathcal{L}_{n_1}(X_1)$ contains a word of all 0's, a word of all 1's, all words with exactly one 1 (by concatenating w_1^1 with itself), all words with exactly one 0 (by concatenating w_2^1 with itself), all words that begin with a continuous string of 0's followed by a continuous string of 1's (by concatenating w_1^1 and w_2^1), and all words with a continuous string of 1's followed by a continuous string of 0's (by concatenating w_2^1 and w_1^1). Thus for $n_1 = \lfloor N_1/2 \rfloor$,

$$P_{X_1}(n_1) = 4n_1 - 2$$

and so if $Y \subseteq X_1$ is any subshift, then $P_Y(n_1) \leq 4n_1 - 2$.

Step 2: the inductive step. Assume we have constructed a nested sequence of subshifts

$$X_k \subseteq X_{k-1} \subseteq X_{k-2} \subseteq \cdots \subseteq X_1 \subseteq \mathcal{A}^{\mathbb{Z}}$$

and an increasing sequence of integers $n_1 < n_2 < \cdots < n_k$ such that we have $P_{X_i}(n_i) \leq \binom{n_i-1}{2} \cdot 2^{i-1} |w_1^{i-1}| \cdot n_i$ for $i = 1, 2, \dots, k$. Moreover, for each $i = 1, 2, \dots, k$, suppose we have constructed distinct words

$$w_1^i, w_2^i, \dots, w_{2^i}^i,$$

all of which have the same length, all of which lie in $\mathcal{L}(X_{i-1})$, all of which are concatenations of words from the set $\{w_1^{i-1}, \dots, w_{2^{i-1}}^{i-1}\}$, and are such that X_i is the subshift of X_{i-1} consisting of all words that can be written as bi-infinite concatenations of $w_1^i, w_2^i, \dots, w_{2^i}^i$. Further assume that for any $1 \leq j < k \leq 2^i$, there is no subword of length $|w_1^i|$ that occurs in both $w_j^i w_j^i$ and $w_k^i w_k^i$. Note that since $w_1^i, \dots, w_{2^i}^i$ are concatenations of words from the set $\{w_1^{i-1}, \dots, w_{2^{i-1}}^{i-1}\}$, all concatenations of $w_1^i, \dots, w_{2^i}^i$ are elements of X_{i-1} . Finally, for $i > 1$, suppose that for $1 \leq j \leq 2^i$, the word w_j^i contains each of the words $w_1^{i-1}, w_2^{i-1}, \dots, w_{2^{i-1}}^{i-1}$ somewhere as a subword.

To construct X_{k+1} , we start by defining

$$w_1^{k+1} = \underbrace{\underbrace{(w_1^k w_1^k \cdots w_1^k w_2^k)}_{S_k \text{ times}} \underbrace{(w_1^k w_1^k \cdots w_1^k w_2^k)}_{S_k \text{ times}} \cdots \underbrace{(w_1^k w_1^k \cdots w_1^k w_2^k)}_{S_k \text{ times}}}_{N_k \text{ times}} w_2^k w_3^k \cdots w_{2^k}^k$$

and

$$w_2^{k+1} = \underbrace{\underbrace{(w_1^k w_1^k \cdots w_1^k w_1^k)}_{S_k \text{ times}} \underbrace{(w_1^k w_1^k \cdots w_1^k w_1^k)}_{S_k \text{ times}} \cdots \underbrace{(w_1^k w_1^k \cdots w_1^k w_1^k)}_{S_k \text{ times}}}_{N_k \text{ times}} w_2^k w_3^k \cdots w_{2^k}^k$$

where the brackets have no mathematical meaning other than to draw attention to the periodic nature of the initial prefix of these words and the fact that the word being periodized in w_1^{k+1} is different from that in w_2^{k+1} . Again, N_k and S_k are large integers to be determined later. We continue the construction: for $1 < j \leq 2^k$, defining

$$v_j^k = \overbrace{w_j^k w_j^k \cdots w_j^k}^{S_k \text{ times}},$$

we define

$$w_{2j-1}^{k+1} = w_1^k w_2^k \cdots w_{j-1}^k \underbrace{(v_j^k w_{j+1}^k)(v_j^k w_{j+1}^k) \cdots (v_j^k w_{j+1}^k)}_{N_k \text{ times}} w_{j+1}^k w_{j+2}^k \cdots w_{2^k}^k$$

and

$$w_{2j}^{k+1} = w_1^k w_2^k \cdots w_{j-1}^k \underbrace{(v_j^k w_j^k)(v_j^k w_j^k) \cdots (v_j^k w_j^k)}_{N_k \text{ times}} w_{j+1}^k \cdots w_{2^k}^k,$$

where $w_{2^{k+1}+1}^{k+1} := w_1^{k+1}$, meaning that subscripts are understood modulo 2^{k+1} .

Note that for $1 \leq j < i \leq 2^{k+1}$, we claim that the words of length $|w_1^{k+1}|$ that occur in the word $w_i^{k+1} w_i^{k+1}$ are distinct from those that occur in the word $w_j^{k+1} w_j^{k+1}$ (provided N_k is sufficiently large). If $i = j+1$, the claim follows because w_{j+1}^k occurs at least N_k times in any subword of length $|w_1^{k+1}|$ in $w_j^{k+1} w_j^{k+1}$ (once in each copy of the periodized word $v_j^k w_{j+1}^k$) and occurs at most $|w_1^k| \cdot 2^k$ times in $w_i^{k+1} w_i^{k+1}$ (since it does not occur anywhere in the periodized word and can only occur in the prefix or suffix, which collectively have length $|w_1^k| \cdot 2^k$). For $N_k > |w_1^k| \cdot 2^k$, no such word can occur in both $w_j^{k+1} w_j^{k+1}$ and $w_i^{k+1} w_i^{k+1}$. If $i > j+1$, the claim follows because w_j^k occurs at least $S_k \cdot N_k$ times in any subword of length $|w_1^k|$ in $w_j^{k+1} w_j^{k+1}$ and occurs at most $|w_1^k| \cdot 2^k$ times in $w_i^{k+1} w_i^{k+1}$. Further note that the frequency with which words of length $|w_j^k|$ that occur in $w_j^k w_j^k$, occur as subwords in w_{2j-1}^{k+1} and w_{2j}^{k+1} is at least $N_k S_k |w_1^k| / |w_1^{k+1}|$. By choosing N_k sufficiently large, this can be made arbitrarily close to 1. Thus, by choosing N_k sufficiently large, any word (or collection of words) that occur with frequency δ in $w_j^k w_j^k$ can be made to occur with frequency arbitrarily close to δ in w_{2j-1}^{k+1} and w_{2j}^{k+1} . Furthermore, for $i \notin \{2j-1, 2j\}$, the frequency with which words of length $|w_j^k|$ that occur in $w_j^k w_j^k$ occur in w_i^{k+1} is at most $1 - N_k S_k |w_1^k| / |w_1^{k+1}|$, as these words do not occur in any $w_t^k w_t^k$ for any $t \neq j$ (and subwords of this form occur with frequency at least $N_k S_k |w_1^k| / |w_1^{k+1}|$ in w_i^{k+1}). Again, by choosing N_k sufficiently large, this frequency can be made arbitrarily close to zero.

Define $X_{k+1} \subseteq X_k$ to be the subshift consisting of all words that can be written as bi-infinite concatenations of the words $w_1^{k+1}, w_2^{k+1}, \dots, w_{2^{k+1}}^{k+1}$. Note that, by construction, every word of the form w_j^{k+1} contains each of the words $w_1^k, w_2^k, \dots, w_{2^k}^k$ as a subword. Furthermore $X_{k+1} \subseteq X_k$ and each element of X_{k+1} can be written as a bi-infinite concatenation of the words $w_1^{k+1}, w_2^{k+1}, \dots, w_{2^{k+1}}^{k+1}$.

Define

$$n_{k+1} := \lfloor |w_1^{k+1}| / 2 \rfloor$$

to be half the (common) length of the words $w_1^{k+1}, \dots, w_{2^{k+1}}^{k+1}$. We claim that $P_{X_{k+1}}(n_{k+1}) \leq \left(\binom{2^k}{2} + 1 \right) n_{k+1}$. Each of the words w_i^{k+1} consists of a *prefix region*

in which w_1^k, \dots, w_{i-1}^k are concatenated in order, then a *periodic region* in which either w_i^k is self-concatenated or $w_i^k w_i^k \dots w_i^k w_{i+1}^k$ is self-concatenated, and finally a *suffix region* in which $w_{i+1}^k, \dots, w_{2^k}^k$ are concatenated in order. Any word in $\mathcal{L}_{n_{k+1}}(X_{k+1})$ occurs either entirely within some word w_i^{k+1} or partially overlaps two words $w_{i_1}^{k+1} w_{i_2}^{k+1}$. For words of the first type, they may occur entirely within the periodic region or they start within the first $2^k |w_1^k|$ letters or they end within the last $2^k |w_1^k|$ letters of w_i^{k+1} . For those in the periodic region, there are at most $2^{k+1} (S_k + 1) |w_1^k|$ many such words since this is the number of words constructed multiplied by the maximal period of the periodic region. For those overlapping two of the regions, there are at most $2^{k+1} |w_1^k|$ such words. Words of the second type must overlap $w_{i_1}^{k+1}$ and $w_{i_2}^{k+1}$ for some $i_1 \neq i_2$ and so must start within the last n_{k+1} letters of $w_{i_1}^{k+1}$, and there are at most $\binom{2^{k+1}}{2} n_{k+1}$ such words. Therefore

$$P_{X_{k+1}}(n_{k+1}) \leq 2^{k+1} (S_k + 1) |w_1^k| + 2^{k+1} |w_1^k| + \binom{2^{k+1}}{2} n_{k+1} \leq \left(\binom{2^{k+1}}{2} + 1 \right) n_{k+1},$$

so long as n_{k+1} is sufficiently large when compared to n_k .

This establishes the assumptions of the inductive hypothesis, and giving us an infinite nested chain of subshifts

$$\mathcal{A}^{\mathbb{Z}} \supseteq X_1 \supseteq X_2 \supseteq X_3 \supseteq \dots \supseteq X_k \supseteq \dots$$

Step 3: construction and growth properties of the subshift X . The word w_1^{k+1} starts with the word w_1^k for all k and so there is a $\{0, 1\}$ -coloring of \mathbb{N} such that for all k , the prefix of length $|w_1^k|$ is w_1^k . Let \tilde{X} be the orbit closure of this word in $\mathcal{A}^{\mathbb{N}}$ and let (X, σ) be its natural extension to a subshift of $\mathcal{A}^{\mathbb{Z}}$. It follows immediately from the construction that

$$X \subseteq \bigcap_{k=1}^{\infty} X_k$$

and (X, σ) is a nonempty subshift. We have constructed an increasing sequence $n_1 < n_2 < \dots < n_k < \dots$ such that

$$P_X(n_k) \leq \left(\binom{2^k}{2} + 1 \right) n_k$$

for all $k > 1$. The integers n_k are on the order of $1/2$ of the parameters $N_k |w_1^k|$, and in particular tend to infinity as N_k tends to infinity. The parameters N_k have not yet been fixed, and we put some constraints on them now. Recall that we are given the sequence (p_n) such that

$$\liminf_{n \rightarrow \infty} \frac{p_n}{n} = \infty.$$

For each $k \geq 1$, there exists M_k such that for all $n \geq M_k$, we have

$$p_n > k \cdot \left(\binom{2^k}{2} + 1 \right) n.$$

Fix an increasing sequence of integers (M_k) with this property. We assume that N_k is chosen to be sufficiently large such that $N_k > 2N_{k-1}$ and $n_k > M_k$. Since $P_X(n_k)/p_{n_k} < 1/k$ for all k , we have that

$$\liminf_{n \rightarrow \infty} \frac{P_X(n)}{p_n} = 0.$$

Moreover, note that for any $k > 1$ we have $X \subseteq X_k$. Since X_k consists only of words that can be obtained as bi-infinite concatenations of the words $w_1^k, \dots, w_{2^k}^k$ and each of these words contains all of the words $w_1^{k-1}, \dots, w_{2^{k-1}}^{k-1}$ as a subword, it follows that each of the words $w_1^{k-1}, \dots, w_{2^{k-1}}^{k-1}$ occurs syndetically in every element of X_k (hence also in every element of X) with gap at most $|w_1^k|$. Since every word in $\mathcal{L}(X)$ is itself a subword of w_1^k for some k , it follows that every word in $\mathcal{L}(X)$ occurs syndetically in every element of X with a bound on the gap that depends on the word but not on the element of X . Hence (X, σ) is minimal.

Step 4: the set of ergodic measures on X . Fix a sequence (δ_i) of positive real numbers in the interval $(0, 1)$ such that

$$\Delta := \prod_{i=1}^{\infty} \delta_i > \frac{9}{10}.$$

Then for any $k > 1$, we also have $\prod_{i=k}^{\infty} \delta_i > 9/10$. Recall that every word of the form w_i^k consists of a prefix region, a periodic region, and a suffix region, where the lengths of the prefix and suffix regions are bounded independently of N_k . Thus we can further choose N_k to grow sufficiently quickly such that

$$\frac{N_k}{|w_i^{k+1}|} \cdot \frac{S_k}{S_k + 1} > \delta_k$$

for all $i = 1, \dots, 2^k$ (recall that all of these words have the same length). It follows from this choice that the frequency with which w_j^k (and the other words of length $|w_j^k|$ that occur when this word is self-concatenated) occur in $w_{2^j-1}^{k+1}$ and $w_{2^j}^{k+1}$ is at least δ_k .

We claim that for each word w_i^k , the set of ergodic measures giving measure at least Δ to the set $[w_i^k w_i^k]$ is nonempty. Moreover, we claim that the set of ergodic measures giving measure at least Δ to the set

$$\bigcup_{j=0}^{|w_i^k|} \sigma^j \underbrace{[w_i^k w_i^k \cdots w_i^k]}_{S_k + 1 \text{ times}}$$

is nonempty and that the set of ergodic measures giving measure at least Δ to the set

$$\underbrace{|w_i^k w_i^k \cdots w_i^k w_{i+1}^k|}_{S_k \text{ times}} \bigcup_{j=0} \sigma^j \underbrace{[(w_i^k w_i^k \cdots w_i^k w_{i+1}^k) (w_i^k w_i^k \cdots w_i^k w_{i+1}^k) \cdots (w_i^k w_i^k \cdots w_i^k w_{i+1}^k)]}_{N_k \text{ times}}$$

is also nonempty. The first claim follows from either of the latter two. We show the former, the latter being similar.

Observe that $w_{2^i}^{k+1}$ has a periodic region which is a long series of self-concatenations of w_i^k , that $w_{4^i}^{k+2}$ has a periodic region which is a long series of self-concatenations of $w_{2^i}^{k+1}$, and similarly $w_{2^j}^{k+j}$ has a periodic region which is a long series of self-concatenations of $w_{2^{j-1}}^{k+j-1}$. As these are all words in the language of X and X is closed, there is an element of X for which the natural frequency of w_i^k (and the other words of length $|w_i^k|$ that occur when it is self-concatenated) is at least $\prod_{j=k}^{\infty} \delta_j > \Delta$. This follows because w_i^k (and the other words of length $|w_i^k|$ that

occur when it is self-concatenated) occur with frequency at least δ_k in $w_{2^i}^{k+1}$, and inductively occurs with frequency at least $\prod_{j=k}^{\ell} \delta_j$ in $w_{w^{\ell-j}i}^{\ell}$ for any ℓ . Therefore there is an invariant probability measure on X that gives the union of these cylinder sets measure at least Δ and so there must be at least one ergodic measure that also has this property. The claim follows.

We next show that the set of ergodic measures on X is uncountable (in fact with cardinality \mathfrak{c}). We have shown that for each k and $1 \leq i \leq 2^k$, there are two disjoint sets of ergodic measures giving large measure to the word w_i^k and its periodic shifts. The first (which we refer to as *type 0*) gives large measure to $\underbrace{w_i^k w_i^k \dots w_i^k}_{N_k \text{ times}}$ (and its periodic shifts), whereas the second (which we refer to as *type*

1) gives small measure to this set and large measure to

$$\underbrace{\underbrace{(w_i^k w_i^k \dots w_i^k w_{i+1}^k)}_{S_k \text{ times}} \underbrace{(w_i^k w_i^k \dots w_i^k w_{i+1}^k)}_{S_k \text{ times}} \dots \underbrace{(w_i^k w_i^k \dots w_i^k w_{i+1}^k)}_{S_k \text{ times}}}_{N_k \text{ times}}$$

(and its periodic shifts). Fix an infinite sequence a_0, a_1, \dots of 0's and 1's. For each j , let ν_j be an element of the set of ergodic measures on X that start by giving large measure to w_1^1 and then are of type a_t for each $t = 1, 2, \dots, j$. Let ν be a weak-* limit of a subsequence of these measures so that ν gives measure at least Δ to the word defining its type for each $t = 1, 2, \dots$. If we had chosen any other sequence of 0's and 1's it would have differed from (a_t) at some finite stage and so there would be a cylinder set the measure resulting from that sequence gives large measure to which was given small measure by ν . Therefore each infinite sequence of 0's and 1's produces its own measure ν . Now, returning to the measure ν , consider the sequence (a_t) and the associate union of cylinder sets (which we call \mathcal{S}_t) that are given measure at least $\prod_{k=t}^{\infty} \delta_k$ by ν . Define

$$A_\nu = \bigcap_{t=1}^{\infty} \mathcal{S}_t.$$

Note that

$$\nu \left(\left(\bigcap_{t=1}^{k-1} \mathcal{S}_t \right) \setminus \left(\bigcap_{t=1}^k \mathcal{S}_t \right) \right) \leq 1 - \delta_k$$

and so we can arrange that

$$\nu \left(\bigcap_{t=1}^k \mathcal{S}_t \right) > 8/10$$

for all k by choosing δ_k to tend to 1 sufficiently rapidly. It follows that $\nu(A_\nu) \geq 8/10$. Therefore there is an ergodic measure giving the set A_ν measure at least $8/10$ and so there is no loss of generality in assuming that ν is ergodic. Therefore the set of ergodic measures is uncountable.

This completes the proof of Theorem 1.1. We note that by modifying the initial words w_1^1 and w_2^1 , we can achieve the same result but ensure that the language has balanced numbers of short patterns. More precisely, replacing the initial use of 0 by the word 01100110 and the initial use of 1 by the word 11100100 and carrying out the same construction, we have a system in which the average number of 0's and 1's on any short range is approximately one half. This follows because all

words later constructed are concatenations of w_1^1 and w_2^1 and so any word in the language of the shift of length larger than fourteen can be made into a word that is a concatenation of w_1^1 's and w_2^1 's by removing at most seven letters from each side of it (and this slightly shorter word has precisely the same number of 0's as 1's).

4. MEASURE-THEORETIC COMPLEXITY

4.1. Definition of slow entropy. We review the definition of slow entropy, as defined by Katok and Thouvenot [6] and Ferenczi [3], adopting Ferenczi's notation in a way more convenient for our setting.

Assume that (X, σ) is a subshift. For $u, v \in \mathcal{L}_n(X)$, the *Hamming distance* $d_H(u, v)$ between u and v is

$$d_H(u, v) = \frac{|\{0 \leq i < n: u_i \neq v_i\}|}{n},$$

and this defines a metric on $\mathcal{L}_n(X)$. For fixed $\varepsilon > 0$ and $u \in \mathcal{L}_n(X)$, define the *ball* $B_\varepsilon(u)$ of radius ε around u by

$$B_\varepsilon(u) = \{v \in \mathcal{L}_n(X): d_H(u, v) < \varepsilon\}.$$

Further assume that μ is an invariant measure on the shift (X, σ) . When slow entropy is defined in [6] and [3], they consider an arbitrary measure preserving system and so a generating partition is a necessary ingredient. As we are restricting ourselves to symbolic systems, we can assume that the space X is partitioned into cylinder sets of length one. We implicitly make this assumption throughout and omit the partition from the notation.

Define $K(n, \varepsilon, \sigma)$ to be the minimum number of words $u_1, u_2, \dots, u_k \in \mathcal{L}_n(X)$ such that

$$\mu \left(\bigcup_{i=1}^k B_\varepsilon(u_i) \right) > 1 - \varepsilon.$$

If $\varepsilon_1 < \varepsilon_2$, then for any $u \in \mathcal{L}_n(X)$, we have $B_{\varepsilon_1}(u) \subseteq B_{\varepsilon_2}(u)$. Thus $K(n, \varepsilon_2, \sigma) \leq K(n, \varepsilon_1, \sigma)$, meaning that $K(n, \varepsilon, \sigma)$ increases as ε decreases. If $(c_n)_{n \in \mathbb{N}}$ is a non-decreasing sequence of positive integers with $c_n \rightarrow \infty$, we say $P_\sigma^-(n) \succ (c_n)$ if

$$\lim_{\varepsilon \rightarrow 0} \liminf_{n \rightarrow \infty} \frac{K(n, \varepsilon, \sigma)}{c_n} \geq 1.$$

Similarly, we say that $P_\sigma^-(n) \prec (c_n)$ if

$$\lim_{\varepsilon \rightarrow 0} \liminf_{n \rightarrow \infty} \frac{K(n, \varepsilon, \sigma)}{c_n} \leq 1.$$

The analogous limits with \liminf replaced by \limsup define the conditions that $P_\sigma^+(n) \succ (c_n)$ and $P_\sigma^+(n) \prec (c_n)$, respectively. It is shown in [6] and [3] that for any fixed sequence (c_n) , the statement $P_\sigma^-(n) \prec (c_n)$ is a measure theoretic conjugacy invariant for (X, μ, σ) (as is the analogous statement for $P_\sigma^+(n) \succ (c_n)$).

These notions of $P_\sigma^-(n)$ and $P_\sigma^+(n)$ clarify the statement of Theorem 1.2, and we reproduce the statement:

Theorem (Theorem 1.2). *Assume $(a_n)_{n \in \mathbb{N}}$ and $(b_n)_{n \in \mathbb{N}}$ are two nondecreasing sequences of positive integers such that $\lim a_n = \infty$, $\lim \frac{1}{n} \cdot \log(b_n) = 0$, and $a_n \leq b_n$ for all $n \in \mathbb{N}$. There exists a minimal subshift (X_∞, σ) of topological entropy zero and an ergodic measure μ supported on X_∞ such that*

$$P_\sigma^-(n) \prec (a_n) \text{ and } P_\sigma^+(n) \succ (b_n).$$

Before turning to the proof, we make a few remarks to place the result in context. Ferenczi showed the following:

Theorem 4.1 (Ferenczi [3, Proposition 3]). *Let (X, σ) be a subshift and suppose μ is an ergodic measure supported on X . Then the following are equivalent:*

- (i) (X, μ, σ) is a Kronecker system;
- (ii) $P_\sigma^-(n) \prec (c_n)$ for any nondecreasing sequences (c_n) that tends to infinity;
- (iii) $P_\sigma^+(n) \prec (c_n)$ for any nondecreasing sequences (c_n) that tends to infinity.

This means that the assumption on the lower bound $(a_n)_{n \in \mathbb{N}}$ in Theorem 1.2 can not be lowered as long as we still require that $P_\sigma^+(n) \succ (b_n)$, as this second condition implies that (X, μ, σ) is not a Kronecker system and so there must be some sequence $(c_n)_{n \in \mathbb{N}}$ that tends to infinity and is such that $P_\sigma^-(n) \succ (c_n)$.

At the other extreme, Katok showed:

Theorem 4.2 (Katok[5, Theorem 1.1]). *Let (X, σ) be a subshift and suppose μ is an ergodic measure supported on X . Then the following are equivalent:*

- (i) (X, μ, σ) has positive entropy;
- (ii) there exists $\lambda > 1$ such that $P_\sigma^-(n) \succ (\lambda^n)$;
- (iii) there exists $\lambda > 1$ such that $P_\sigma^+(n) \succ (\lambda^n)$.

In particular, since $(a_n)_{n \in \mathbb{N}}$ grows subexponentially and $P_\sigma^-(n) \prec (a_n)$, this implies that (X, μ, σ) has zero entropy and so we cannot have $P_\sigma^+(n) \succ (b_n)$ for any sequence $(b_n)_{n \in \mathbb{N}}$ with positive exponential growth rate. Theorem 1.2 implies that, even subject to the requirement that $P_\sigma^-(n) \prec (a_n)$, $P_\sigma^+(n)$ can grow as quickly as we want, subject to the necessary condition that it grow subexponentially, as given by Katok's Theorem.

Katok and Thouvenot [6] and Serafin [9] give constructions of a subshift with zero topological entropy and $P_\sigma^+(n)$ growing arbitrarily quickly (of course still subject to the condition that the growth be subexponential), but without a requirement that $P_\sigma^-(n)$ grow slowly. In showing that we achieve our upper bound (what we refer to as the loud phase), our construction and the derivation of its properties has many features in common with their constructions. However, as we generalize this approach, we include all details for the sake of clarity.

4.2. Large-scale features of the construction. We fix the nondecreasing sequences $(a_n)_{n \in \mathbb{N}}$ and $(b_n)_{n \in \mathbb{N}}$ of positive integers such that $\lim_{n \rightarrow \infty} a_n = \infty$, $\lim_{n \rightarrow \infty} \frac{1}{n} \cdot \log(b_n) = 0$, and $a_n \leq b_n$ for all $n \in \mathbb{N}$. Let $(\varepsilon_n)_{n \rightarrow \infty}$ be a decreasing sequence of positive real numbers such that $\lim_{n \rightarrow \infty} \varepsilon_n = 0$. We inductively construct a descending sequence of positive entropy subshifts:

$$\mathcal{A}^{\mathbb{Z}} =: X_0 \supseteq X_1 \supseteq X_2 \supseteq X_3 \supseteq \cdots \supseteq X_n \supseteq \cdots$$

and an increasing sequence of positive integers

$$N_1 < P_1 < N_2 < P_2 < N_3 < P_3 < \cdots .$$

In our construction, we show that

$$X_\infty := \bigcap_{i=1}^{\infty} X_i$$

is nonempty and show that if μ is any ergodic measure supported on X_∞ , then

$$(1) \quad K(N_i, 1/8, \sigma) > b_{N_i}$$

and

$$(2) \quad K(P_i, \varepsilon_i, \sigma) \leq a_{P_i}.$$

Since $K(n, \varepsilon, \sigma)$ increases as ε decreases, it follows that for sufficient small $\varepsilon > 0$,

$$\limsup_{n \rightarrow \infty} \frac{K(n, \varepsilon, \sigma)}{b_n} \geq \limsup_{i \rightarrow \infty} \frac{K(N_i, \varepsilon, \sigma)}{b_{N_i}} \geq \limsup_{i \rightarrow \infty} \frac{K(N_i, 1/8, \sigma)}{b_{N_i}} \geq 1,$$

meaning that

$$\lim_{\varepsilon \rightarrow 0} \limsup_{n \rightarrow \infty} \frac{K(n, \varepsilon, \sigma)}{b_n} \geq 1,$$

and so $P_\sigma^+(n) \succ (b_n)$. Similarly, since $\lim_{i \rightarrow \infty} \varepsilon_i = 0$ and again passing to the subsequence $(P_i)_{i \in \mathbb{N}}$, it follows that

$$\liminf_{n \rightarrow \infty} \frac{K(n, \varepsilon, \sigma)}{a_n} \leq \liminf_{i \rightarrow \infty} \frac{K(P_i, \varepsilon, \sigma)}{a_{P_i}} \leq \liminf_{i \rightarrow \infty} \frac{K(P_i, \varepsilon_i, \sigma)}{a_{P_i}} \leq 1.$$

This means that

$$\lim_{\varepsilon \rightarrow 0} \liminf_{n \rightarrow \infty} \frac{K(n, \varepsilon, \sigma)}{a_n} \leq 1$$

and so $P_\sigma^-(n) \prec (a_n)$. Therefore, to prove Theorem 1.2, it suffices to construct the shift X_∞ , the ergodic measure μ and show that they satisfy (1) and (2). To show that this construction can be carried out to produce a minimal shift, we note that by the Jewett-Krieger Theorem there is a strictly ergodic model for (X_∞, μ, σ) and this model must obey the same slow entropy bounds because they are invariants of measure theoretic conjugacy. We start in Section 4.3 by providing estimates on how the words in the language of the shift must be constructed and then in Section 4.4 complete the construction of the subshift and verify its properties.

4.3. Estimates for the language. We start with a lemma for use in the proof of the main theorem:

Lemma 4.3. *Let $k, n \in \mathbb{N}$ be fixed and let $A_1, \dots, A_{2k-1} \subseteq \{1, 2, \dots, n\}$ be any collection of subsets satisfying $|A_i| \geq n/2$ for $i = 1, \dots, 2k-1$. Then there exist distinct indices $1 \leq i_1 < i_2 < \dots < i_k \leq 2k-1$ and $s \in \{1, 2, \dots, n\}$ such that $s \in A_{i_j}$ for all $j = 1, 2, \dots, k$.*

Proof. For contradiction, suppose $A_1, \dots, A_{2k-1} \subseteq \{1, 2, \dots, n\}$ are a collection of subsets satisfying $|A_i| \geq n/2$ for $i = 1, \dots, 2k-1$, but no subcollection of k of these sets have nonempty intersection. For each $x \in \{1, 2, \dots, n\}$, let $i(x)$ denote the

number of distinct indices j such that $x \in A_j$. Then $i(x) \leq k - 1$ for all x and so

$$\begin{aligned} n(k-1) &< \sum_{i=1}^{2k-1} |A_i| = \sum_{m=0}^{k-1} m \cdot |\{x: i(x) = m\}| \\ &\leq \sum_{m=0}^{k-1} (k-1) \cdot |\{x: i(x) = m\}| = n(k-1), \end{aligned}$$

where the last equality holds since the level sets of $i(x)$ partition $\{1, 2, \dots, n\}$. Thus no such collection of sets exists. \square

Lemma 4.4. *Let \mathcal{A} be a finite alphabet and suppose $k, N, M \in \mathbb{N}$ are fixed and assume that $M > 1$. Let $w_1, \dots, w_k \in \mathcal{L}_N(\mathcal{A}^{\mathbb{Z}})$ and for each $1 \leq i \leq k$, let*

$$v_i = \underbrace{w_i w_i w_i \cdots w_i}_{M \text{ times}}.$$

Let (X, σ) denote the subshift (with alphabet \mathcal{A}) defined by the words v_1, \dots, v_k . Assume $P \in [1, NM)$ is an integer and let $\mathcal{W}_P \subseteq \mathcal{L}_P(X)$ be the set of all words u for which there exists i such that u is a subword of v_i . Then if μ is any ergodic measure supported on X , we have

$$\mu \left(\bigcup_{w \in \mathcal{W}_P} [w] \right) \geq 1 - \frac{P-1}{NM}.$$

Proof. Assume μ is an ergodic measure supported on X and set $\mathcal{S} = \bigcup_{w \in \mathcal{W}_P} [w]$. By the pointwise ergodic theorem, there exists $x \in X$ such that

$$\mu(\mathcal{S}) = \lim_{n \rightarrow \infty} \frac{1}{2n+1} \sum_{i=-n}^n \mathbf{1}_{\mathcal{S}}(\sigma^i x).$$

By definition of the subshift (X, σ) , the element x can be parsed into a bi-infinite concatenation of the words v_1, \dots, v_k . Fix one such way to parse x and let $\mathcal{I} \subseteq \mathbb{Z}$ be the set of indices at which these words begin; note that \mathcal{I} is an arithmetic progression in \mathbb{Z} with gap NM . For each $i \in \mathbb{Z}$, the cylinder set of length P that contains $\sigma^i x$ is contained in \mathcal{S} , unless i lies within distance $P-1$ of the smallest element of \mathcal{I} larger than i . Thus for any $n \in \mathbb{N}$,

$$\sum_{i=-n}^n \mathbf{1}_{\mathcal{S}}(\sigma^i x) \geq 2n+1 - (P-1) \cdot |\mathcal{I} \cap [-n, n]|.$$

Since \mathcal{I} is an arithmetic progression with gap NM , it follows that

$$\mu(\mathcal{S}) \geq \lim_{n \rightarrow \infty} \frac{2n+1 - (P-1) \cdot |\mathcal{I} \cap [-n, n]|}{2n+1} \geq 1 - \frac{P-1}{NM}. \quad \square$$

Lemma 4.5. *Let $N \in \mathbb{N}$ and $\varepsilon > 0$ be fixed. For each n , let $\mathcal{W}_n \subseteq \{1, 2, \dots, N\}^n$ be the set of words $w = (w_0, w_1, \dots, w_{n-1})$ for which*

$$(1-\varepsilon) \frac{n}{N} < |\{i \in [0, n): w_i = a\}| < (1+\varepsilon) \frac{n}{N}$$

for all $a \in \{1, 2, \dots, N\}$. Then there exists M such that for all $n > M$, we have $|\mathcal{W}_n| > (1-\varepsilon)N^n$.

Proof. Let ν be the $(1/N, 1/N, \dots, 1/N)$ -Bernoulli measure on $\{1, 2, \dots, N\}^{\mathbb{N}}$. By the pointwise ergodic theorem, for almost all $x \in \{1, 2, \dots, N\}^{\mathbb{N}}$ and for each $a \in \{1, 2, \dots, N\}$ we have

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} \mathbf{1}_{[a]}(\sigma^i x) = \nu([a]) = \frac{1}{N}.$$

Therefore there exists $M(x)$ such that for all $n > M(x)$ we have

$$\frac{1 - \varepsilon}{N} < \frac{1}{n} \sum_{i=0}^{n-1} \mathbf{1}_{[a]}(\sigma^i x) < \frac{1 + \varepsilon}{N}$$

for all $a \in \{1, 2, \dots, N\}$. Thus there is some M and a set \mathcal{S} of ν -measure at least $1 - \varepsilon$ such that these inequalities hold for any $x \in \mathcal{S}$ and any $n > M$. Setting $w_n(x) = (x_0, x_1, \dots, x_{n-1})$, we have that

$$\nu \left(\bigcup_{x \in \mathcal{S}} [w_n(x)] \right) \geq \nu(\mathcal{S}) \geq 1 - \varepsilon.$$

Since the ν -measure of each word of length n is $1/N^n$, it follows that the number of distinct words of length n that can be written as $w_n(x)$ for some $x \in \mathcal{S}$ is at least $(1 - \varepsilon)N^n$. \square

We combine these to derive our key estimate on the statistics in the language:

Proposition 4.6. *Let $N \in \mathbb{N}$, $\varepsilon > 0$, and $0 < \alpha < \frac{N-1}{N}$ be fixed. Then there exists $M \in \mathbb{N}$ and $\lambda = \lambda(N, \varepsilon, \alpha) > 1$ such that for any $n > M$, there is a set of words $w_1, \dots, w_k \in \{1, 2, \dots, N\}^n$ satisfying*

$$d_H(w_i, w_j) > \alpha \text{ for all } i \neq j$$

with $k > \lambda^n$ and for all $a \in \{1, 2, \dots, N\}$,

$$(1 - \varepsilon) \frac{n}{N} < |\{i \in [0, n) : w_i = a\}| < (1 + \varepsilon) \frac{n}{N}.$$

Moreover, these words can be chosen such that for any $1 \leq j_1 < j_2 \leq k$, no word of length n that occurs as a subword of $w_{j_1} w_{j_1}$ is also a subword of $w_{j_2} w_{j_2}$.

Proof. Let $w \in \{1, 2, \dots, N\}^n$ be fixed. A classical use of Stirling's Formula (see for example [5, Equation (1.3)]) shows that, since $0 < \alpha < \frac{N-1}{N}$,

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{1}{n} \log |\{u \in \{1, 2, \dots, N\}^n : d_H(u, w) < \alpha\}| \\ = \alpha \log(N - 1) - \alpha \log \alpha - (1 - \alpha) \log(1 - \alpha). \end{aligned}$$

Set $f(x) = x \log(N - 1) - x \log x - (1 - x) \log(1 - x)$. Then for $x < (N - 1)/N$, the derivative of $f(x)$ is positive and $\lim_{x \rightarrow [(N-1)/N]^-} f(x) = \log(N)$. Thus $f(\alpha) < \log(N)$, and so there exists $M \in \mathbb{N}$ and $\delta > 0$ sufficiently small such that $(1 + \delta)f(\alpha) < \log(N)$ and such that for all $n > M$,

$$|\{u \in \{1, 2, \dots, N\}^n : d_H(u, w) < \alpha\}| < 2^{n \cdot (1 + \delta)f(\alpha)}.$$

By Lemma 4.5, if \mathcal{W}_n is the set of all words $w = (w_0, w_1, \dots, w_{n-1}) \in \{1, 2, \dots, N\}^n$ for which

$$(1 - \varepsilon) \frac{n}{N} < |\{i \in [0, n) : w_i = a\}| < (1 + \varepsilon) \frac{n}{N},$$

then $|\mathcal{W}_n| > (1 - \varepsilon)N^n$ for all sufficiently large n . Adjusting the value of M if necessary, we can assume this holds for all $n > M$. But for each $w \in \mathcal{W}_n$, we have

$$|\{u \in \mathcal{W}_n : d_H(u, w) < \alpha\}| \leq |\{u \in \{1, 2, \dots, N\}^n : d_H(u, w) < \alpha\}| \leq 2^{n \cdot (1+\delta)f(\alpha)}.$$

This means there is a set of at least

$$\left\lfloor \frac{(1 - \varepsilon)N^n}{2^{n \cdot (1+\delta)f(\alpha)}} \right\rfloor$$

elements of \mathcal{W}_n that are pairwise at least α separated in the Hamming distance. If u, v are two words in this set and if some word w of length $|u| = |v|$ occurs as a subword of uu and vv , then u is itself a subword of vv . Thus there is a subset of size at least

$$k(n) := \frac{1}{n} \cdot \left\lfloor \frac{(1 - \varepsilon)N^n}{2^{n \cdot (1+\delta)f(\alpha)}} \right\rfloor$$

with the additional property that for any u, v in this list, no word of length n occurs as a subword of both uu and vv .

Since $(1 + \delta)f(\alpha) < \log(N)$, it follows that

$$g := \lim_{n \rightarrow \infty} \frac{\log k(n)}{n} = \log(N) - (1 + \delta)f(\alpha) > 0.$$

Thus if $\lambda := 2^{g/2}$, then $\lambda > 1$ and $k(n) \geq \lambda^n$ for all $n > M$. \square

4.4. Construction of the subshifts. The remainder of this section is devoted to proof of Theorem 1.2.

We construct a sequence of subshifts inductively. Each step of the construction involves the construction of two nested subshifts, with the larger one being referred to as the “noisy” phase and the smaller one being referred to as the “quiet” phase. At each stage of the construction, we appeal to Proposition 4.6, and this necessitates the definition of two auxiliary sequences. Set $\alpha_0 = 1/3$ and let $(\alpha_i)_{i \geq 0}$ be an increasing sequence of real numbers with $0 < \alpha_i < 1$ for all $i \in \mathbb{N}$ and such that

$$(3) \quad \prod_{i=1}^{\infty} \alpha_i > \frac{3}{4}.$$

This sequence provides the parameter α appearing in Proposition 4.6 for each step of the construction. Next let $(\varepsilon_i)_{i \geq 0}$ be a decreasing sequence of real numbers with $0 < \varepsilon_i < 1$ for all i and such that

$$(4) \quad \prod_{i=1}^{\infty} (1 - \varepsilon_i) > \frac{99}{100}.$$

This sequence provides the parameter ε appearing Proposition 4.6.

Base Loud Phase. Let $X_0 := \{1, 2\}^{\mathbb{Z}}$ and $M_0 = 1$. Thus X_0 is a subshift on $N_0 := 2$ letters. Since $1/3 = \alpha_0 < \frac{N_0 - 1}{N_0}$, we can apply Proposition 4.6 with parameters $N = N_0$, $\varepsilon = \varepsilon_0$, and $\alpha = \alpha_0$. Thus there exists $\lambda_0 > 1$ such that for any sufficiently large integer n , we can find a set of words $w_1(n), w_2(n), \dots, w_{k(n)}(n) \in \{1, 2\}^n$ where $k(n) > \lambda_0^n$ and such that for each $a \in \{1, 2\}$, the following conditions are satisfied:

- (i) We have the estimate $(1 - \varepsilon_0)\frac{n}{2} < B_a < (1 + \varepsilon_0)\frac{n}{2}$ for all i , where B_a denotes the number of locations where the letter a occurs in $w_i(n)$;

- (ii) We have the distances separated, meaning that $d_H(w_i(n), w_j(n)) > \alpha_0$ for any $i \neq j$;
- (iii) No word of length n occurs as a subword of both $w_i(n)w_i(n)$ and $w_j(n)w_j(n)$ for some $i \neq j$.

Let N_1 be an integer which is sufficiently large that we can choose such a set of words, such that $\lambda_0^{N_1} > 4b_{N_1}$, and such that $\alpha_1 < \frac{N_1-1}{N_1}$. Let $k_1 := k(N_1)$ be the number of words constructed in this way and let $w_1, \dots, w_{k_1} \in \{1, 2\}^{N_1}$ be the words produced by the construction. Finally let $L_1 \subseteq X_0$ be the subshift of X_0 defined by the words w_1, w_2, \dots, w_{k_1} .

Base Quiet Phase. Choose an integer $P_1 > N_1$ sufficiently large that

$$|w_1| + |w_2| + \dots + |w_{k_1}| = k_1 N_1 < a_{P_1}$$

and then choose an integer M_1 such that $(P_1-1)/(N_1 M_1) < \varepsilon_0$. For each $1 \leq i \leq k_1$, define

$$v_i := \underbrace{w_i w_i w_i \dots w_i}_{M_1 \text{ times}}.$$

Let $X_1 \subseteq L_1$ be the subshift defined by the words v_1, \dots, v_{k_1} . We apply Lemma 4.4 with parameters $\mathcal{A} = \{1, 2\}$, $k = k_1$, $N = N_1$, $M = M_1$, and $P = P_1 \in [1, NM)$. If $\mathcal{W} = \mathcal{W}_{P_1}$ is the set of words of length P_1 that occur as subwords of the words v_1, \dots, v_{k_1} , then for any ergodic measure μ supported on X_1 we have

$$\mu \left(\bigcup_{w \in \mathcal{W}} [w] \right) > 1 - \frac{P_1}{N_1 M_1} > 1 - \varepsilon_0$$

and if $\widetilde{\mathcal{W}} = \widetilde{\mathcal{W}}_{N_1}$ is the set of words of length N_1 that occur as subwords of the words v_1, \dots, v_{k_1} , then

$$\mu \left(\bigcup_{w \in \widetilde{\mathcal{W}}} [w] \right) > 1 - \frac{N_1}{N_1 M_1} > 1 - \varepsilon_0.$$

Inductive Loud Phase. Assume we have constructed subshifts

$$\{1, 2\}^{\mathbb{Z}} =: X_0 \supseteq L_1 \supseteq X_1 \supseteq L_2 \supseteq X_2 \supseteq \dots \supseteq L_i \supseteq X_i,$$

a sequence of integers $N_1 < P_1 < N_2 < P_2 < \dots < N_i < P_i$, a sequence of integers $M_1 < M_2 < \dots < M_i$, such that that following hold: for each $1 \leq j \leq i$,

- (i) N_j is sufficiently large such that $\alpha_j < \frac{N_j-1}{N_j}$;
- (ii) There exists $\lambda_j > 1$ and an integer k_j such that $k_j > \lambda_j^{N_j} > 4b_{N_j}$;
- (iii) There exist words $w_1^j, w_2^j, \dots, w_{k_j}^j \in \{1, 2\}^{N_j \cdot \prod_{s=0}^{j-1} N_s M_s}$ such that L_j is comprised of all elements of X_{j-1} defined by the words $w_1^j, \dots, w_{k_j}^j$ and for $i_1 \neq i_2$ we have $d_H(w_{i_1}^j, w_{i_2}^j) > \prod_{s=1}^j \alpha_s$ and additionally no word of length $|w_1^j|$ occurs as a subword of both $w_{i_1}^j w_{i_1}^j$ and $w_{i_2}^j w_{i_2}^j$;
- (iv) For each $1 \leq t \leq k_j$, there is a word

$$v_t^j := \underbrace{w_t^j w_t^j w_t^j \dots w_t^j}_{M_j \text{ times}}$$

where for any $t_1 \neq t_2$ no subword of length $N_j \cdot \prod_{s=0}^{j-1} N_s M_s$ in $v_{t_1}^j$ is also a subword of $v_{t_2}^j$, and $X_j \subseteq L_j$ is the subshift of L_j defined by the words $v_1^j, v_2^j, \dots, v_{k_j}^j$;

- (v) If $j > 1$, then for each $1 \leq t \leq k_j$ the word w_t^j can be written as a concatenation of the words $v_1^{j-1}, \dots, v_{k_{j-1}}^{j-1}$ and so by identifying this set of k_{j-1} words with the alphabet $\{1, 2, \dots, k_{j-1}\}$, we can identify w_t^j with a word of length N_j written in these letters. With this identification, for each $a \in \{1, 2, \dots, k_{j-1}\}$, we have

$$\begin{aligned} \frac{N_j}{k_{j-1}} \cdot (1 - \varepsilon_{j-1}) &< |\{i \in [0, n): \text{the } i^{\text{th}} \text{ letter in } w_t^j \text{ is } a\}| \\ &< \frac{N_j}{k_{j-1}} \cdot (1 + \varepsilon_{j-1}); \end{aligned}$$

- (vi) We have $P_j > k_j$, $M_j > |w_1^j| k_j / \varepsilon_j$, and $(P_j - 1) / (N_j M_j) < \varepsilon_j$ and if $\mathcal{W} = \mathcal{W}_{P_j}$ is the set of all words of length P_j that occur in X_j as subwords of $v_1^j, v_2^j, \dots, v_{k_j}^j$ and if μ is any ergodic measure supported on X_j , then

$$\mu \left(\bigcup_{w \in \mathcal{W}} [w] \right) > (1 - \varepsilon_j)$$

and if $\widetilde{\mathcal{W}} = \widetilde{\mathcal{W}}_{N_j}$ is the set of all words of length N_j that occur in X_j as subwords of $v_1^j, v_2^j, \dots, v_{k_j}^j$ then

$$\mu \left(\bigcup_{w \in \widetilde{\mathcal{W}}} [w] \right) > (1 - \varepsilon_j).$$

Since $\alpha_i < \frac{N_i - 1}{N_i}$, we can apply Proposition 4.6 with parameters $N = k_i$, $\varepsilon = \varepsilon_i$, and $\alpha = \alpha_i$. Thus there exists $\lambda_i > 1$ such that for any sufficiently large integer n , there is a set of words $w_1(n), w_2(n), \dots, w_{k(n)}(n) \in \{1, 2, \dots, k_i\}^n$ where $k(n) > \lambda_i^n$ and such that for any $a \in \{1, 2, \dots, k_i\}$, the following conditions are satisfied:

- (i) We have the estimate $(1 - \varepsilon_i) \frac{n}{k_i} < B_a < (1 + \varepsilon_i) \frac{n}{k_i}$, where again B_a denotes the number of locations where the letter a occurs in $w_i(n)$;
- (ii) We have the distances separated, meaning that $d_H(w_i(n), w_j(n)) > \alpha_i$ for any $i \neq j$;
- (iii) No word of length n occurs as a subword of both $w_i(n)w_i(n)$ and $w_j(n)w_j(n)$ for any $i \neq j$.

Let N_{i+1} be a sufficiently large integer such that we can find such a set of words, such that $\lambda_i^{N_{i+1}} > 4b_{N_{i+1}}$, and such that $\alpha_{i+1} < \frac{N_{i+1} - 1}{N_{i+1}}$. Let $k_{i+1} := k(N_{i+1})$ be the number of words constructed in this way. Finally let

$$w_1^{i+1}, w_2^{i+1}, \dots, w_{k_{i+1}}^{i+1} \in \{1, 2\}^{N_{i+1} \cdot \prod_{s=0}^i N_s M_s}$$

be the words constructed by concatenating $w_1^i, \dots, w_{k_i}^i$ according to the letters of the words $w_1(n), \dots, w_{k_{i+1}}(n)$: for $1 \leq j \leq k_{i+1}$ if $w_j(n) = a_1 a_2 \dots a_{k_i}$ then we define

$$w_j^{i+1} = w_{a_1}^i w_{a_2}^i \dots w_{a_{k_i}}^i.$$

Finally let $L_{i+1} \subseteq X_i$ be the subshift of X_i defined by the words $w_1^{i+1}, w_2^{i+1}, \dots, w_{k_{i+1}}^{i+1}$.

Inductive Quiet Phase. Choose an integer $P_{i+1} > N_{i+1}$ sufficiently large such that

$$(5) \quad |w_1^{i+1}| + |w_2^{i+1}| + \cdots + |w_{k_{i+1}}^{i+1}| = k_{i+1}N_{i+1} \cdot \prod_{s=0}^i N_s M_s < a_{P_{i+1}}.$$

Find an integer $M_{i+1} > |w_1^{i+1}|k_{i+1}/\varepsilon_{i+1}$ such that $(P_{i+1} - 1)/(N_{i+1}M_{i+1}) < \varepsilon_i$. For each $1 \leq j \leq k_{i+1}$, define

$$v_j^{i+1} := \underbrace{w_j^{i+1} w_j^{i+1} w_j^{i+1} \cdots w_j^{i+1}}_{M_{i+1} \text{ times}}.$$

Let $X_{i+1} \subseteq L_{i+1}$ be the subshift of L_{i+1} defined by the words $v_1^{i+1}, \dots, v_{k_{i+1}}^{i+1}$. As in the base case, for each $1 \leq j \leq k_{i+1}$, we choose $w_j^{i+1} = w_{a_1}^i w_{a_2}^i \cdots w_{a_{N_i}}^i$ as a way to parse w_j^{i+1} into a concatenation of words with superscript i . Then define

$$\tilde{v}_j^{i+1} = \underbrace{(a_1 a_2 \cdots a_{N_i})(a_1 a_2 \cdots a_{N_i}) \cdots (a_1 a_2 \cdots a_{N_i})}_{M_{i+1} \text{ times}}$$

to be the identification of v_j^{i+1} with a concatenation of letters $\{1, 2, \dots, k_i\}$, rather than words $\{w_1^i, w_2^i, \dots, w_{k_i}^i\}$. Let \tilde{X}_{i+1} be the subshift of $\{1, 2, \dots, k_i\}^{\mathbb{Z}}$ defined by the words $\tilde{v}_1^{i+1}, \dots, \tilde{v}_{k_{i+1}}^{i+1}$. We then apply Lemma 4.4 with parameters $\mathcal{A} = \{1, 2, \dots, k_i\}$, $k = k_{i+1}$, $N = N_{i+1}$, $M = M_{i+1}$, and we choose $P = P_{i+1} \in [1, N_{i+1}M_{i+1})$. Then if $\mathcal{W} = \mathcal{W}_{P_{i+1}}$ denotes the set of words of length P_{i+1} that occur as subwords of $\tilde{v}_1^{i+1}, \tilde{v}_2^{i+1}, \dots, \tilde{v}_{k_{i+1}}^{i+1}$ and if ν is any ergodic measure supported on \tilde{X}_{i+1} , then

$$\nu \left(\bigcup_{w \in \mathcal{W}} [w] \right) > 1 - \frac{P_{i+1}}{N_{i+1}M_{i+1}} > (1 - \varepsilon_i).$$

If $\tilde{\mathcal{W}} = \tilde{\mathcal{W}}_{N_i}$ is the set of words of length N_i that occur as subwords of $\tilde{v}_1^{i+1}, \dots, \tilde{v}_{k_{i+1}}^{i+1}$, then

$$\nu \left(\bigcup_{w \in \tilde{\mathcal{W}}} [w] \right) > 1 - \frac{N_{i+1}}{N_{i+1}M_{i+1}} > (1 - \varepsilon_i).$$

Therefore conditions (i)–(vi) of the induction hypothesis are satisfied for $j = i + 1$.

Thus, by induction, we obtain an infinite descending sequence of subshifts

$$\{1, 2\}^{\mathbb{Z}} =: X_0 \supseteq L_1 \supseteq X_1 \supseteq L_2 \supseteq X_2 \supseteq \cdots \supseteq L_i \supseteq X_i \supseteq \cdots$$

We define

$$X_\infty := \bigcap_{i=0}^{\infty} X_i.$$

Since $\{1, 2\}^{\mathbb{Z}}$ is a Baire space (with the usual metric), the intersection of any nested sequence of subshifts is nonempty and so X_∞ is nonempty.

We now assume that μ is an ergodic measure supported on X_∞ and we study its properties.

Analysis of $P^+(n)$. Our goal is to show that

$$K(\alpha, N_i, 1/8, \sigma) > b_{N_i}$$

for all i . Fix $i \in \mathbb{N}$. First we recall the definition of $K(N_i, 1/8, \sigma)$. For $u \in \mathcal{L}_{N_i}(X_\infty)$, let

$$B_{1/8}(u) = \{w \in \mathcal{L}_{N_i}(X_\infty) : d_H(u, w) < 1/8\}$$

be the $(1/8)$ -Hamming ball around u . Define

$$[B_{1/8}(u)] := \bigcup_{w \in B_{1/8}(u)} [w].$$

With this notation, $K(N_i, 1/8, \sigma)$ is the smallest cardinality of a set $\mathcal{U} \subseteq \mathcal{L}_{N_i}(X_\infty)$ such that

$$(6) \quad \mu \left(\bigcup_{u \in \mathcal{U}} [B_{1/8}(u)] \right) > \frac{7}{8}.$$

Fix such a set $\mathcal{U} \subseteq \mathcal{L}_{N_i}(X_\infty)$. To establish (1), we are left with showing that $|\mathcal{U}| > b_{N_i}$.

Since μ is an ergodic measure supported on X_∞ , it is also an ergodic measure supported on X_i (albeit not a measure of full support). The shift X_i is constructed by first constructing a set of words $w_1^i, w_2^i, \dots, w_{k_i}^i$ of length N_i and using them to construct words $v_1^i, v_2^i, \dots, v_{k_i}^i$ via the formula

$$v_j^i = \underbrace{w_j^i w_j^i w_j^i \cdots w_j^i}_{M_i \text{ times}}$$

and M_i is a parameter chosen during the construction. Thus the language of X_i is defined to be all elements of $\{1, 2\}^{\mathbb{Z}}$ that can be written as bi-infinite concatenations of $v_1^i, v_2^i, \dots, v_{k_i}^i$. The choice of M_i guarantees, by induction hypothesis (vi), that for any ergodic measure supported on X_i (in particular, for μ) if $\widetilde{\mathcal{W}}$ is the set of words of length N_i that occur as subwords of one of $v_1^i, v_2^i, \dots, v_{k_i}^i$, then

$$(7) \quad \mu \left(\bigcup_{w \in \widetilde{\mathcal{W}}} [w] \right) > 1 - \varepsilon_i.$$

The construction also guarantees, by induction hypothesis (iii), that for $j_1 \neq j_2$ we have $d_H(w_{j_1}^i, w_{j_2}^i) > \prod_{s=0}^i \alpha_s \geq \prod_{s=0}^\infty \alpha_s > 1/4$ (recall that $\alpha_0 = 1/3$ and equation (3)). Now observe that from (6) and (7), if $\mathcal{V} \subseteq \widetilde{\mathcal{W}}$ is the set of all $w \in \widetilde{\mathcal{W}}$ such that there exists $u \in \mathcal{U}$ such that $d_H(u, w) < 1/8$, then

$$\mu \left(\bigcup_{w \in \mathcal{V}} [w] \right) > \frac{7}{8} - \varepsilon_i \geq \frac{3}{4}$$

provided i is sufficiently large. Next observe that if $w \in \mathcal{V}$, then w is a word of length N_i that occurs as a subword of one of $v_1^i, v_2^i, \dots, v_{k_i}^i$. In particular, this means there exists $1 \leq j \leq k_i$ such that w is a subword of $w_j^i w_j^i$ (recall that v_j^i is just the concatenation of a large number of copies of w_j^i and $|w_j^i| = N_i$). We write

$$[[w_j^i]] = \bigcup_{w \hookrightarrow w_j^i w_j^i} [w]$$

where $w \hookrightarrow w_j^i w_j^i$ means w is a word of length N_i that occurs as a subword of $w_j^i w_j^i$. Therefore there exists a smallest integer $1 \leq t(w) < N_i$ such that the subword of $w_j^i w_j^i$ of length N_i that starts on the $t(w)^{\text{th}}$ letter is w ; let $s(w)$ denote the set whose only element is $t(w)$ (if $w_j^i w_j^i$ is periodic of period smaller than $|w_j^i|$, let $s(w)$ denote the set of starting points of w , excluding $|w_j^i|$ if $w = w_j^i$). Next recall that $X_\infty \subseteq L_{i+1} \subseteq X_i$. The subshift L_{i+1} is defined by the words $w_1^{i+1}, \dots, w_{k_{i+1}}^{i+1}$ and L_{i+1} is defined by the words $w_1^{i+1}, \dots, w_{k_{i+1}}^{i+1}$. The words $w_1^{i+1}, \dots, w_{k_{i+1}}^{i+1}$ are themselves concatenations of the words $u_1^i, \dots, u_{k_i}^i$ and induction hypothesis (v) guarantees that for any $1 \leq j \leq k_i$ and any $1 \leq t \leq k_{i+1}$, the relative frequency with which w_j^i appears in the concatenation defining w_t^{i+1} lies between $(1 - \varepsilon_i)/k_i$ and $(1 + \varepsilon_i)/k_i$. Recall that no word of length $|w_{j_1}^i|$ occurs as a subword of both $w_{j_1}^i w_{j_1}^i$ and $w_{j_2}^i w_{j_2}^i$, for $j_1 \neq j_2$, and so words of length $|w_{j_1}^i|$ that occur as subwords of $w_{j_1}^i w_{j_1}^i$ can occur only in $w_{j_1}^i w_{j_1}^i$ or possibly as a subword of $u_{j_2}^i u_{j_3}^i$ for some $j_2 \neq j_3$ and in this case the occurrence must partially overlap both of the concatenated words. Since μ is ergodic, if w is a subword of some $w_j^i w_j^i$ of length $|w_j^i|$, then

$$\mu([w]) = \frac{1}{2n+1} \sum_{m=-n}^n \mathbf{1}_{[w]}(\sigma^m x)$$

for μ -almost every $x \in X_\infty$. Fix some such x and choose some way to parse x into a concatenation of the words $u_1^i, \dots, u_{k_i}^i$. Let $\mathcal{I} \subseteq \mathbb{Z}$ be the locations where the words in this concatenation begin; this is an arithmetic progression in \mathbb{Z} with gap $N_i M_i$. The frequency with which a shift of x brings one of the elements of \mathcal{I} within distance N_i of the origin (meaning when the word of length N_i determined by this shift of x is a word that partially overlaps the break between two of the words in our concatenation) is $N_i/N_i M_i = 1/M_i$. Thus we can check:

$$\begin{aligned} \frac{(1 - 2\varepsilon_i)|s(w)|}{k_i|w_1^i|} &\leq \frac{(1 - \varepsilon_i)|s(w)|}{k_i|w_1^i|} - \frac{N_i}{N_i M_i} \\ &\leq \lim_{n \rightarrow \infty} \frac{1}{2n+1} \sum_{m=-n}^n \mathbf{1}_{[w]}(\sigma^m x) \leq \frac{(1 + \varepsilon_i)|s(w)|}{k_i|w_1^i|} + \frac{N_i}{N_i M_i} \\ &\leq \frac{(1 + 2\varepsilon_i)|s(w)|}{k_i|w_1^i|} \end{aligned}$$

since $M_i > |w_1^i|k_i/\varepsilon_i$ by induction hypothesis (vi) (recall that $s(w)$ is 1 unless it counts the number of occurrences of w in the concatenation $w_j^i w_j^i$ where it occurs). Therefore,

$$\frac{(1 - 2\varepsilon_i)|s(w)|}{k_i|w_1^i|} \leq \mu([w_j^i]) \leq \frac{(1 + 2\varepsilon_i)|s(w)|}{k_i|w_1^i|}$$

for all $1 \leq j \leq k_i$. Since

$$\mu\left(\bigcup_{w \in \mathcal{V}} [w]\right) \geq \frac{3}{4}$$

and since $\mu[w] \leq (1 + 2\varepsilon_i)/k_i$ for each $w \in \mathcal{V}$, it follows that $|\mathcal{V}| \geq (3k_i)/(4 + 8\varepsilon_i)$.

For each $1 \leq j \leq k_i$, let $A_j \subseteq \{1, 2, \dots, |w_j^i|\}$ be the set

$$A_j = \{s(w) : w \text{ is a word of length } |w_j^i| \text{ that occurs } w_j^i w_j^i \text{ and is in } \mathcal{V}\}.$$

Since

$$\mu \left(\bigcup_{w \in \mathcal{V}} [w] \right) \geq \frac{3}{4}$$

and since

$$\frac{(1 - 2\varepsilon_i)|s(w)|}{k_i|w_1^i|} \leq \mu[w] \leq \frac{(1 + 2\varepsilon_i)|s(w)|}{k_i|w_1^i|}$$

for all $w \in \mathcal{V}$, the number of elements of \mathcal{V} is at least

$$\frac{3}{4} \cdot \frac{k_i|w_1^i|}{(1 + 2\varepsilon_i)} \geq \frac{k_i|w_1^i|}{2},$$

where we count each $w \in \mathcal{V}$ with multiplicity $|s(w)|$. Therefore for at least half of the integers, $1 \leq j \leq k_i$ we have

$$|A_j| \geq \frac{k_i}{2}.$$

By Lemma 4.3, there exists some integer $1 \leq s \leq k_i$ such that $s \in A_j$ for at least $1/4$ of the integers $1 \leq j \leq k_i$. Let \mathcal{T} be this set of integers. Then \mathcal{V} contains the subword of length $|w_1^i|$ that occurs in $w_t^i w_t^i$, beginning at location s for all $t \in \mathcal{T}$. Since $d_H(w_{t_1}^i, w_{t_2}^i) > 1/4$ for all $t_1 \neq t_2$, it follows that the subword of length $|w_{t_1}^i|$ that occurs in $w_{t_1}^i w_{t_1}^i$ starting at location s is Hamming distance at least $1/4$ from the analogous subword in $w_{t_2}^i w_{t_2}^i$. It follows from our construction that every element of \mathcal{V} is within Hamming distance $1/8$ of some element of \mathcal{U} . Since two words of Hamming distance $1/4$ cannot be within distance $1/8$ of the same element of \mathcal{U} , it follows that $|\mathcal{U}|$ is at least $|\mathcal{T}| \geq k_i/4$. But by construction, $k_i/4 > b_{N_i}$, by induction hypothesis (ii). Thus $|\mathcal{U}| > b_{N_i}$, and since \mathcal{U} was arbitrary, (1) holds.

Analysis of $P_\sigma^-(n)$. Our goal is to show that

$$K(P_i, \varepsilon_i, \sigma) < a_{P_i}.$$

Fix $i \in \mathbb{N}$. Note that if $w \in \mathcal{L}_{P_i}(X_\infty)$, then $w \in \mathcal{L}_{P_i}(X_{i+1})$. By induction hypothesis (vi) recall that if \mathcal{W} is the set of words in the language of X_{i+1} that occur as subwords of $v_1^{i+1}, \dots, v_{k_{i+1}}^{i+1}$ then for any ergodic measure μ supported on X_{i+1} we have

$$\mu \left(\bigcup_{w \in \mathcal{W}} [w] \right) > 1 - \varepsilon_i.$$

Therefore we can take ε_i -Hamming balls centered on words in \mathcal{W} as a way to cover a subset of μ -measure at least $1 - \varepsilon_i$. But by construction, the words $v_1^{i+1}, \dots, v_{k_{i+1}}^{i+1}$ are all periodic words of period $|w_1^i| = |w_2^i| = \dots = |w_{k_i}^i|$ and so

$$|\mathcal{W}| \leq |w_1^i| + |w_2^i| + \dots + |w_{k_i}^i|.$$

By (5) we know that this quantity is at most a_{P_i} . Therefore it is possible to cover a subset X_∞ of μ -measure at least $1 - \varepsilon_i$ with at most a_{P_i} many ε_i -Hamming balls around words of length P_i . Therefore

$$K(P_i, \varepsilon_i, \sigma) < a_{P_i}$$

and so (2) holds.

Our construction produces a subshift (X_∞, σ) with the property that for any ergodic measure μ supported on σ , we have $P_\sigma^-(n) \prec (a_n)$ and $P_\sigma^+(n) \succ (b_n)$. Since all ergodic measures supported on X_∞ satisfy $P_\sigma^-(n) \prec (a_n)$ and since (a_n)

grows subexponentially, Katok's theorem guarantees that X_∞ supports only zero entropy measures. Furthermore, the Jewett-Krieger Theorem (see for example, Petersen [8]) guarantees that if (Y, T, μ) is an ergodic system of (measure theoretic) entropy less than $\log(N)$, then there is a minimal and uniquely ergodic system that is measure theoretically isomorphic to our system. Taking this model for the system, we have the existence of a subshift with all of the desired properties.

This completes the proof of Theorem 1.2. \square

REFERENCES

- [1] M. BOSHERNITZAN. A unique ergodicity of minimal symbolic flows with linear block growth. *Journal d'Analyse Math.* **44**, no. 1 (1984) 77–96.
- [2] V. CYR & B. KRA. Counting generic measures for a shift of linear growth. *J. Eur. Math. Soc. (JEMS)*, **21** (2019), no. 2, 355–380.
- [3] S. FERENCZI. Measure-theoretic complexity of ergodic systems. *Israel J. Math.*, **100** (1997) 189–207.
- [4] N. FRANTZIKINAKIS & B. HOST. The logarithmic Sarnak conjecture for ergodic weights. *Ann. Math.* **187**, no. 3 (2018) 869–931.
- [5] A. KATOK. Lyapunov exponents, entropy and periodic orbits for diffeomorphisms. *Pub. Math. de l'I.H.E.S.* **51** (1980) 137–173.
- [6] A. KATOK & J. P. THOUVENOT. Slow entropy type invariants and smooth realization of commuting measure-preserving transformations. *Ann. Inst. H. Poincaré Probab. Statist.* **33** (1997), no. 3, 323–338.
- [7] R. MCNAMARA. Sarnak's Conjecture for Sequences of Almost Quadratic Word Growth. [arXiv:1901.06460](https://arxiv.org/abs/1901.06460)
- [8] K. PETERSEN. Ergodic theory. Cambridge Studies in Advanced Mathematics, **2**. Cambridge University Press, Cambridge, 1983.
- [9] J. SERAFIN. Non-existence of a universal zero-entropy system. *Israel J. Math.* **194** (2013), no. 1, 349–358.

BUCKNELL UNIVERSITY, LEWISBURG, PA 17837 USA
E-mail address: `van.cyr@bucknell.edu`

NORTHWESTERN UNIVERSITY, EVANSTON, IL 60208 USA
E-mail address: `kra@math.northwestern.edu`