

HEAT SEMIGROUP ON A COMPLETE RIEMANNIAN MANIFOLD

BY PEI HSU

University of Illinois at Chicago

Let M be a complete Riemannian manifold and $p(t, x, y)$ the minimal heat kernel on M . Let P_t be the associated semigroup. We say that M is stochastically complete if $\int_M p(t, x, y) dy = 1$ for all $t > 0, x \in M$; we say that M has the C_0 -diffusion property (or the Feller property) if $P_t f$ vanishes at infinity for all $t > 0$ whenever f is so. Let $x_0 \in M$ and let $\kappa(r)^2 \geq -\inf\{\text{Ric}(x): \rho(x, x_0) \leq r\}$ (ρ is the Riemannian distance). We prove that M is stochastically complete and has the C_0 -diffusion property if $\int_c^\infty \kappa(r)^{-1} dr = \infty$ by studying the radial part of the Riemannian Brownian motion on M .

1. Introduction. Let (M, g) be a noncompact, connected Riemannian manifold and Δ the Laplace–Beltrami operator. The Riemannian Brownian motion X on M is the minimal diffusion process on M associated with the operator $\Delta/2$. The transition density $p(t, x, y)$ of the Brownian motion is the minimal fundamental solution of the heat operator $\partial/\partial t - \Delta/2$. Let $M \cup \{\partial\}$ be the one-point compactification of M . As usual, we regard X as a continuous Markov process on $M \cup \{\partial\}$. Let e be the explosion time of X , that is,

$$e = \inf\{t > 0: X_t = \partial\}.$$

When $P_x[e = \infty] = 1$ for one $x \in M$ (hence for all $x \in M$ since $P_x[e = \infty]$ is harmonic on M), we say M is stochastically complete. Intuitively, stochastic completeness means that the Brownian motion will not drift to infinity in finite amount of time. Since clearly

$$P_x[e > t] = \int_M p(t, x, y) dy,$$

stochastic completeness also means that the heat kernel on M is conservative:

$$\int_M p(t, x, y) dy = 1 \quad \text{for all } t > 0, x \in M.$$

Stochastic completeness is equivalent to the uniqueness of solution of heat equation with L^∞ initial data.

Let $C_0(M)$ be the space of continuous functions on M vanishing at infinity. Consider the diffusion semigroup

$$P_t f(X) = \int_M f(y) p(t, x, y) dy$$

Received December 1987; revised June 1988.

AMS 1980 subject classification. 58J32.

Key words and phrases. Riemannian manifold, comparison theorems, Riemannian Brownian motion, stochastic completeness, Ricci curvature, C_0 -diffusion.

for $f \in C_0(M)$. We say that P_t is a C_0 -diffusion or M has the C_0 -diffusion property (or the Feller property) if $P_t C_0(M) \subset C_0(M)$, namely if the function space $C_0(M)$ is invariant under P_t for any $t > 0$. Intuitively, P_t is a C_0 -diffusion if the Brownian motion starting from very far has small probability of visiting a fixed compact set before fixed time (see the precise statement in Lemma 3.1). Thus again C_0 -diffusion property means the Brownian motion will not drift away from where it starts too fast. We would like to seek a geometric condition on M for stochastic completeness and the C_0 -diffusion property.

We will assume that M is geometrically complete, namely, complete under its Riemannian distance ρ . Let $x_0 \in M$ be a fixed reference point on M , and $\rho(x) = \rho(x, x_0)$. Set $\rho(x, \partial) = \infty$ for all $x \in M$. Then we have $e = \inf\{t > 0: \rho(X_t) = \infty\}$. Geometrically, the speed at which the Brownian motion wanders away from its starting point is controlled by the lower bound of the Ricci curvature. A quantitative version of this statement can be found in Lemma 3.2 below. Let

$$\kappa(r)^2 \geq -\inf\{\text{Ric}(x): \rho(x) \leq r\}.$$

Then it is reasonable to expect that conditions for stochastic completeness and the C_0 -diffusion property can be expressed as growth conditions on the function κ .

Stochastic completeness and the C_0 -diffusion property has been discussed by various authors. We mention Azencott [1] and Yau [10] (also Doziuk [2]), who proved, among other things, that a geometrically complete manifold with Ricci curvature bounded from below by a constant [namely, $\kappa(r) \leq c < \infty$] is stochastically complete and has the C_0 -diffusion property. Azencott [1] pointed out that if a Cartan–Hadamard manifold has sectional curvature bounded from above by $-\rho(x)^{2+\epsilon}$, then it is not stochastically complete. He also noted that every Cartan–Hadamard manifold has the C_0 -diffusion property. The work of Hsu and March [5] implies that M is stochastically complete if $\kappa(r) \leq L(1+r)$ for some $L > 0$. Karp and Li [7] in an unpublished article showed that M is stochastically complete if the volume of the geodesic ball $B_R(x_0)$ of radius R centered at x_0 satisfies the growth condition $\text{vol}(B_R(x_0)) \leq \exp[cR^2]$ for some constant c . This was improved by Grigor'yan [4] to $\text{vol}(B_R(x_0)) \leq \exp f(R)$ for any increasing f such that $\int_c^\infty rf(r)^{-1} dr = \infty$. It was also shown in Karp and Li [7] that M has the C_0 -diffusion property if there exists a constant L such that $\kappa(r) \leq L(1+r)$.

Our result in this article can be simply stated as follows.

THEOREM. *If M is complete and if $\int_c^\infty \kappa(r)^{-1} dr = \infty$, then M is stochastically complete and has the C_0 -diffusion property.*

The stochastic completeness part of our result is implied by the result of Grigor'yan [4]. Our result of the C_0 -diffusion property improves the condition of Karp and Li [7]. We show our results by studying the radial part of the Riemannian Brownian motion of M .

2. Stochastic completeness. Consider the radial process $\rho(X_t)$. According to Kendall [8], there exists a standard Brownian motion β_t and a *nondecreasing* process L_t with initial value zero which increases only when X_t belongs to the cut-locus $C(x_0)$ of x_0 , such that for $t < e$,

$$(2.1) \quad \rho(X_t) = \beta_t + \frac{1}{2} \int_0^t \Delta \rho(X_s) ds - L_t.$$

Now $\Delta \rho(x)$ is smooth on $M \setminus C(x_0)$. By the Hessian comparison theorem (Greene and Wu [3], pages 19–28), $\Delta \rho$ is locally bounded away from point x_0 . We also notice that the Riemannian volume measure of $C(x_0)$ is zero. As a consequence the Brownian motion spends zero amount of time on $C(x_0)$ and the term $\int_0^t \Delta \rho(X_s) ds$ in (2.1) is well defined. (2.1) is essentially the result of applying Itô's formula to function $\rho(x)$. See the appendix to this section for a proof.

Without loss of generality, we assume that function κ satisfies the following three extra conditions:

- (i) $\kappa(c) > 0$.
- (ii) κ is nondecreasing.
- (iii) $\lim_{s \rightarrow \infty} \kappa(s) = \infty$.

Let $G: [0, \infty) \rightarrow [0, \infty)$ be the unique solution of the equation

$$G''(r) = \kappa(r)^2 G(r), \quad G(0) = 0, \quad G'(0) = 1.$$

Let (R^n, g^*) be the Riemannian manifold with the metric $g^* = dr^2 + G(r)^2 d\theta^2$ [(ρ, θ) is the usual polar coordinates on R^n]. The radial Ricci curvature of (R^n, g^*) is

$$\text{Ric}(\dot{\rho}, \dot{\rho}) = -(n - 1) \frac{G''(\rho)}{G(\rho)} = -(n - 1) \kappa(\rho)^2.$$

Now we use the Laplacian comparison theorem (Greene and Wu [3], page 26) to manifolds (M, g) and (R^n, g^*) , and conclude that on $M \setminus C(x_0)$,

$$(2.2) \quad \Delta \rho \leq (n - 1) \frac{G'(\rho)}{G(\rho)}.$$

Let ρ_t^* be the process on $[0, \infty)$ determined by the stochastic differential equation

$$(2.3) \quad \rho_t^* = \beta_t + \frac{n - 1}{2} \int_0^t \frac{G'(\rho_s^*)}{G(\rho_s^*)} ds, \quad \rho_0^* = 0.$$

Let e^* be the explosion time of ρ_t^* . By a comparison theorem for solutions of stochastic differential equations (Ikeda and Watanabe [6], pages 352–356) and the positivity of L_t , (2.1), (2.2) and (2.3) imply $\rho(X_t) \leq \rho_t^*$ for all $t > 0$. It follows that $e^* \leq e$ a.s. Thus it suffices to show that $e^* = \infty$ a.s. Now we are dealing with a one-dimensional Itô-type diffusion (2.3). The condition for nonexplosion in this case is known (Ikeda and Watanabe [6], pages 365–367): $e^* = \infty$ a.s. if

and only if

$$(2.4) \quad I(G) \stackrel{\text{def}}{=} \int_c^\infty G(r)^{1-n} dr \int_c^r G(s)^{n-1} ds = \infty.$$

The above condition (2.4) is the best condition for the nonexplosion of Riemannian Brownian motion in general. To convert this condition into the more explicit condition on the function $\kappa(s)$ stated in our theorem, we need to show that $\int_c^\infty \kappa(r)^{-1} ds = \infty$ implies $I(G) = \infty$.

Observe first that $\kappa(r)^2 = G''(r)/G(r)$ is nondecreasing. Integrating by parts, we have

$$\int_0^r G(s)^2 d\kappa(s)^2 = G''(r)G(r) - G'(r)^2 + 1.$$

Hence $G''(r)G(r) - G'(r)^2 \geq -1$, or

$$\left[\frac{G'(r)}{G(r)} \right]^2 \leq \frac{G''(r)}{G(r)} + \frac{1}{G(r)^2} \leq c_1 \kappa(r)^2$$

for $r \geq c$ and $c_1 = 1 + \kappa(c)^{-1}G(c)^{-2}$. Integrating by parts again, we have

$$\int_c^r G(s)^{n-1} ds = \frac{1}{n} \int_c^r \frac{dG^n(s)}{G'(s)} \geq \frac{1}{n} \frac{G(r)^n}{G'(r)} - \frac{1}{n} \frac{G(c)^n}{G'(c)}.$$

Since G grows at least exponentially, we have $\int_c^\infty G(r)^{1-n} dr < \infty$. Therefore

$$I(G) \geq \frac{1}{n} \int_c^\infty \frac{G(r)}{G'(r)} dr - c_2 \geq \frac{c_1^{-1/2}}{n} \int_c^\infty \kappa(r)^{-1} ds - c_2 = \infty.$$

This shows M is stochastically complete.

Appendix to Section 2. We give a more direct proof of (2.1) (compare with Kendall [8]). The key to the proof is the following remarkable property of the distance function ρ . Let ϕ be a smooth, nonnegative function on M with compact support, then

$$(A) \quad \int_M \rho(x) \Delta\phi(x) dx \leq \int_{M \setminus C(x_0)} \phi(x) \Delta\rho(x) dx.$$

[Recall that $C(x_0)$ is the cut-locus of x_0 .] Inequality (A) is proved in the appendix of Yau [11]. Now let $\Delta\rho$ denote the Laplacian of ρ in the *distributional sense*. The above inequality implies that the distribution

$$(2.5) \quad \mu = \Delta\rho I_{M \setminus C(x_0)} - \Delta\rho$$

is nonnegative on nonnegative test functions. It follows from the Riesz representation theorem that μ is a Radon measure on M concentrated on the cut-locus $C(x_0)$. Thus $\Delta\rho$, the distributional Laplacian of ρ , is itself realized as a measure on M . A generalized Itô formula (Meyer [9]) applied to the function ρ gives immediately formula (2.1) with L_t equal to the continuous positive additive functional associated with the measure μ .

It is interesting to point out that (2.1) and (A) are in fact equivalent. Thus Kendall's proof of (2.1) amounts to a probabilistic proof of (A).

3. C_0 -diffusion property. We need two lemmas. The first is elementary and a proof can be found in Azencott [1]. The proof of the second is essentially contained in Hsu and March [5].

LEMMA 3.1. P_t is a C_0 -diffusion semigroup if and only if for any compact set K and any fixed $t > 0$, we have

$$(3.1) \quad \lim_{\rho(x) \rightarrow \infty} P_x[T_K \leq t] = 0,$$

where T_K is the first hitting time of K :

$$T_K = \inf\{t > 0: X_t \in K\}.$$

LEMMA 3.2. Let

$$\tau = \inf\{t > 0: \rho(X_0, X_t) = 1\}.$$

If the Ricci curvature of M in the geodesic ball $B_1(X_0)$ centered at X_0 with radius 1 is bounded from below by $-L^2 \leq -1$, then

$$P_x[\tau \leq c_1 L^{-1}] \leq e^{-c_2 L}$$

for some universal positive constants c_1 and c_2 .

PROOF. This is essentially Lemma 4 of Hsu and March [5]. In the present case, the geodesic ball $B_1(X_0)$ may not entirely lie within the cut-locus. However, we notice the following three facts:

(a) From (2.1) we have for any $s \leq t$,

$$\rho(X_t) - \rho(X_s) \leq \beta_t - \beta_s + \frac{1}{2} \int \Delta \rho(X_s) ds.$$

(b) Away from the cut-locus $C(X_0)$, the Laplacian comparison theorem (Greene and Wu [3], page 26) gives

$$\Delta \rho \leq (n - 1)L \coth L\rho.$$

(c) The Brownian motion spends zero amount of time in the cut-locus.

The above three facts enable us to carry out the proof of the lemma by repeating, mutatis mutandis, the proof of Hsu and March [5]. \square

We now prove the second part of our theorem. By Lemma 3.1, it is enough to show (3.1). We may assume that $K = \overline{B_R(x_0)}$, the geodesic ball centered at x_0 with radius R . Let τ be defined as in Lemma 3.2. Consider the following

stopping times:

$$\begin{aligned}
 s_0 &= 0, \\
 \tau_1 &= \tau, \\
 s_1 &= \inf\{t \geq \tau_1: \rho(x_0, X_t) = \rho(x) - 1\}, \\
 \tau_2 &= \tau \circ \theta_{s_1}, \\
 s_2 &= \inf\{t \geq \tau_2 + s_1: \rho(x_0, X_t) = \rho(x) - 2\}, \\
 &\dots \\
 \tau_n &= \tau \circ \theta_{s_{n-1}}, \\
 s_n &= \inf\{t \geq \tau_n + s_{n-1}: \rho(X_t) = \rho(x) - n\}
 \end{aligned}$$

(θ is the shift operator). It is clear that

$$(3.2) \quad T_K \geq s_{[\rho(x)-R]} \geq \tau_1 + \tau_2 + \dots + \tau_{[\rho(x)-R]}$$

($[a]$ denotes the integral part of a). Now since $\rho(X_{s_{k-1}}) = \rho(x) - k + 1$, we have by Lemma 3.2

$$(3.3) \quad P_x[\tau_k \leq c_1 \kappa(\rho(x) - k + 2)^{-1}] \leq e^{-c_2 \kappa(\rho(x) - k + 2)}.$$

Choose $n(x, t)$ to be the first integer n such that

$$\sum_{k=1}^n \kappa(\rho(x) - k + 2)^{-1} > \frac{t}{c_1}.$$

In view of the condition $\int_c^\infty \kappa(s)^{-1} ds = \infty$ such $n(x, t)$ exists for sufficiently large $\rho(x)$, and furthermore we must have $[\rho(x) - R] \geq n(x, t)$. Hence

$$\{T_K \leq t\} \subset \{\tau_1 + \tau_2 + \dots + \tau_{[\rho(x)-R]} \leq t\} \subset \bigcup_{k=1}^{n(x,t)} \{\tau_k \leq c_1 \kappa(\rho(x) - k + 2)^{-1}\}.$$

Using (3.3), we have

$$(3.4) \quad P_x[T_K \leq t] \leq \sum_{k=1}^{n(x,t)} e^{-c_2 \kappa(\rho(x) - k + 2)} \leq \int_{m(x,t)}^{\rho(x)+1} e^{-c_2 \kappa(r)} dr.$$

Here we have set $m(x, T) = \rho(x) - n(x, t) + 1$ for simplicity. The choice of $n(x, t)$ implies that

$$\sum_{k=1}^{n(x,t)-1} \kappa(\rho(x) - k + 2)^{-1} \leq \frac{t}{c_1}$$

or

$$(3.5) \quad \int_{m(x,t)+2}^{\rho(x)+2} \kappa(r)^{-1} \leq \frac{t}{c_1}.$$

Using (3.4), (3.5) and the elementary inequality $e^{-c_2 \kappa} \leq a e^{-c_2 a} / \kappa$ for $\kappa \geq a \geq c_2^{-1}$,

we have

$$P_x[T_k \leq t] \leq \kappa(m(x, t))e^{-c_2\kappa(m(x, t))} \left[2\kappa(m(x, t))^{-1} + \frac{t}{c_1} \right],$$

provided that $m(x, t) \geq c_2^{-1}$. Now the assumption $\int_c^\infty \kappa(r)^{-1} dr = \infty$ implies by (3.5) that $m(x, t) \rightarrow \infty$ as $\rho(x) \rightarrow \infty$. (3.1) follows immediately from the above inequality by letting $\rho(x) \rightarrow \infty$. The theorem is proved.

Acknowledgment. I thank Harold Donnelly for showing me reference [4].

REFERENCES

- [1] AZENCOTT, R. (1974). Behavior of diffusion semigroups at infinity. *Bull. Soc. Math. France* **102** 193–240.
- [2] DODZIUK, J. (1983). Maximum principle for parabolic inequalities and the heat flow on open manifolds. *Indiana Univ. Math. J.* **32** 703–716.
- [3] GREENE, R. E. and WU, H. (1979). *Function Theory on Manifolds which Possess a Pole. Lecture Notes in Math.* **699**. Springer, New York.
- [4] GRIGOR'YAN, A. A. (1987). On stochastic completeness of complete manifolds. *Soviet Math. Dokl.* **34** 310–313.
- [5] HSU, P. and MARCH, P. (1985). The limiting angle of certain Riemannian Brownian motions. *Comm. Pure Appl. Math.* **38** 755–768.
- [6] IKEDA, N. and WATANABE, S. (1981). *Stochastic Differential Equations and Diffusion Processes*. Kodansha, Tokyo/North-Holland, Amsterdam.
- [7] KARP, L. and LI, P. (1984). On the heat equation on complete manifolds. Preprint.
- [8] KENDALL, W. S. (1987). The radial part of Brownian motion on a manifold: A semimartingale property. *Ann. Probab.* **15** 1491–1500.
- [9] MEYER, P.-A. (1978). La formule d'Itô pour le mouvement brownien d'après Brosamler. *Séminaire de Probabilités XII, 1976 / 77. Lecture Notes in Math.* **649** 764–769. Springer, New York.
- [10] YAU, S.-T. (1978). On the heat kernel of a complete Riemannian manifold. *J. Math. Pures Appl.* **57** 191–201.
- [11] YAU, S.-T. (1976). Some function-theoretic properties of complete Riemannian manifolds and their applications to geometry. *Indiana Univ. Math. J.* **25** 659–670.

DEPARTMENT OF MATHEMATICS
NORTHWESTERN UNIVERSITY
EVANSTON, ILLINOIS 60208