

# INVERSE SPECTRAL PROBLEM FOR ANALYTIC $(\mathbb{Z}/2\mathbb{Z})^n$ -SYMMETRIC DOMAINS IN $\mathbb{R}^n$

HAMID HEZARI AND STEVE ZELDITCH

ABSTRACT. We prove that bounded real analytic domains in  $\mathbb{R}^n$  with the symmetries of an ellipsoid, and with one axis length fixed, are determined by their Dirichlet or Neumann eigenvalues among other bounded real analytic domains with the same symmetries and axis length. Some non-degeneracy conditions are also imposed on the class of domains. It follows that bounded, convex analytic domains are determined by their spectra among other such domains. This seems to be the first positive result on the well-known Kac problem, can one hear the shape of a drum?, in higher dimensions.

## 1. INTRODUCTION AND THE STATEMENT OF RESULTS

The purpose of this article is to prove that bounded analytic domains  $\Omega \subset \mathbb{R}^n$  with  $\pm$  reflection symmetries across all coordinate axes, and with one axis height fixed (and also satisfying some generic non-degeneracy conditions) are spectrally determined among other such domains. This inverse result (Theorem 1) gives a higher dimensional analogue of the main result of [Z2] that “bi-axisymmetric” real analytic plane domains are spectrally determined among other bounded analytic plane domains with the symmetry of an ellipse. To our knowledge, it is the first positive higher dimensional inverse spectral result for Euclidean domains which is not restricted to balls. Negative results (i.e. constructions of non-isometric isospectral pairs) are given in [U, GW, GWW] (see also [GS] for some non-Euclidean domains). Higher dimensional inverse results for semi-classical Schrödinger operators with similar symmetries have recently been proved in [GU, H].

1.1. **Statement of results.** We consider the eigenvalue problem on the domain  $\Omega$  with the Euclidean Laplacian  $\Delta_\Omega^B$  and with boundary conditions  $B$ :

$$(1) \quad \begin{cases} \Delta_\Omega^B \varphi_j(x) = -\lambda_j^2 \varphi_j(x), & \langle \varphi_i, \varphi_j \rangle = \delta_{ij}, \quad (x \in \Omega) \\ B\varphi_j(y) = 0, & y \in \partial\Omega. \end{cases}$$

The boundary conditions could be either Dirichlet  $B\varphi = \varphi|_{\partial\Omega}$ , or Neumann  $B\varphi = \partial_\nu \varphi|_{\partial\Omega}$  where  $\partial_\nu$  is the interior unit normal.

The  $(\mathbb{Z}/2\mathbb{Z})^n$  symmetries of the title are the maps

$$(2) \quad \sigma_j : (x_1, \dots, x_n) \rightarrow (x_1, \dots, -x_j, x_{j+1}, \dots, x_n)$$

and we assume that they are isometries of  $\Omega$ . The symmetry assumption implies that the intersections of the coordinate axes with  $\Omega$  are projections of bouncing ball orbits preserved by the symmetries. We recall that a bouncing ball orbit  $\gamma$  is a 2-link periodic trajectory of

---

Research partially supported by NSF grant #DMS-06-03850.

the billiard flow, i.e. a reversible periodic billiard trajectory that bounces back and forth along a line segment orthogonal to the boundary at both endpoints. The endpoints of the projection to  $\Omega$  of the bouncing ball orbit are fixed points of all but one of the isometries  $\sigma_j$ ; the remaining one fixes the projected orbit setwise but interchanges the endpoints. We add the generic condition that at least one of these bouncing ball orbits is *non-degenerate* (see (3) for the conditions). We also fix the length  $L_\gamma = 2L$  of this bouncing ball orbit  $\gamma$ .

We denote by  $\mathcal{D}_L$  to be the class of all bounded real-analytic domains  $\Omega \subset \mathbb{R}^n$  satisfying these assumptions. Thus,  $\mathcal{D}_L$  is the class of domains such that:

- (3)  $\left\{ \begin{array}{l} \text{(i)} \quad \sigma_j : \Omega \rightarrow \Omega \text{ is an isometry for all } j = 1, \dots, n; \\ \text{(ii)} \quad \text{one of the coordinate axis bouncing ball orbits, called } \gamma, \text{ is of length } 2L \\ \text{(iii)} \quad \text{the lengths } 2rL \text{ of all iterates } \gamma^r (r = 1, 2, 3, \dots) \text{ have multiplicity one in } Lsp(\Omega); \\ \text{(iv)} \quad \gamma \text{ is non-degenerate, i.e. } 1 \text{ is not an eigenvalue of its Poincaré map } P_\gamma; \\ \quad \text{if } \gamma \text{ is elliptic and } \{e^{\pm i\alpha_1}, \dots, e^{\pm i\alpha_{n-1}}\} \text{ are the eigenvalues of } P_\gamma, \text{ we} \\ \quad \text{further require that } \{\alpha_1, \dots, \alpha_{n-1}\} \text{ are linearly independent over } \mathbb{Q}. \text{ We assume the} \\ \quad \text{same independence condition in the Hyperbolic case or mixed cases.} \end{array} \right.$

Here,  $Lsp(\Omega)$  is the length spectrum of  $\Omega$ , i.e. the set of lengths of closed billiard trajectories (cf. [PS, Z3]). Multiplicity one means that there exists precisely one closed billiard trajectory of the given length up to time reversal. Let  $\text{Spec}_B(\Omega)$  denote the spectrum of the Laplacian  $\Delta_\Omega^B$  of the domain  $\Omega$  with boundary conditions  $B$  (Dirichlet or Neumann).

**THEOREM 1.** *For Dirichlet (or Neumann) boundary conditions  $B$ , the map  $\text{Spec}_B : \mathcal{D}_L \rightarrow \mathbb{R}_+^{\mathbb{N}}$  is 1-1.*

In other words, if two bounded real analytic domains  $\Omega_1, \Omega_2 \subset \mathbb{R}^n$  possessing the symmetries of an ellipsoid and satisfying the non-degeneracy and length assumptions of (3) have the same Dirichlet (resp. Neumann) spectra, then they are isometric. To our knowledge, the only prior positive result on the inverse spectral problem for higher dimensional bounded domains is that a domain with the Dirichlet (or Neumann) spectrum of a ball must be a ball [KAC]. In that case the proof is based on the trace of the heat semi-group rather than the wave group or resolvent kernel. The heat trace for the Dirichlet (or Neumann) Laplacian  $\Delta_\Omega^D$  of a bounded domain has the singularity expansion,

$$\text{Tre}^{t\Delta_\Omega^D} \sim t^{-n/2}(C_n \text{Vol}_n(\Omega) + C'_n \text{Vol}_{n-1}(\partial\Omega)t^{-1/2} + \dots), \quad t \rightarrow 0^+,$$

where  $C_n, C'_n$  are constants depending only on the dimension. Hence the volume and surface measure are spectral invariants. The ball is determined as the unique domain where the isometric inequality  $\text{Vol}_{n-1}(\partial\Omega) \geq A_n \text{Vol}_n(\Omega)^{\frac{n-1}{n}}$  (for a certain constant  $A_n$ ) is an equality.

Our proof of Theorem 1 has a similar form in that we calculate some special spectral invariants and then use the invariants to uniquely determine the domain. But instead of the heat semi-group we use the wave group  $e^{it\sqrt{-\Delta_\Omega^D}}$  or more precisely the semi-classical resolvent  $R_\Omega^D(k) = -(\Delta_\Omega^D + k^2)^{-1}$  for  $k \in \mathbb{C}$ , which is a semi-classical Laplace transform of the wave group (see §2.3). Here, we are assuming that the boundary conditions are Dirichlet, but the methods and results are valid for the Neumann Laplacian  $\Delta_\Omega^N$  with only minor modifications. The spectral invariants we study are the ‘wave invariants’ associated to one of the bouncing ball orbits defined by the coordinate axes. The key advantage of these wave invariants is

that they are localized at the endpoints of the projected orbit, whereas heat invariants are integrals of curvature invariants over  $\Omega$  or  $\partial\Omega$ . In Theorem 2, the wave invariants of bouncing ball orbits are expressed in terms of the Taylor coefficients of the defining function of  $\Omega$  near the endpoints. Under our symmetry assumptions, the Taylor coefficients are determined from the wave invariants. That proves Theorem 1.

As a corollary, we obtain a result for convex analytic domains that does not require any length to be marked.

**COROLLARY 1.** *Let  $\mathcal{C}$  be the class of analytic convex domains with  $(\mathbb{Z}/2\mathbb{Z})^n$  symmetry, such that the shortest closed billiard trajectory  $\gamma_0$  is non-degenerate and satisfies the conditions (iii) and (iv) of (3). Then  $\text{Spec}_B: \mathcal{C} \rightarrow \mathbb{R}_+^{\mathbb{N}}$  is 1-1.*

This follows from Theorem 1 and a result of M. Ghomi [Gh] that the shortest closed trajectory of a centrally-symmetric convex domain is automatically a bouncing ball orbit. Hence the length of this orbit is self-marked, and it is not necessary to mark the length  $L_\gamma = 2L$  of an invariant bouncing ball orbit  $\gamma$ .

**1.2. Balian-Bloch and wave invariants at a bouncing ball orbit.** As mentioned above, the proof of Theorem 1 is based the study of spectral invariants of the Dirichlet or Neumann Laplacian of  $\Omega$  known as the Balin-Bloch (or wave trace) invariants at the closed billiard trajectories  $\gamma$  of  $\Omega$ . The Balian-Bloch invariants  $B_{\gamma,j}$  are coefficients of the regularized trace expansion

$$(4) \quad \text{Tr} R_{\Omega,\rho}^D(k) \sim \mathcal{D}_{D,\gamma}(k) \sum_{j=0}^{\infty} B_{\gamma,j} k^{-j}, \quad \Re k \rightarrow \infty,$$

of the smoothed semi-classical resolvent  $R_{\Omega,\rho}^D(k)$  where  $\hat{\rho}$  is localized at the length of the closed orbit  $\gamma$ . The smoothed semi-classical resolvent is defined in (14) and the precise statement of (4) is given in Theorem 2.1. The factor  $\mathcal{D}_{D,\gamma}(k)$  is a well-known symplectic factor that is reviewed in §2.3 and discussed in more detail in [GM, PS].

The Balian-Bloch invariants are named after the physicists who introduced them in [BB1, BB2] and studied them on a somewhat non-rigorous formal level. Since then, a long stream of mathematical works have been produced on the dual singularity expansion of the trace  $\text{Tr} \cos t\sqrt{-\Delta_\Omega^D}$  of the wave group. The classical results on wave trace invariants on compact Riemannian manifolds without boundary are due to Colin de Verdière, Chazarain and Duistermaat-Guillemin. The wave trace expansion was then generalized to manifolds with boundary by Guillemin-Melrose in [GM] (see also [PS] for a very thorough study). The semi-classical resolvent and wave group are related by a Laplace transform (see (13)) and so the semi-classical (i.e. large  $k$ ) expansion (4) is essentially the same as the singularity expansion of  $\text{Tr} \cos t\sqrt{-\Delta_\Omega^D}$ . Algorithms for calculating the coefficients in the boundaryless case were given in [G, G2, Z1]. In [Z2, Z3] an algorithm was given for calculating the invariants in the boundary case but it was only implemented for plane domains. In this article, the algorithm is developed for higher dimensional domains and explicit formulae for the Balian-Bloch or wave trace invariants are given in Theorem 2. The calculations also draw on the analysis in [H] of similar invariants for semi-classical Schrödinger operators. This result is of independent interest and is valid without any symmetry assumptions.

We now state the formulae for the Balian-Bloch invariants. They require some more notation which will be further discussed in §2 and §7. Almost the same notation is used in [Z3]. We align the axes so that the bouncing ball orbit  $\gamma$  is a vertical segment of length  $L$  with endpoints at  $A = (0, \frac{L}{2})$  and  $B = (0, -\frac{L}{2})$ , where 0 denotes the origin in the orthogonal  $x' = (x^1, \dots, x^{n-1}) \in \mathbb{R}^{n-1}$  plane. In a metric tube  $T_\epsilon(\overline{AB})$  of radius  $\epsilon$  around  $\gamma$ , we may locally express  $\partial\Omega = \partial\Omega^+ \cup \partial\Omega^-$  as the union of two graphs over a ball  $B_\epsilon(0)$  around 0 in the  $x'$ -hyperplane, namely

$$(5) \quad \partial\Omega^+ = \{x^n = f_+(x'), \quad |x'| \leq \epsilon\}, \quad \partial\Omega^- = \{x^n = f_-(x'), \quad |x'| \leq \epsilon\}.$$

We will use the standard shorthand notations for multi-indices, i.e.  $\vec{\gamma} = (\gamma_1, \dots, \gamma_{n-1})$ ,  $|\vec{\gamma}| = \gamma_1 + \dots + \gamma_{n-1}$ ,  $\vec{X}^{\vec{\gamma}} = X_1^{\gamma_1} \dots X_{n-1}^{\gamma_{n-1}}$ , and by  $\overrightarrow{h_{\pm, 2r}^{pq}}$ , we mean the  $(n-1)$ -vector

$$\overrightarrow{h_{\pm, 2r}^{pq}} = (h_{\pm, 2r}^{11, pq}, h_{\pm, 2r}^{22, pq}, \dots, h_{\pm, 2r}^{(n-1, n-1), pq}),$$

where  $[h_{\pm, 2r}^{ij, pq}]_{1 \leq i, j \leq n-1, 1 \leq p, q \leq 2r}$  is the inverse Hessian matrix of the length functional  $\mathcal{L}_\pm(x'_1, \dots, x'_{2r})$  given in (11).

The following generalizes Theorem 5.1 in [Z3] from two to higher dimensions.

**THEOREM 2.** *Let  $\Omega$  be a smooth domain with a bouncing ball orbit  $\gamma$  of length  $2L$  and let  $B_{\gamma^r, j}$  be the wave invariants associated to  $\gamma^r$  (see cf. 2.2). Then for each  $r = 1, 2, \dots$ , and  $j$  there exists a polynomial  $P_{r, j}$  such that:*

- (1)  $B_{\gamma^r, j} = P_{r, j}(\{D_{\vec{\gamma}}^{|\gamma|} f_+(0)\}, \{D_{\vec{\gamma}}^{|\gamma|} f_-(0)\})$  with  $|\gamma| \leq 2j + 2$ ; i.e. the highest order of derivatives appearing in  $B_{\gamma^r, j}$  is  $2j + 2$ .
- (2) In the polynomial expansion of  $B_{\gamma^r, j}$  the Taylor coefficients of order  $2j + 2$  appear in the form  $\{D_{2\vec{\gamma}}^{2j+2} f_\pm(0)\}$ .
- (3)  $B_{\gamma^r, 0}$  is only a function of  $r$ ,  $L$  and  $n$ , and for  $j \geq 1$

$$B_{\gamma^r, j} = \frac{B_{\gamma^r, 0}}{(2i)^{j+1}} \sum_{|\gamma|=j+1} \frac{r}{\vec{\gamma}!} \{(\overrightarrow{h_{+, 2r}^{11}})^{\vec{\gamma}} D_{2\vec{\gamma}}^{2j+2} f_+(0) - (\overrightarrow{h_{-, 2r}^{11}})^{\vec{\gamma}} D_{2\vec{\gamma}}^{2j+2} f_-(0)\} \\ + R_{2r, j}(\mathcal{J}^{2j+1} f_+(0), \mathcal{J}^{2j+1} f_-(0)),$$

where the remainder  $R_{2r, j}(\mathcal{J}^{2j+1} f_+(0), \mathcal{J}^{2j+1} f_-(0))$  is a polynomial in the designated jet of  $f_\pm$ .

- (4) In the  $(\mathbb{Z}/2\mathbb{Z})$ -symmetric case, where  $f_+ = f = -f_-$ , we have the simplified formula

$$B_{\gamma^r, j} = \frac{B_{\gamma^r, 0}}{(2i)^{j+1}} \sum_{|\gamma|=j+1} \frac{r}{\vec{\gamma}!} \left( \frac{1}{\sin \frac{r\vec{\alpha}}{2}} \cot \frac{r\vec{\alpha}}{2} \right)^{\vec{\gamma}} D_{2\vec{\gamma}}^{2j+2} f(0) + R_{2r, j}(\mathcal{J}^{2j+1} f(0)).$$

- (5) In the  $(\mathbb{Z}/2\mathbb{Z})^n$ -symmetric case, formula (4) holds with remainder in  $R_{2r, j}(\mathcal{J}^{2j} f(0))$ .

In the above notation, the  $(\mathbb{Z}/2\mathbb{Z})^n$ -symmetry assumptions in (5) are that

$$(6) \quad f_+(x') = -f_-(x'), \quad f_\pm(\sigma_j(x')) = f_\pm(x'),$$

where  $\sigma_j$  denotes the reflections in the coordinate hyperplanes of  $\mathbb{R}^{n-1}$ . The first assumption implies that there exists a function  $f(x')$  so that the top of the domain is defined by  $x_n = f(x')$  and the bottom is defined by  $x_n = -f(x')$ . The further symmetry assumptions then say that  $f$  is an even function in every variable  $x^i$ ,  $1 \leq i \leq n-1$ , i.e.

$$(7) \quad f(x^1, \dots, x^{n-1}) = F((x^1)^2, \dots, (x^{n-1})^2), \quad F \in C^\omega(\mathbb{R}^{n-1}).$$

The value of these explicit formulae is demonstrated by applications such as Theorem 1. Theorem 2 may be viewed as an alternative to the use of Birkhoff normal forms methods for calculating wave trace coefficients as in [G, G2, Z1, Z2, ISZ, SZ]. Further discussion and comparison of methods is given at the end of the introduction.

Our proof of Theorem 2 is rather different from that in [Z3, Z4]. It is based on the construction and analysis of a microlocal monodromy operator associated to  $\gamma$ , inspired by the works of Sjöstrand-Zworski [SZ] and Cardoso-Popov [CP] (see also [ISZ]), but employing a layer potential analysis more closely related to that in [Z3, HZ]. The trace asymptotics are eventually reduced to those of a boundary integral operator in Proposition 5.6 and Corollary 5.7, and then to the stationary phase asymptotics of a certain oscillatory integral in Theorem 6.3. Since the method and results give a higher dimensional generalization of the analogous results of [Z3, Z4], there is some overlap in the arguments from the two-dimensional case; we have tried to minimize the overlap, but it is necessary to give complete details on the formulae in  $n$  dimensions since they differ in numerous ways from the two-dimensional case.

**1.3. Determining Taylor coefficients from wave invariants.** To prove Theorem 1, it is only necessary to determine the Taylor coefficients of the defining function  $f = f_+ = -f_-$  of  $\Omega$  at the endpoints of a symmetric bouncing ball orbit  $\gamma$  from the Balian-Bloch invariants given in Theorem 2 for iterates of this orbit. This is done in §7.6. The proof builds on the methods of [Z3, H].

In fact, our method could be extended to show that analytic domains with fewer symmetries are spectrally determined as in [Z3, H], but for the sake of brevity we do not prove that here.

**1.4. Discussion and Comparison of Methods.** We obtain the formulae for the wave invariants by applying the stationary phase method to the trace of a well-constructed parametrix for the monodromy operator. A secondary purpose of this article is to connect the very conceptual but somewhat abstract monodromy method of [ISZ, SZ] with the methods of [Z1, Z2, Z3]. The articles [Z1, Z2] implicitly used the monodromy approach in the form given in [BBa, L, LT]. In this article, we construct the monodromy operator explicitly in terms of layer potentials, using in part the methods of [CP] and in part those of [Z3, HZ]. The monodromy approach connects nicely with the ‘Balian-Bloch’ approach of [Z3] and simplifies remainder estimates for the Balian-Bloch (i.e. Neumann) expansion of the resolvent.

In calculating the trace asymptotics, we do not put the monodromy operator into normal form, but rather apply a direct stationary phase analysis to the parametrix. Terms of the stationary phase expansion correspond to Feynman diagrams and the main idea (as in [Z3]) is to isolate the diagrams which are necessary and sufficient to determine the Taylor coefficients of the boundary defining function at the endpoints of  $\gamma$  from the wave trace invariants of iterates of  $\gamma$ . In this  $(\mathbb{Z}/2\mathbb{Z})^n$  symmetric case, there is a unique such diagram and that is why the symmetries simplify the problem. It is an interesting but difficult problem to ‘invert’ the spectrum when some or all of the symmetries are absent.

An alternative to the approach of this article is to use quantum Birkhoff normal forms around the bouncing ball orbit as in [G, G2, Z1, Z2, ISZ, SZ]. It suffices to prove the abstract result that the quantum normal form of the Laplacian or wave group at the invariant bouncing ball orbit, hence that the classical Birkhoff normal form of the Poincaré map, is a spectral invariant. At that point, one could generalize the result of Y. Colin de Verdière [CV] that the classical normal form determines the Taylor coefficients of  $f$  at the endpoints of the bouncing ball orbit when  $f$  has the  $(\mathbb{Z}/2\mathbb{Z})^n$  symmetries. We plan to carry out the details in a follow-up to this article. The methods of this article go further, since Theorem 2 determines much more than the classical Birkhoff normal form.

It would be interesting to obtain a more direct connection between the ‘normal forms’ approach and the ‘parametrix approach’. In general terms, normal forms for Hamiltonians and for canonical transformations belong to the canonical Hamiltonian formulation of quantum mechanics, while parametrix constructions, stationary phase methods and Feynman diagrams belong to the Lagrangian or path integral approach. Normal forms are of course canonical, while parametrices are not: there are many possible parametrices (finite dimensional approximations to path integrals), and in the inverse problem it is essential to construct computable ones. The two approaches are dual, and although they contain the same information, it is formatted in different ways. In particular, the two approaches highlight different features of the geometry and dynamics.

At the present time, explicit calculations and spectral inversion for boundary problems have only been carried out in the Lagrangian approach, despite the existence of a quantum normal form along bouncing ball orbits [Z2]. In the simpler setting of semi-classical Schrödinger operators at equilibrium points, one may compare the normal forms approach of [GU, CVG] to the Lagrangian approach of [H]. In this inverse problem, one has a one-parameter family of isospectral operators depending on a Planck’s constant  $h$ , whereas in the boundary problem one has only one operator and spectrum to work with. The Lagrangian calculations in [H] reproduced the inverse results of [GU, CVG], and gave stronger ones where some of the symmetries were removed. It directly gives formulae for wave invariants, which are linear combinations of normal form invariants.

It is interesting to observe that formula in Theorem 2 is very similar to the formula in [H] (Theorem 2.1) for the wave invariants at an equilibrium point for a Schrödinger operator on  $\mathbb{R}^n$  with a unique equilibrium point at  $(x, \xi) = (0, 0)$ . This perhaps indicates a similarity between the quantum normal form of the Schrödinger operator at the equilibrium point and that of the Laplacian at a bouncing ball orbit. The formula is also similar to a trace asymptotics formula of T. Christiansen for an inverse problem for wave-guides [Chr], but that is less surprising.

The methods of this paper have further applications. In a work in progress [HeZ], we use the wave invariants to prove a certain spectral rigidity result for analytic deformations of an ellipse.

Finally, we would like to thank the referees for their suggestions on improving the exposition.

## 2. BACKGROUND

In this section, we go over the basic set-up of the problem. It is very similar to that of [Z3] but requires some higher dimensional generalizations. We use the same notation as in [Z3] and refer there for many details.

**2.1. Billiard map.** The billiard map  $\beta$  is defined on

$$B^*\partial\Omega = \{(y, \eta); y \in \partial\Omega, \eta \in T^*\partial\Omega, |\eta| \leq 1\}$$

as follows: given  $(y, \eta) \in B^*\partial\Omega$ , with  $|\eta| \leq 1$ , let  $(y, \zeta) \in S^*\Omega$  be the unique inward-pointing unit covector at  $y$  which projects to  $(y, \eta)$  under the map  $T_{\partial\Omega}^*\bar{\Omega} \rightarrow T^*\partial\Omega$ . Then follow the geodesic (straight line) determined by  $(y, \zeta)$  to the first place it intersects the boundary again; let  $y' \in \partial\Omega$  denote this first intersection. (If  $|\eta| = 1$ , then we let  $y' = y$ .) Denoting the inward unit normal vector at  $y'$  by  $\nu_{y'}$ , we let  $\zeta' = \zeta + 2(\zeta \cdot \nu_{y'})\nu_{y'}$  be the direction of the geodesic after elastic reflection at  $y'$ , and let  $\eta'$  be the projection of  $\zeta'$  to  $B_y^*\partial\Omega$ . Then we define

$$\beta(y, \eta) = (y', \eta').$$

The billiard map is a symplectic, hence measure preserving, map with respect to the standard symplectic form on  $T^*\partial\Omega$ . We denote its graph of  $\beta$  by

$$(8) \quad C_{\text{billiard}} := \text{graph } \beta \equiv \{(\beta(z), z) \mid z \in B^*\partial\Omega\}.$$

In the case of a convex domain,

$$(9) \quad C_{\text{billiard}} = \Gamma_d := \{(y, -\nabla_y d(y, y'), y', \nabla_{y'} d(y, y'))\},$$

i.e. the Euclidean distance function  $d(y, y')$  is a generating function for  $\beta$ . For non-convex domains, this graph is larger due to ‘ghost’ billiard trajectories which exit and re-enter  $\Omega$  but satisfy the reflection law of equal angles at each intersection point. Such ghost orbits are the price one pays for using a parametrix and distance function  $d(y, y')$  defined on the ambient space  $\mathbb{R}^n$ .

**2.2. Length functional.** We define the length functional on  $(\partial\Omega)^m$  (the Cartesian product), by

$$(10) \quad L(y_1, \dots, y_m) = |y_1 - y_2| + \dots + |y_{m-1} - y_m| + |y_m - y_1|.$$

Then  $L$  is a smooth away from the ‘large diagonals’  $\Delta_{p,p+1} := \{y_p = y_{p+1}\}$ . The condition that  $dL = 0$  is the classical condition that each 2-link defined by the triplet  $(y_{p-1}, y_p, y_{p+1})$  makes equal angles with the normal at  $y_p$ . Hence a smooth critical point corresponds to a closed  $m$ -link billiard trajectory. See for instance §2.1 of [PS].

**2.2.1. Length functional in graph coordinates near the iterates of a bouncing ball orbit.** A bouncing ball orbit  $\gamma$  is a 2-link periodic trajectory of the billiard flow, i.e. a reversible periodic billiard trajectory that bounces back and forth along a line segment orthogonal to the boundary at both endpoints. As in the Introduction we orient  $\Omega$  so that  $\overline{AB}$  lies along the vertical  $x_n$  axis, with  $A = (0, \frac{L}{2}), B = (0, -\frac{L}{2})$ . We parameterize the boundary locally as two graphs  $x_n = f_{\pm}(x')$  over the  $x'$ -hyperplane. Thus, in a small tube  $T_\epsilon(\gamma)$  around  $\overline{AB}$ , the boundary consists of two components, which are graphs of the form  $y = f_+(x')$  near  $A$  and  $y = f_-(x')$  near  $B$ .

We then define the length functionals in Cartesian coordinates for the two possible orientations of the  $r$ th iterate of a bouncing ball orbit by

$$(11) \quad \mathcal{L}_\pm(x'_1, \dots, x'_{2r}) = \sum_{p=1}^{2r} \sqrt{(x'_{p+1} - x'_p)^2 + (f_{w_\pm(p+1)}(x'_{p+1}) - f_{w_\pm(p)}(x'_p))^2}.$$

Here,  $w_\pm : \mathbb{Z}_{2r} \rightarrow \{\pm\}$ , where  $w_+(p)$  (resp.  $w_-(p)$ ) alternates sign starting with  $w_+(1) = +$  (resp.  $w_-(1) = -$ ). Obviously the point  $(x'_1, \dots, x'_{2r}) = (0, \dots, 0)$ , corresponding to the  $r$ -th iteration of the bouncing ball orbit, is a critical point of  $\mathcal{L}_\pm$ .

**2.3. Resolvent and Wave group.** By the Dirichlet Laplacian  $\Delta_\Omega^D$  we mean the Laplacian  $\Delta = \sum_{j=1}^n \frac{\partial^2}{\partial x_j^2}$  with domain  $\{u \in H_0^1(\Omega) : \Delta u \in L^2\}$ ; thus, in our notation,  $\Delta_\Omega$  is a negative operator. The resolvent  $R_\Omega^D(k)$  of the Laplacian  $\Delta_\Omega^D$  on  $\Omega$  with Dirichlet boundary conditions is the family of operators on  $L^2(\Omega)$  defined for  $k \in \mathbb{C}$ ,  $\Im k > 0$  by

$$R_\Omega^D(k) = -(\Delta_\Omega^D + k^2)^{-1}, \quad \Im k > 0.$$

The resolvent kernel, which we refer to as the *Dirichlet Green's function*  $G_\Omega^D(k, x, y)$  of  $\Omega \subset \mathbb{R}^n$ , is by definition the solution of the boundary problem:

$$(12) \quad \begin{cases} (\Delta_x + k^2)G_\Omega^D(k, x, y) = -\delta(x - y), & (x, y \in \Omega) \\ G_\Omega^D(k, x, y) = 0, & x \in \partial\Omega. \end{cases}$$

The discussion is similar for Neumann boundary conditions except that its domain is  $\{u \in H^2(\Omega) : \partial_\nu u|_{\partial\Omega} = 0\}$ , and  $\partial_\nu G_\Omega^N(k, x, y) = 0$  for  $x \in \partial\Omega$ , where  $\partial_\nu$  is the interior unit normal. As in the introduction, we superscript functions of the Laplacian with  $B$  to denote either boundary condition.

The resolvent with either boundary condition may be expressed in terms of the even wave operator  $E_\Omega^B(t) = \cos(t\sqrt{-\Delta_\Omega^B})$  as

$$(13) \quad R_\Omega^B(k) = \frac{i}{k} \int_0^\infty e^{ikt} E_\Omega^B(t) dt, \quad (\Im k > 0)$$

In this paper we will consider the resolvent  $R_\Omega^B(k)$  along the logarithmic ray  $k = \lambda + i\tau \log \lambda$ , where  $\lambda > 1$  and  $\tau \in \mathbb{R}^+$ .

Given  $\hat{\rho} \in C_0^\infty(\mathbb{R}^+)$ , we define the smoothed resolvent  $R_{\Omega, \rho}^B(k)$  by

$$(14) \quad R_{\Omega, \rho}^B(k) := \rho * (\mu R_\Omega^B(\mu)) = \int_{\mathbb{R}} \rho(k - \mu) (\mu R_\Omega^B(\mu)) d\mu.$$

We note that  $\rho(k - \mu)$  is well-defined since  $\rho$  is an entire function. Let us discuss in what sense the integral in (14) is defined. We notice that since  $\mu \in \mathbb{R}$ , we have defined the resolvent  $R_\Omega^B(\mu)$  by  $R_\Omega^B(\mu + i0^+)$ . Hence we can write

$$\begin{aligned} R_{\Omega, \rho}^B(k) &= \lim_{\epsilon \rightarrow 0^+} \int_{\mathbb{R}} \rho(k - \mu) \mu R_\Omega^B(\mu + i\epsilon) d\mu \\ &= -\lim_{\epsilon \rightarrow 0^+} \int_{k-\mathbb{R}} \rho(\mu') (k - \mu') R_\Omega^B(k + i\epsilon - \mu') d\mu' \\ &= \int_{\mathbb{R}} \rho(\mu) (k - \mu) R_\Omega^B(k - \mu) d\mu, \end{aligned}$$



where the last equality is obtained by taking an appropriate contour. Thus we can also take this last integral as our definition of the smoothed resolvent.

Now by (13) we can rewrite  $R_{\Omega,\rho}^B(k)$  in terms of the wave operator as:

$$\begin{aligned}
 R_{\Omega,\rho}^B(k) &= i \int_0^\infty \int_{\mathbb{R}} \rho(\mu) e^{i(k-\mu)t} E_\Omega^B(t) dt d\mu \\
 (15) \qquad &= i \int_0^\infty \hat{\rho}(t) e^{ikt} E_\Omega^B(t) dt \\
 &= \frac{i}{2} (\rho(k + \sqrt{-\Delta_\Omega^B}) + \rho(k - \sqrt{-\Delta_\Omega^B})).
 \end{aligned}$$

The Poisson formula for manifolds with boundary [AM, GM, PS] gives the existence of a singularity expansion for the trace of  $E_\Omega^B(t)$  near a transversal reflecting ray. If we substitute this singularity expansion into the trace of (15) we obtain an asymptotic expansion in inverse powers of  $k$  for the smoothed resolvent trace:

**THEOREM 2.1.** [AM, GM, PS] *Assume that  $\gamma$  is a non-degenerate periodic reflecting ray, and let  $\hat{\rho} \in C_0^\infty(L_\gamma - \epsilon, L_\gamma + \epsilon)$ , such that  $\hat{\rho} \equiv 1$  on  $(L_\gamma - \epsilon/2, L_\gamma + \epsilon/2)$  and with no other lengths in its support. Then for  $k = \lambda + i\tau \log \lambda$  with  $\tau \in \mathbb{R}^+$ ,  $Tr R_{\Omega,\rho}^B(k)$  admits a complete asymptotic expansion of the form (4) where*

- $\mathcal{D}_{D,\gamma}(k)$  is the symplectic pre-factor

$$\mathcal{D}_{D,\gamma}(k) = C_0 \epsilon_D(\gamma) \frac{e^{ikL_\gamma} e^{i\frac{\pi}{4}m_\gamma}}{\sqrt{|\det(I - P_\gamma)|}};$$

- $P_\gamma$  is the Poincaré map associated to  $\gamma$ ;
- $\epsilon_D(\gamma)$  is the signed number of intersections of  $\gamma$  with  $\partial\Omega$ ;
- $m_\gamma$  is the Maslov index of  $\gamma$ ;
- $C_0$  is a universal constant (e.g. factors of  $2\pi$ ).

**DEFINITION 2.2.** *The coefficients  $B_{\gamma,j}$  are called the wave trace invariants (or Balian-Bloch invariants) associated to the periodic orbit  $\gamma$ .*

As emphasized above, the discussion in the case of the smoothed Neumann resolvent  $R_{\Omega,\rho}^N(k)$  is essentially the same.

### 3. RESOLVENT AND THE LAYER POTENTIALS

The method of layer potentials ([T] II, §7. 11) solves (12) in terms of the ‘layer potentials’  $G_0(k, x, y), \partial_{\nu_y} G_0(k, x, y) \in \mathcal{D}'(\Omega \times \partial\Omega)$ , where  $\nu$  is the interior unit normal to  $\Omega$ , and  $\partial_\nu = \nu \cdot \nabla$ , and where  $G_0(k, x, y)$  is the ‘free’ Green’s function of  $\mathbb{R}^n$ , i.e. of the kernel of the free resolvent  $-(\Delta_0 + k^2)^{-1}$  of the Laplacian  $\Delta_0$  on  $\mathbb{R}^n$ . A key point, first recognized in [BB1, BB2] and put on a rigorous mathematical basis in [Z3, HZ], is that the layer potentials are semi-classical (i.e. non-homogeneous) Lagrangian distributions in the  $k$  parameter, with additional homogeneous singularities on the diagonal. In effect we wish to make use of the first property and suppress the second. This will be explained in §3.1.

First let us briefly recall the method of layer potentials. The free outgoing Green’s function in dimension  $n$  is given by:

$$(16) \qquad G_0(k, x, y) = \frac{i}{4} k^{n-2} (2\pi k |x - y|)^{-(n-2)/2} \text{Ha}_{n/2-1}^{(1)}(k|x - y|).$$

In general, the Hankel function of index  $\nu$  has the integral representation ([T], Chapter 3, §6)

$$(17) \quad \text{Ha}_\nu^{(1)}(z) = \left(\frac{2}{\pi z}\right)^{1/2} \frac{e^{i(z-\pi\nu/2-\pi/4)}}{\Gamma(\nu+1/2)} \int_0^\infty e^{-s} s^{-1/2} \left(1 - \frac{s}{2iz}\right)^{\nu-1/2} ds.$$

The single and double *layer potentials*, as operators from the boundary  $\partial\Omega$  to the interior  $\Omega$ , are then respectively defined by

$$(18) \quad \begin{cases} \mathcal{S}\ell(k)f(x) = \int_{\partial\Omega} G_0(k, x, y)f(y)dS(y), & x \in \Omega, \\ \mathcal{D}\ell(k)f(x) = \int_{\partial\Omega} 2\partial_{\nu_y}G_0(k, x, y)f(y)dS(y), & x \in \Omega. \end{cases}$$

Similarly, for a function  $f$  on  $\partial\Omega$ , the *boundary layer potentials*  $S(k)$  and  $N(k)$ , as operators from the boundary  $\partial\Omega$  to itself are denoted by:

$$(19) \quad \begin{cases} S(k)f(x) = \int_{\partial\Omega} G_0(k, x, y)f(y)dS(y), & x \in \partial\Omega, \\ N(k)f(x) = \int_{\partial\Omega} 2\partial_{\nu_y}G_0(k, x, y)f(y)dS(y), & x \in \partial\Omega. \end{cases}$$

Given any function  $g$  on  $\mathbb{R}^n \setminus \partial\Omega$  and any  $x \in \partial\Omega$ , we denote by  $g_+(x)$  (resp.  $g_-(x)$ ) the limits of  $g(w)$  as  $w \rightarrow x \in \partial\Omega$  from  $w \in \Omega$  (resp.  $w \in \mathbb{R}^n \setminus \bar{\Omega}$ ). The layer potentials and the boundary layer potentials introduced above are related by the following

$$(20) \quad \begin{aligned} (\mathcal{S}\ell(k)f)_+(x) &= (\mathcal{S}\ell(k)f)_-(x) = S(k)f(x), \\ (\mathcal{D}\ell(k)f)_\pm(x) &= (\pm I + N(k))f(x), \quad (\text{The jump formula}). \end{aligned}$$

Using the above notation we also have the following interesting formula of Fredholm-Neumann

$$(21) \quad R_\Omega^D(k) - R_0(k) = -\mathcal{D}\ell(k)(I + N(k))^{-1}\mathcal{S}\ell(k)^t,$$

where  $R_0(k)$  is the free resolvent. This is true because the kernels of both sides of the equation are solutions to the Helmholtz equation and also the restrictions to the boundary of these kernels are the same by the jump formula (20). The formula follows by the uniqueness of the solutions of the Helmholtz equation.

**3.1. Structure of the operator  $N(k)$ .** We now state the precise sense in which  $N(k)$  is a semi-classical Fourier integral operator quantizing the billiard map of  $\partial\Omega$  when  $k = \lambda + i\tau \log \lambda$  and  $\lambda = \Re k \rightarrow \infty$ . It additionally has homogeneous singularities on the diagonal. The discussion here is adapted from [HZ]. Note that here our semiclassical parameter is  $\frac{1}{k}$  which is a complex parameter.

We denote  $S^*\partial\Omega = \{(y, \eta) \in T^*\partial\Omega; |\eta| = 1\}$  and define the diagonal set

$$(22) \quad \Delta_{S^*\partial\Omega} = \{(z, z); z \in S^*\partial\Omega\} \subset T^*\partial\Omega \times T^*\partial\Omega.$$

**PROPOSITION 3.1.** [HZ] *Assume that  $\Omega$  is a smooth domain. Let  $U$  be any neighborhood of  $\Delta_{S^*\partial\Omega}$ . Then there is a decomposition  $N(k)$  as*

$$N(k) = N_0(k) + N_1(k) + N_2(k),$$

where  $N_0(k)$  is a pseudodifferential operator of order  $-1$ ,  $N_1(k)$  is a semi-classical Fourier integral operator of order zero associated with the canonical relation  $\Gamma_d$  (cf. (9)), and  $N_2(k)$  has operator wavefront set contained in  $U$ .

This proposition is valid whether or not  $\Omega$  is convex, but in the convex case  $\Gamma_d = C_{\text{billiard}}$  is the graph of the billiard map (cf §2.1).

From the integral formula (17), the Hankel function  $\text{Ha}_{n/2-1}^{(1)}(z)$  is conormal at  $z = 0$  and as  $z \rightarrow \infty$ ,

$$(23) \quad a(z) := e^{-iz} \text{Ha}_{n/2-1}^{(1)}(z) \sim z^{-1/2} \sum_{j=0}^{\infty} a_j z^{-j}.$$

The kernel of the single-layer potential then has the form

$$S(k, x, y) = G_0(k, x, y) = Ck^{n-2} (k|x-y|)^{-(n-2)/2} \text{Ha}_{n/2-1}^{(1)}(k|x-y|).$$

For the double layer potential it follows from the identity

$$\frac{d}{dz} \text{Ha}_{\nu}^{(1)}(z) = \frac{\nu}{z} \text{Ha}_{\nu}^{(1)}(z) - \text{Ha}_{\nu+1}^{(1)}(z),$$

that

$$N(k, x, y) = 2 \partial_{\nu_y} G_0(k, x, y) = Ck^{n-1} (k|x-y|)^{-(n-2)/2} \text{Ha}_{n/2}^{(1)}(k|x-y|) \left\langle \frac{x-y}{|x-y|}, \nu_y \right\rangle.$$

We now introduce cutoff functions as in [HZ]. When  $k = \lambda + i\tau \log \lambda$ , we put

$$1 = \varphi_1(|x-y|) + \varphi_0(|k|^{3/4}|x-y|) + \varphi_2(|x-y|, |k|)$$

where  $\varphi_1(t)$  is supported in  $t \geq t_0$  for some sufficiently small  $t_0 > 0$  (see [HZ] for the choice of this  $t_0$ ), and  $\varphi_0(t)$  is equal to 1 for  $t \leq 1$  and equal to 0 for  $t \geq 2$ . (The power 3/4 in  $\varphi_0$  could be replaced by any other power strictly between 1/2 and 1). We define the operator  $N_1(k)$  to be the one with the kernel  $\varphi_1(|x-y|)N(k, x, y)$  which has the form

$$(24) \quad N_1(k, x, y) = Ck^{(n-1)} e^{ik|x-y|} \varphi_1(|x-y|) a_1(k|x-y|) \left\langle \frac{x-y}{|x-y|}, \nu_y \right\rangle,$$

where  $a_1(z) = z^{-\frac{n-2}{2}} a(z)$  has an expansion in inverse powers of  $z$  as  $z \rightarrow \infty$  (see (23)), with leading term  $z^{-(n-1)/2}$ . The operator  $N_1(k)$  is manifestly a semiclassical FIO of order 0 with the phase function  $d(x, y) = |x-y|$ , and thus associated with the canonical relation  $\Gamma_d$  (cf. (9)). The operators  $N_0(k)$  and  $N_2(k)$  are constructed similarly from the cut offs  $\varphi_0$  and  $\varphi_2$  respectively. See [HZ] for the details.

#### 4. MONODROMY OPERATORS AND BOUNDARY INTEGRAL OPERATORS

It simplifies the resolvent trace calculation considerably to reduce it to a boundary trace. In [Z4, Z3] this was done by taking the direct sum  $R_{\Omega, \rho}^D(k) \oplus R_{\Omega^c, \rho}^N(k)$  of the interior Dirichlet and exterior Neumann resolvents, and verifying that

$$\text{Tr}(R_{\Omega, \rho}^D(k) \oplus R_{\Omega^c, \rho}^N(k) - R_{0, \rho}(k)) = \rho * \frac{d}{dk} \log \det(I + N(k)).$$

The construction for the interior Neumann resolvent and exterior Dirichlet resolvent is essentially the same and is omitted for brevity. This relation was stated in the physics literature, and we refer to [Z4] for references. In this article, we take a related but somewhat different method to reduce the trace to the boundary using monodromy operator ideas similar to those of Cardoso-Popov [CP] and of Sjöstrand-Zworski [SZ]. A novel feature is that we relate the monodromy operators to the boundary integral operators  $N_1(k)$  and  $N_0(k)$ .

Before going into the details, let us note some motivating ideas. First, both the monodromy operator  $M(k)$  and the boundary integral operator  $N_1(k)$  are quantizations of the billiard map in the sense of being semi-classical Fourier integral operators whose canonical relation is  $C_{\text{billiard}}$ . This suggests that they must be closely related. However,  $M(k)$  is microlocally constructed while  $N_1(k)$  is global. Further,  $N_1(k)$  is just a piece of  $N(k)$ , which has the more complicated structure described in §3.1, and in the boundary reduction  $N(k)$  is the primary object. On the other hand, there is a simple exact formula for  $N(k)$  while  $M(k)$  is only known through microlocal conjugation to normal form. Hence our purpose here is to construct a monodromy operator resembling  $M(k)$  using  $N(k)$  and the layer potentials. In doing so, we follow the approach to monodromy operators of [CP]. Since the resulting monodromy operator does not seem to arise from the abstract set-up of Grushin reductions, the proof that the trace reduces to the boundary is not the same as in [SZ].

**4.1. Definition of the monodromy operator.** In this section we will use the basic terminologies in semi-classical analysis such as semi-classical pseudodifferential operators, Fourier integral operators and semi-classical wave front sets. We refer to [GS, EZ, SZ, Al] for all the definitions and properties. We just mention that in the following sections when we write  $T(k) \sim S(k)$ , for two operators  $T(k)$  and  $S(k)$ , we mean  $T(k) - S(k)$  is a negligible (residual) operator in the sense that its kernel is of order  $O(k^{-\infty})$  in all  $C^s$  norms.

Let  $\gamma$  be an  $m$ -link periodic reflecting ray, with vertices at  $v_j$  in  $\partial\Omega$ , where there are  $m+1$  vertices and  $v_{m+1} = v_1$ . Let  $\eta_j$  be the projection to  $B_{v_j}^* \partial\Omega$  of the direction of the ray  $\gamma$  where it hits the boundary at  $v_j$ . We denote

$$\partial\gamma := \{(v_1, \eta_1), \dots, (v_m, \eta_m)\}$$

Let  $\Gamma_j$  be microlocal neighborhoods of  $(v_j, \eta_j)$  in  $B^* \partial\Omega = \{(y, \eta) | y \in \partial\Omega, \eta \in T^* \partial\Omega, |\eta| \leq 1\}$ . We then define a microlocalization to  $\Gamma_j$  of the double layer potential operator by

$$H_j(k) := \mathcal{D}\ell(k) \Psi_{\Gamma_j}(k) : C(\partial\Omega) \rightarrow C(\Omega),$$

where  $\Psi_{\Gamma_j}(k)$  is a  $k$ -pseudodifferential operator microsupported in  $\Gamma_j$ . In fact we will choose,

$$\Psi_{\Gamma_j}(k) = (I + N_0(k))^{-1} \chi_{\Gamma_j}(k),$$

where  $\chi_{\Gamma_j}$  is a microlocal cut off supported in  $\Gamma_j$ . More precisely  $\chi_{\Gamma_j}(k) = Op_{\frac{1}{k}}(a_j(y, \eta))$ , where  $\text{supp}(a_j(y, \eta)) \subset \Gamma_j$  and  $a_j(y, \eta) = 1$  in a neighborhood of  $(v_j, \eta_j)$ . Here  $(I + N_0(k))^{-1}$  is a parametrix for  $I + N_0(k)$ . Notice a parametrix exists because  $N_0(k)$  is a pseudodifferential of order  $-1$ .

Clearly, by the jump formula (20), for any  $u \in C(\partial\Omega)$ ,

$$\begin{cases} (-\Delta - k^2)H_j(k)u = 0, \\ H_j(k)(u)|_{\partial\Omega} = (I + N(k))\Psi_{\Gamma_j}(k)u. \end{cases}$$

Next we put for every  $1 \leq i \leq m$ ,

(25)

$$P_i(k) = H_i(k) - H_{i+1}(k)r_{\partial\Omega}H_i(k) + \dots + (-1)^{m-1}H_{i+(m-1)}(k)r_{\partial\Omega}H_{i+(m-2)}(k) \cdots r_{\partial\Omega}H_i(k),$$

where  $r_{\partial\Omega} : C(\Omega) \rightarrow C(\partial\Omega)$  is the operator of restriction to the boundary. Also notice by our notation the indices  $j+m$  and  $j$  are identified. Then we define

$$(26) \quad P(k) = \frac{1}{m} \sum_{i=1}^m P_i(k).$$

**PROPOSITION 4.1.** *We have:  $(-\Delta - k^2)P(k)u = 0$ .*

*Proof.* In fact for every  $i$  we have  $(-\Delta - k^2)P_i(k)u = 0$ . We observe that  $P_i(k) = \mathcal{D}\ell(k)Q_i(k)$  with

$$(27) \quad Q_i(k) = \Psi_{\Gamma_i} - \Psi_{\Gamma_{i+2}}[(I + N(k))\Psi_{\Gamma_{i+1}}] + \cdots + (-1)^{m-1}\Psi_{\Gamma_{i+m-1}}[(I + N(k))\Psi_{\Gamma_{i+m-2}} \cdots (I + N(k))\Psi_{\Gamma_i}].$$

The statement follows since  $\mathcal{D}\ell(k)$  maps  $C(\partial\Omega)$  into the solutions of the Helmholtz equation.  $\square$

We now make a couple of useful technical observations:

**PROPOSITION 4.2.** *We have:*

$$r_{\partial\Omega} H_j(k) \sim (I + N_0(k) + N_1(k))\Psi_{\Gamma_j}.$$

*Proof.* By the jump formula we know that  $r_{\partial\Omega} H_j(k) = (I + N(k))\Psi_{\Gamma_j}$ . The missing term  $N_2$  has its wave front set contained in  $U \times U$  where  $U$  is a small neighborhood  $S^*\partial\Omega$ . But then  $WF'(N_2) \circ WF'(\Psi_{\Gamma_j}) = \emptyset$ . So this term of the composition is negligible in  $k$ .  $\square$

**PROPOSITION 4.3.** *For all  $1 \leq j < m$ , we have:*

$$\Psi_{\Gamma_{j+1}}[(I + N(k))\Psi_{\Gamma_j} \cdots (I + N(k))\Psi_{\Gamma_1}] \sim \Psi_{\Gamma_{j+1}}[N_1(k)\Psi_{\Gamma_j} \cdots N_1(k)\Psi_{\Gamma_1}].$$

*Proof.* We argue inductively. For  $j = 1$ , using the proof of Proposition 4.2 we have

$$\Psi_{\Gamma_2}[(I + N(k))\Psi_{\Gamma_1}] \sim \Psi_{\Gamma_2}N_1(k)\Psi_{\Gamma_1} + \Psi_{\Gamma_2}[(I + N_0(k))\Psi_{\Gamma_1}].$$

But the last term in the expression above is negligible in  $k$ , as the two semi-classical pseudodifferentials  $\Psi_{\Gamma_2}$  and  $[(I + N_0(k))\Psi_{\Gamma_1}]$  are microsupported in the two disjoint open sets  $\Gamma_2$  and  $\Gamma_1$  respectively. Now assume the statement is true for  $j - 1$ . Then we write

$$\begin{aligned} \Psi_{\Gamma_{j+1}}[(I + N(k))\Psi_{\Gamma_j} \cdots (I + N(k))\Psi_{\Gamma_1}] &\sim \Psi_{\Gamma_{j+1}}[N_1(k)\Psi_{\Gamma_j} \cdots N_1(k)\Psi_{\Gamma_1}] \\ &\quad + \Psi_{\Gamma_{j+1}}(I + N_0(k))\Psi_{\Gamma_j}[N_1\Psi_{\Gamma_{j-1}} \cdots N_1(k)\Psi_{\Gamma_1}]. \end{aligned}$$

Similarly,  $\Psi_{\Gamma_{j+1}}(I + N_0(k))\Psi_{\Gamma_j}$  is negligible in  $k$  and the second term above is negligible.  $\square$

Next we have the following important proposition which implicitly defines the monodromy operator:

**PROPOSITION 4.4.** *We have*

$$r_{\partial\Omega} P(k) \sim I + M(k), \quad (\text{microlocally near } \partial\gamma)$$

where

$$(28) \quad M(k) = \frac{1}{m} \sum_{i=1}^m (-1)^{m-1} N_1(k)\Psi_{\Gamma_{i+m-1}} \cdots N_1(k)\Psi_{\Gamma_i}.$$

**DEFINITION 4.5.** *The monodromy operator for the  $m$ -link periodic reflecting ray  $\gamma$  is the operator  $M(k)$  on  $L^2(\partial\Omega)$  defined by (28).*

*Proof.* We define

$$M_i(k) = (-1)^{m-1} N_1(k) \Psi_{\Gamma_{i+m-1}} \cdots N_1(k) \Psi_{\Gamma_i},$$

and we show that for every  $i$

$$(29) \quad r_{\partial\Omega} P_i(k) \sim \chi_{\Gamma_i} + M_i(k), \quad (\text{microlocally near } \partial\gamma).$$

Since

$$\frac{1}{m} \sum_{i=1}^m M_i(k) = M(k), \quad \text{and} \quad \frac{1}{m} \sum_{i=1}^m \chi_{\Gamma_i} = I \quad (\text{microlocally near } \partial\gamma),$$

taking averages of equations (29) over all  $i$  implies Proposition 4.4. So we only prove (29) for  $i = 1$ . Since  $P_1(k) = \mathcal{D}\ell(k)Q_1(k)$ , by the jump formula (20) and by (27), we have

$$r_{\partial\Omega} P_1(k) = (I + N(k)) \sum_{j=0}^{m-1} (-1)^j \Psi_{\Gamma_{j+1}} [(I + N(k)) \Psi_{\Gamma_j} \cdots (I + N(k)) \Psi_{\Gamma_1}],$$

where the 0-th term of the sum is defined to be  $\Psi_{\Gamma_1}$ . Now we apply Proposition 4.3 to each term of the above sum and we get

$$r_{\partial\Omega} P_1(k) \sim (I + N(k)) \sum_{j=0}^{m-1} (-1)^j \Psi_{\Gamma_{j+1}} [N_1(k) \Psi_{\Gamma_j} \cdots N_1(k) \Psi_{\Gamma_1}].$$

Hence by substituting  $I + N = (I + N_0) + N_1$  and collecting the terms corresponding to the same number of iterations of the billiard map  $\beta$  (this number is the same as the number of factors  $N_1$  in each term) we obtain

$$1_{\partial\Omega} P_1(k) \sim \sum_{j=1}^{m-1} (-1)^{j-1} (I - \chi_{\Gamma_{j+1}}) [N_1(k) \Psi_{\Gamma_j} \cdots N_1(k) \Psi_{\Gamma_1}] + (\chi_{\Gamma_1} + M_1(k)).$$

The  $j$ -th term of the sum above is a  $k$ -FIO corresponding to  $\beta^j$ . We show that each of these terms is microlocally equivalent to 0 near  $\partial\gamma$ . Hence only  $\chi_{\Gamma_1}$  corresponding to  $\beta^0$  and  $M_1(k)$  corresponding to  $\beta^m$  survive and the proposition follows.

Let us discuss why

$$(I - \chi_{\Gamma_{j+1}}) [N_1(k) \Psi_{\Gamma_j} \cdots N_1(k) \Psi_{\Gamma_1}] \sim 0. \quad (\text{microlocally near } \partial\gamma)$$

This is true because

$$WF'(N_1(k) \Psi_{\Gamma_j} \cdots N_1(k) \Psi_{\Gamma_1}) \subset \{(\beta^j(y, \eta), (y, \eta)); (y, \eta) \in \Gamma_1\} \subset \Gamma_{j+1} \times \Gamma_1,$$

and because  $I - \chi_{\Gamma_{j+1}}$  is micro-supported away from  $(v_j, \eta_j)$ . Thus by passing to a smaller microlocal neighborhood of  $\partial\gamma$  in  $B^*\partial\Omega$ , we obtain a negligible operator.  $\square$

#### 4.2. Microlocal parametrix for the interior Dirichlet problem in terms of $M(k)$ .

In this section we construct a microlocal parametrix for the Dirichlet resolvent of  $\Omega$  near  $\gamma$  in terms of the monodromy operator  $M(k)$ . It is a microlocal version of the global formula (21) of Fredholm-Neumann. The discussion in the Neumann case is similar.

**PROPOSITION 4.6.** *Microlocally in a neighborhood of  $\gamma \times \gamma \subset T^*\Omega \times T^*\Omega$ , we have*

$$R_\Omega^D(k) - R_0(k) \sim -P(k)(I + M(k))^{-1}\mathcal{S}\ell(k)^t.$$

*Proof.* Let us look at the Schwartz kernels of the both hand sides in a microlocal neighborhood of  $\gamma \times \gamma$ . We show that

$$G_\Omega^D(k, x, y) - G_0(k, x, y) \sim -(P(k)(I + M(k))^{-1})(\mathcal{S}\ell(k)^t(k, x, y)), \quad (\text{microlocally near } \gamma \times \gamma)$$

where the operator  $P(k)(I + M(k))^{-1} : C(\partial\Omega) \rightarrow C(\Omega)$  acts on the first component  $x$  of the kernel  $\mathcal{S}\ell(k)^t(k, x, y)$ . To prove this, since microlocal solutions are unique, it is enough to show that for all  $y$  the right side is a solution of the Dirichlet problem

$$\begin{cases} (-\Delta_x - k^2)(P(k)(I + M(k))^{-1})(\mathcal{S}\ell(k)^t(k, x, y)) = 0, \\ r_{\partial\Omega}(P(k)(I + M(k))^{-1})(\mathcal{S}\ell(k)^t(k, x, y)) \sim -(G_\Omega^D - G_0)(k, x, y), \quad \text{microlocally for } x \text{ near } \partial\gamma. \end{cases}$$

But this is clear because by Proposition 4.4, microlocally for  $x \in \partial\Omega$  near  $\partial\gamma$ , we have for all  $y \in \Omega$

$$\begin{aligned} r_{\partial\Omega}(P(k)(I + M(k))^{-1})(\mathcal{S}\ell(k)^t(k, x, y)) &\sim (I + M(k))(I + M(k))^{-1}\mathcal{S}\ell(k)^t(k, x, y) \\ &\sim -(G_\Omega^D - G_0)(k, x, y). \end{aligned}$$

We note that the operator  $I + M(k)$  is invertible by Lemma 5.5.  $\square$

### 5. TRACE FORMULA AND MONODROMY OPERATORS

Here we assume  $L_\gamma$  is the only length in the support of  $\hat{\rho} \in C_0^\infty$  and  $\hat{\rho}(t) = 1$  near  $L_\gamma$ .

From Proposition 4.6, we immediately have a reduction of the trace to the boundary.

**PROPOSITION 5.1.** *We have:*

$$Tr_\Omega(R_{\Omega,\rho}^D(k) - R_{0,\rho}(k)) \sim -Tr_{\partial\Omega}(\rho * k \mathcal{S}\ell(k)^t \mathcal{D}\ell(k) Q(k) (I + M(k))^{-1}).$$

*Proof.* First of all we note that the regularized trace  $Tr_{\partial\Omega}(R_{\Omega,\rho}^D(k) - R_{0,\rho}(k))$  can be microlocalized to  $\gamma$ , i.e. if  $\chi_\gamma$  is a microlocal cutoff around  $\gamma$ , then

$$Tr_{\partial\Omega}(R_{\Omega,\rho}^D(k) - R_{0,\rho}(k)) \sim Tr_{\partial\Omega}(\chi_\gamma(R_{\Omega,\rho}^D(k) - R_{0,\rho}(k))\chi_\gamma).$$

For a proof of this fact, see [Z4] §3.3. Now by Proposition 4.6, we have

$$\begin{aligned} Tr_\Omega(R_{\Omega,\rho}^D(k) - R_{0,\rho}(k)) &\sim -Tr_\Omega(\rho * k P(k)(I + M(k))^{-1}\mathcal{S}\ell(k)^t) \\ &\sim -Tr_{\partial\Omega}(\rho * k \mathcal{S}\ell(k)^t P(k)(I + M(k))^{-1}). \end{aligned}$$

The formula follows by substituting  $P(k) = \mathcal{D}\ell(k)Q(k)$ .  $\square$

This formula is useful but somewhat unwieldy. As proved in [Z3],  $\mathcal{S}\ell(k)^t \mathcal{D}\ell(k) = D_0 + D_1$  where  $D_0$  is a  $k$ -pseudodifferential operator and where  $D_1$  quantizes  $\beta$ . The same analysis could be used here. But it is simpler to use the alternative in the next section.

**5.1. Interior plus exterior.** If we take the direct sum of the interior Dirichlet and exterior Neumann resolvents, then the trace formula simplifies in that we can sum up the interior and exterior  $\mathcal{S}\ell(k)^t \circ \mathcal{D}\ell(k)$  operators to obtain  $\frac{1}{2k}N'(k) := \frac{1}{2k}(\partial/\partial k)N(k)$ .

**PROPOSITION 5.2.** *We have:*

$$Tr_{\mathbb{R}^n}(R_{\Omega,\rho}^D(k) \oplus R_{\Omega^c,\rho}^N(k) - R_{0,\rho}(k)) = - \int_{\mathbb{R}} \rho(\mu) Tr(N'(k-\mu)Q(k-\mu)(I+M(k-\mu))^{-1})d\mu.$$

*Proof.* We first derive an analogue of Proposition 4.6 for the exterior Neumann problem. We then take the trace of the direct sum of the interior Dirichlet and exterior Neumann resolvents.

We construct a parametrix for the exterior Neumann problem by a modification of the method used for the interior Dirichlet problem. The discussion of the interior was motivated by the double layer representation for the interior Dirichlet Green's function. For the exterior Neumann problem we use the single layer representation of the exterior Neumann Green's function,  $R_{\Omega^c}^N(k) - R_0(k) = -\mathcal{S}\ell(k)(I+N^t(k))^{-1}\mathcal{D}\ell(k)^t$ , where the superscript  $t$  denotes the transpose. This formula is proved by expressing the left side as  $\mathcal{S}\ell(k)\psi$  for some  $\psi$ , taking the normal derivative from the exterior and solving for  $\psi$ . We then consider

$$r_{\Omega^c}(R_{\Omega^c}^N(k) - R_0(k))r_{\Omega^c},$$

where  $r_X$  is the characteristic function of  $X$ . We observe that this operator is symmetric, i.e. equals its transpose. It follows that

$$(r_{\Omega^c}(R_{\Omega^c}^N(k) - R_0(k))r_{\Omega^c})(x, y) = -(r_{\Omega^c}\mathcal{D}\ell(k)(I+N)^{-1}\mathcal{S}\ell(k)^tr_{\Omega^c})(y, x).$$

Therefore, at least on the diagonal, we can use the same parametrix formula in the exterior.

We now complete the proof of Proposition 5.2 by taking the trace on  $\mathbb{R}^n$  of the direct sum of the two operators,

$$(30) \quad \begin{cases} r_{\Omega}(R_{\Omega}^D(k) - R_0(k))r_{\Omega} \sim -r_{\Omega}P(k)(I+M(k))^{-1}\mathcal{S}\ell(k)^tr_{\Omega} & \text{(microlocally near } \gamma) \\ r_{\Omega^c}(R_{\Omega^c}^N(k) - R_0(k))r_{\Omega^c} \sim -r_{\Omega^c}P(k)(I+M(k))^{-1}\mathcal{S}\ell(k)^tr_{\Omega^c} & \text{(microlocally near } \gamma) \end{cases}$$

In taking the trace we may cycle  $\mathcal{S}\ell(k)^tr_{\Omega}$  to the front in the first trace and  $\mathcal{S}\ell(k)^tr_{\Omega^c}$  to the front in the second trace. We then add them to get,

$$(31) \quad Tr_{\mathbb{R}^n}(R_{\Omega}^D(k) \oplus R_{\Omega^c}^N(k) - R_0(k)) \sim -Tr_{\partial\Omega}\mathcal{S}\ell(k)^t\mathcal{D}\ell(k)Q(k)(I+M(k))^{-1},$$

where  $\mathcal{S}\ell(k)^t \circ \mathcal{D}\ell(k)$  has the kernel

$$(32) \quad \int_{\mathbb{R}^n} G_0(k, x, w)\partial_{\nu_y}G_0(k, w, y)dw = \frac{1}{2k}N'(k, x, y).$$

For a proof of this simple fact see equation (19) of [Z5]. This indeed is why the interior Dirichlet and exterior Neumann problems were combined and explains the sense in which they are complementary. We then convolve with  $\rho$ .  $\square$

**REMARK 5.3.** *We notice that here the exterior trace  $Tr(r_{\Omega^c}(R_{\Omega^c,\rho}^N(k) - R_{0,\rho}(k))r_{\Omega^c})$  is negligible in  $k$  and it is added only to simplify the expression in Prop 5.1 to the more convenient expression in Prop 5.2.*



We now use Proposition 5.2 to obtain asymptotics of the trace. The next step is to expand  $(I + M)^{-1}$  in a finite geometric (Neumann) series with remainder. We have

$$(33) \quad (I+M)^{-1} = \sum_{n=0}^{n_0} (-1)^n M^n + (-1)^{n_0+1} M^{n_0+1}(I+M)^{-1}.$$

The following proposition shows that, in calculating a given order of Balian-Bloch invariant  $B_{\gamma,j}$ , we may neglect a sufficiently high remainder of the expansion (33).

**PROPOSITION 5.4.** *Assume that  $k = \lambda + i\tau \log \lambda$ . For each order  $|k|^{-J}$  in the trace expansion there exists  $n_0(J)$  such that*

$$(i) \quad \text{Tr} \int_{\mathbb{R}} \rho(\mu) M(k - \mu)^{n_0(J)+1} (I + M(k - \mu))^{-1} N'(k - \mu) Q(k - \mu) d\mu = O(|k|^{-J-1}),$$

$$(ii) \quad \text{Tr}_{\Omega} R_{\Omega, \rho}^D(k) = \sum_{n=0}^{n_0(J)} (-1)^n \text{Tr} \int_{\mathbb{R}} \rho(\mu) M(k - \mu)^n N'(k - \mu) Q(k - \mu) d\mu + O(|k|^{-J-1}).$$

*Proof.* Part (ii) is easily proved by combining part (i), Proposition 5.2 and (33). It remains to estimate the remainder and show part (i). For this, we need to establish an  $L^2$ - norm estimate for the operator norm of  $M(k - \mu) = M(\lambda - \mu + i\tau \log \lambda)$  for sufficiently large  $\tau$ .

The proof is implied by the following norm estimate, which is analogous to Lemma 6.2 of [SZ]. Let  $t_0$  be the constant in §3.1. We note that the monodromy operator depends on a choice of  $t_0$  although it is not indicated in the notation.

**LEMMA 5.5.** *Let  $k = \lambda + i\tau \log \lambda$ . For every  $a > 0$  there exists  $\tau > 0$  and constants  $b, C > 0$  such that,*

$$\|M(k - \mu)\|_{L^2} \leq C|k|^{-a} < \mu >^b.$$

To prove the Lemma, we observe that by Proposition 4.4,

$$(34) \quad \|M(k - \mu)\| \leq C(|k| < \mu >)^b \|N_1(k - \mu)\|^m,$$

For some integers  $C$  and  $b$ . We will not relabel these constants in the course of our estimates.

We recall (see (24)) that  $N_1(k - \mu)$  has Schwartz kernel

$$(35) \quad C(k - \mu)^{n-1} e^{i(k-\mu)|x-y|} \varphi_1(|x-y|) a_1((k - \mu)|x-y|) \left\langle \frac{x-y}{|x-y|}, \nu_y \right\rangle,$$

where  $\varphi_1(t)$  is supported in  $t \geq t_0$  for some  $t_0 > 0$ . We estimate its norm by the Schur estimate,

$$(36) \quad \begin{aligned} \|N_1(k - \mu)\| &\leq C|k - \mu|^{n-1} \sup_{x \in \partial\Omega} \int_{\partial\Omega} |e^{i(\lambda+i\tau \log \lambda)|x-y|} \varphi_1(|x-y|) a_1((k - \mu)|x-y|)| dS(y) \\ &\leq C|k - \mu|^{n-1} e^{-\tau \log \lambda t_0} \sup_{x \in \partial\Omega} \int_{\partial\Omega} |\varphi_1(|x-y|) a_1((k - \mu)|x-y|)| dS(y) \\ &\leq C < \mu >^{2n} e^{-(\tau t_0 - \epsilon) \log \lambda}, \end{aligned}$$

where we estimate  $|k - \mu|^{n-1} \sup_{x,y \in \partial\Omega, |x-y| \geq t_0} |a_1((k - \mu)|x-y|)| \leq C|k|^{2n} < \mu >^{2n}$ .

Since we can choose any small  $\epsilon > 0$  and also any large  $\tau > 0$ , it is clear that, for any  $a > 0$ , there exist  $\epsilon$  and  $\tau$  such that  $\|M(k - \mu)\| \leq C|k|^{-a} < \mu >^b$ . This proves the Lemma and hence the first part of Proposition 5.4. □

**5.2. Trace for the iterations of a bouncing ball orbit.** We now analyze the trace in part (ii) of Prop 5.4 when it is specialized to the  $r$ th iterate  $\gamma^r$  of a bouncing ball orbit, which has  $m = 2r$  links. We observe that  $Q(k - \mu)$  is a sum of terms quantizing  $\beta^0, \beta^1, \dots, \beta^{2r}$ . Let us write  $q_j(k - \mu)$  for the term quantizing  $\beta^j$ . We note that based on this notation, we have  $q_{2r}(k - \mu) = M(k - \mu)$  where  $M(k - \mu)$  is the monodromy operator for  $\gamma^r$ . It follows that

$$N'(k - \mu) Q(k - \mu)$$

is a sum of terms quantizing  $\beta^0, \dots, \beta^{2r+1}$ . On the other hand if we use the monodromy operator for  $\gamma^r$ , then  $(I + M(k - \mu))^{-1}$  is a sum of terms quantizing  $\beta^0, \beta^{2r}, \beta^{4r}, \dots$ . Therefore only three terms of  $N'(k - \mu) Q(k - \mu) (I + M(k - \mu))^{-1}$  are associated to  $\gamma^{2r}$  and can contribute to the trace:

- (1)  $-N'_0(k - \mu) M(k - \mu)$ ;
- (2)  $N'_0(k - \mu) q_{2r}(k - \mu) = N'_0(k - \mu) M(k - \mu)$ ;
- (3)  $N'_1(k - \mu) q_{2r-1}(k - \mu)$ .

Here, we use that  $N'_0$  is associated to  $\beta^0$  and  $N'_1$  is associated to  $\beta$ . We notice the terms (1) and (2) cancel and hence only the term (3) contributes to the trace. Also we notice that for the  $r$ -th iteration of a bouncing ball orbit we have only two vertices and therefore for  $i$  odd we have  $\Gamma_i = \Gamma_1$  and for  $i$  even we have  $\Gamma_i = \Gamma_2$ . Let us denote by

$$\Gamma_+ = \Gamma_1, \quad \text{and} \quad \Gamma_- = \Gamma_2,$$

the microlocal neighborhoods corresponding to the top and bottom vertex respectively. Thus, by these notations we have

**PROPOSITION 5.6.** *Let  $\hat{\rho} \in C_0^\infty(\mathbb{R})$  be a cut off satisfying  $\text{supp } \hat{\rho} \cap \text{Lsp}(\Omega) = \{rL_\gamma\}$ . Then*

$$\begin{aligned} (37) \quad & \text{Tr}(R_{\Omega, \rho}^D(k)) \sim -\text{Tr} \int_{\mathbb{R}} \rho(\mu) N'_1(k - \mu) q_{2r-1}(k - \mu) d\mu \\ & \sim \frac{1}{2} \sum_{\pm} \text{Tr} \int_{\mathbb{R}} \rho(\mu) N'_1(k - \mu) \\ & \quad \times \overbrace{(N_1(k - \mu) \Psi_{\Gamma_{\mp}})(N_1(k - \mu) \Psi_{\Gamma_{\pm}}) \cdots (N_1(k - \mu) \Psi_{\Gamma_{\mp}})(N_1(k - \mu) \Psi_{\Gamma_{\pm}})}^{(2r-1) \text{ times}} d\mu. \end{aligned}$$

We now express this trace as an explicit oscillatory integral. We consider both principal (we will define the principal terms in the next section) and non-principal terms. All terms arise as composition of  $2r$  Fourier integral operators quantizing  $\beta$ , hence may be expressed as compositions of  $2r$  oscillatory integrals. We recall that  $\Psi_{\Gamma_j} = (I + N_0)^{-1} \chi_{\Gamma_j}$ . Next we expand

$$(I + N_0(k))^{-1} = I + N_{-1}(k),$$

where  $N_{-1}(k)$  is a  $(-1)$ st order pseudo-differential operator. If we plug  $(I + N_{-1}(k)) \chi_{\Gamma_{\pm}}$  for  $\Psi_{\Gamma_{\pm}}$  into the expression (37), after expanding we get

**COROLLARY 5.7.** *Let  $k = \lambda + i\tau \log \lambda$ . Up to  $O(|k|^{-\infty})$ , the trace  $\text{Tr}(R_{\Omega, \rho}^D(k))$  is a sum of  $2r$  oscillatory integrals of the form*

$$I_{2r, \rho}^\sigma(k) = \int_{\mathbb{R}} \int_{\mathbb{R}} \int_{(\partial\Omega)^{2r}} e^{i[\mu(t - \mathcal{L}(y_1, \dots, y_{2r})) + k\mathcal{L}(y_1, \dots, y_{2r})]} A_{2r}^\sigma(k - \mu, y_1, \dots, y_{2r}) \hat{\rho}(t) dt d\mu dS(y_1) \cdots dS(y_r),$$

where the superscript  $\sigma$ ;  $0 \leq \sigma \leq 2r - 1$ , denotes the sum of the terms which contain  $\sigma$  factors of  $N_{-1}$ , and where

$$\mathcal{L}(y_1, \dots, y_{2r}) = |y_1 - y_2| + \cdots + |y_{2r} - y_1|,$$

and  $A_{2r}^\sigma(k - \mu, y_1, \dots, y_{2r}) \in S_\delta^{-|\sigma|}((\partial\Omega)^{2r})$ .

## 6. PRINCIPAL TERMS

The goal of this section is to identify the *principal terms*, which generate the highest derivative data, and to prove that non-principal terms contribute only lower order derivative data.

As in [Z3], we separate out a single oscillatory integral (the principal term  $I_{2r, \rho}^0$ ) which generates all terms of the wave trace (or Balian-Bloch) expansion which contain the maximal number of derivatives of the boundary defining function per power of  $k$  (i.e. order of wave invariant).

**DEFINITION 6.1.** *Let  $\gamma$  be a 2-link periodic orbit, and let  $\gamma^r$  be its  $r$ th iterate. The principal term is the term of (37) in which  $\Psi_{\Gamma_\pm}$  is replaced by  $\chi_{\Gamma_\pm}$ . Thus, the principal term is*

$$I_{2r, \rho}^0 = - \sum_{\pm} \text{Tr} \rho * N_1' \left( \overbrace{N_1 \chi_{\Gamma_\mp} N_1 \chi_{\Gamma_\pm} \cdots N_1 \chi_{\Gamma_\mp} N_1 \chi_{\Gamma_\pm}}^{(2r-1) \text{ times}} \right) d\mu.$$

This oscillatory integral corresponds to  $I_{2r, \rho}^0$ , i.e. the one in 5.7 corresponding to  $\sigma = 0$ . By Corollary 5.7, the oscillatory integral  $I_{2r, \rho}^0$  has the phase function  $\mathcal{L}(y_1, \dots, y_{2r}) = |y_1 - y_2| + \cdots + |y_{2r} - y_1|$ , where  $y_p \in \partial\Omega$ . We may write each  $y_p$  in graph coordinates as  $(x'_p, f_\pm(x'_p))$ . We will use superscripts for the  $n - 1$  components of  $x'_p$ , i.e.  $x'_p = (x_p^1, \dots, x_p^{n-1})$ . Hence the integral is localized to  $[(-\epsilon, \epsilon)^{n-1}]^{2r}$ . We notice that  $I_{2r, \rho}^0$  is the sum of  $I_{2r, \rho}^{0,+}$  and  $I_{2r, \rho}^{0,-}$ , where they correspond to the  $+$  and  $-$  term respectively. It is clear that the phase function of  $I_{2r, \rho}^{0,\pm}$  is given by

$$\mathcal{L}_\pm(x'_1, \dots, x'_{2r}) = \sum_{p=1}^{2r} \sqrt{(x'_{p+1} - x'_p)^2 + (f_{w_\pm(p+1)}(x'_{p+1}) - f_{w_\pm(p)}(x'_p))^2}.$$

Here,  $w_\pm : \mathbb{Z}_{2r} \rightarrow \{\pm\}$ , where  $w_+(p)$  (resp.  $w_-(p)$ ) alternates sign starting with  $w_+(1) = +$  (resp.  $w_-(1) = -$ ).

Now we have the following Theorem 6.3. First we have a definition:

**DEFINITION 6.2.** Let  $\gamma$  be an  $m$ -link periodic reflecting ray, and let  $\hat{\rho} \in C_0^\infty(\mathbb{R})$  be a cut off satisfying  $\text{supp } \hat{\rho} \cap Lsp(\Omega) = \{rL_\gamma\}$  for some fixed  $r \in \mathbb{N}$ . Given an oscillatory integral  $I(k)$ , we write

$$TrR_{\Omega,\rho}^B(k) \equiv I(k) \text{ mod } \mathcal{O}\left(\sum_j k^{-j}(\mathcal{J}^{2j}f)\right)$$

if

$$TrR_{\Omega,\rho}^B(k) - I(k)$$

has a complete asymptotic expansion of the form (4), and if the coefficient of  $k^{-j}$  depends on  $\leq 2j$  derivatives of the defining functions  $f$  at the reflection points.

The following Theorem is the higher dimensional generalization of Theorem 4.2 of [Z3].

**THEOREM 6.3.** Let  $k = \lambda + i\tau \log \lambda$ . Let  $\gamma$  be a primitive non-degenerate 2-link periodic reflecting ray, whose reflection points are points of non-zero curvature of  $\partial\Omega$ , and let  $\hat{\rho} \in C_0^\infty(\mathbb{R})$  be a cut off satisfying  $\text{supp } \hat{\rho} \cap Lsp(\Omega) = \{rL_\gamma\}$  and equals one near  $rL_\gamma$  for some fixed  $r \in \mathbb{N}$ . Orient  $\Omega$  so that  $\gamma$  is the vertical segment  $\{x' = 0\} \cap \Omega$ , and so that  $\partial\Omega$  is a union of two graphs over  $[-\epsilon, \epsilon]^{n-1}$ . Then

$$(1) \quad TrR_{\Omega,\rho}^B(k) \equiv I_{2r,\rho}^0 \text{ mod } \mathcal{O}\left(\sum_j k^{-j}(j^{2j}f_\pm)\right)$$

(2) We also have the following integral representation for  $I_{2r,\rho}^0$  in the  $x'_p$  coordinates

$$(38) \quad I_{2r,\rho}^0 = \sum_{\pm} \int_{([-\epsilon,\epsilon]^{n-1})^{2r}} e^{i(k+i\tau)\mathcal{L}_\pm(x'_1,\dots,x'_{2r})} a_{2r}^{pr,\pm}(k, x'_1, x'_2, \dots, x'_{2r}) dx'_1 \cdots dx'_{2r},$$

where the phase  $\mathcal{L}_\pm(x'_1, \dots, x'_{2r})$  is given in (11), and where the amplitude is given by:

$$a_{2r}^{pr,\pm}(k, x'_1, \dots, x'_{2r}) = \mathcal{L}_{w_\pm}(x'_1, \dots, x'_{2r}) A_{2r}^{pr,\pm}(k, x'_1, \dots, x'_{2r}) + \frac{1}{i} \frac{\partial}{\partial k} A_{2r}^{pr,\pm}(k, x'_1, \dots, x'_{2r}).$$

Here

$$(39) \quad \begin{aligned} A_{2r}^{pr,\pm}(k, x'_1, \dots, x'_{2r}) &= \prod_{p=1}^{2r} \left\{ a_1 \left( k \sqrt{(x'_p - x'_{p+1})^2 + (f_{w_\pm(p)}(x'_p) - f_{w_\pm(p+1)}(x'_{p+1}))^2} \right) \right. \\ &\times \left. \frac{\langle x'_p - x'_{p+1}, \nabla_{x'_p} f_{w_\pm(p)}(x'_p) \rangle - (f_{w_\pm(p)}(x'_p) - f_{w_\pm(p+1)}(x'_{p+1}))}{\sqrt{(x'_p - x'_{p+1})^2 + (f_{w_\pm(p)}(x'_p) - f_{w_\pm(p+1)}(x'_{p+1}))^2}} \right\} \end{aligned}$$

where  $a_1$  is the Hankel amplitude in (24). Here  $w_+(p) = (-1)^{p+1}$  and  $w_-(p) = -w_+(p)$ . Also we have identified  $x'_{2r+1} = x'_1$ .

*Proof.* To prove the first part of Theorem it is enough to show that for a given  $\sigma \geq 1$ , the coefficient of  $k^{-j}$  in the stationary phase expansion of  $I_{2r,\rho}^\sigma(k)$ , has only Taylor coefficients of order at most  $2j - \sigma + 1$ . This is shown in §5.4 of [Z3]. The second part of Theorem follows from the proof of Proposition 3.10 of [Z3]. It is basically just eliminating the variables  $t$  and  $\mu$  in the integral in Corollary (5.7) using the stationary phase lemma.  $\square$

Theorem 6.3 is a crucial ingredient in the proof of Theorem 1. It gives explicit formula for the phase and amplitude of the principal oscillatory integrals that determine the highest order jet of  $\Omega$  in each wave invariant. The notation  $A_{2r}^{pr,\pm}, a_{2r}^{pr,\pm}$  refers to the amplitude of the principal terms of the  $2r$ th integral; these amplitudes contain terms of all orders in  $k$  and principal here does not refer to the principal symbol, i.e. the leading order term in the semi-classical expansion.

## 7. STATIONARY PHASE CALCULATIONS OF $I_{2r,\rho}^0$ AND THE WAVE INVARIANTS

It is easy to see that (see Proposition 4.4 of [Z3]) we have  $I_{2r,\rho}^{0,+} = I_{2r,\rho}^{0,-}$  and therefore  $I_{2r,\rho}^0 = 2I_{2r,\rho}^{0,+}$ . Hence it suffices to consider the  $+$  term. The oscillatory integrals  $I_{2r,\rho}^{0,+}$  have the form (38) with the phase  $\mathcal{L}_+$  and the amplitude  $a_{2r}^{pr,+}$ .

The only critical point occurs when  $x'_p = 0$  for all  $p$ . We denote by  $\text{Hess } \mathcal{L}_\pm(0)$  the  $2r(n-1) \times 2r(n-1)$  matrix with components

$$(40) \quad \text{Hess } \mathcal{L}_\pm(0) = \left( \frac{\partial^2 \mathcal{L}_\pm}{\partial x_p^i \partial x_q^j} \right), \quad i, j = 1, \dots, n-1; \quad p, q = 1, \dots, 2r.$$

It is the Hessian of  $\mathcal{L}_\pm$  at its critical point  $(x'_1, \dots, x'_{2r}) = 0$  in Cartesian coordinates.

We denote by  $\mathcal{H}_+$  the inverse Hessian operator in the variables  $(x'_1, \dots, x'_{2r})$  at this critical point. That is  $\mathcal{H}_+ = \langle \text{Hess}(\mathcal{L}_+)^{-1}(0)D, D \rangle$ , where  $D$  is short for  $(\frac{\partial}{\partial x_1^1}, \dots, \frac{\partial}{\partial x_{2r}^{n-1}})$ . More precisely,

$$(41) \quad \mathcal{H}_+ = \sum_{p,q=1}^{2r} \sum_{i,j=1}^{n-1} h^{(i,j),(p,q)} \left( \frac{\partial^2}{\partial x_p^i \partial x_q^j} \right).$$

Before we apply the Stationary Phase Lemma, in two subsections we state some properties of the inverse Hessian matrix of  $\mathcal{L}_+$ , and also some properties of the phase function  $\mathcal{L}_+$  and principal amplitude  $a_{2r}^{pr,+}$  which may be derived directly from the formula in Theorem 6.3.

**7.1. Properties of Hess  $(\mathcal{L}_+)^{-1}$ .** Let  $\{\nu_{j,\pm}\}_{j=1}^{n-1}$  denote the eigenvalues of the second fundamental form of  $\partial\Omega$  at the endpoints of the bouncing ball orbit. Without loss of generality we can assume

$$(42) \quad \nu_{j,\pm} = D_{x^j}^2 f_\pm(0), \quad j = 1, \dots, n-1.$$

This is because by an orthogonal change of variable (i.e. an isometry of the plane) we can make  $\text{Hess } f_\pm$  a diagonal matrix. Of course when all the symmetry assumptions are satisfied, then  $\text{Hess } f_\pm$  is automatically diagonal.

The following generalizes Proposition 2.2 of [Z3].

**PROPOSITION 7.1.** *Put  $a_{j,\pm} = -2(1 \pm L\nu_{j,\pm})$ , and let  $A_\pm = \text{Diag}(a_{j,\pm})$  be the  $(n-1) \times (n-1)$  diagonal matrix with the diagonal entries  $a_{j,\pm}$ . Then the Hessian  $H_{2r}$  of  $\mathcal{L}_+$  at  $x' = 0$  has the*

form:

$$H_{2r} = \frac{-1}{L} \left\{ \begin{array}{cccccc} A_+ & I & 0 & \dots & I & \\ I & A_- & I & \dots & 0 & \\ 0 & I & A_+ & I & 0 & \\ \dots & \dots & \dots & \dots & \dots & \\ I & 0 & 0 & \dots & A_- & \end{array} \right\},$$

where there are  $2r \times 2r$  blocks and each block is of size  $(n-1) \times (n-1)$ .

*Proof.* There are  $2r$  sets of variables  $x'_p$  and therefore there are  $2r \times 2r$  blocks and the  $(p, q)$ -th block is given by  $D^2_{x'_p x'_q} \mathcal{L}_+(0)$ . We have:

$$(43) \quad \nabla_{x'_p} \mathcal{L}_\pm = \frac{(x'_p - x'_{p+1}) + (f_{w_\pm(p)}(x'_p) - f_{w_\pm(p+1)}(x'_{p+1})) \nabla_{x'_p} f_{w_\pm(p)}(x'_p)}{\sqrt{(x'_p - x'_{p+1})^2 + (f_{w_\pm(p)}(x'_p) - f_{w_\pm(p+1)}(x'_{p+1}))^2}} - \frac{(x'_{p-1} - x'_p) + (f_{w_\pm(p-1)}(x'_{p-1}) - f_{w_\pm(p)}(x'_p)) \nabla_{x'_p} f_{w_\pm(p)}(x'_p)}{\sqrt{(x'_p - x'_{p-1})^2 + (f_{w_\pm(p)}(x'_p) - f_{w_\pm(p-1)}(x'_{p-1}))^2}}.$$

A simple calculation (using (43)) shows that all the blocks  $D^2_{x'_p x'_q} \mathcal{L}_+(0)$  are zero except the ones with  $p = q, p = q + 1$  and  $q = p + 1$ . From (43) we obtain

$$\begin{cases} D^2_{x'_p x'_p} \mathcal{L}_+(0) = 2\left(\frac{1}{L} I + w_+(p) \text{Hess } f_{w_+(p)}(0)\right) = \frac{-1}{L} A_{w_+(p)}, \\ D^2_{x'_p x'_{p+1}} \mathcal{L}_+(0) = \frac{-1}{L} I. \end{cases}$$

□

In the elliptic case,  $\det H_{2r}$  is a polynomial in  $\cos \alpha_j/2$  (in  $\cosh \alpha_j/2$  in the hyperbolic case) of degree  $2r(n-1)$ . Here, in the elliptic case,  $\{e^{\pm i\alpha_1}, \dots, e^{\pm i\alpha_{n-1}}\}$  are the eigenvalues of the Poincaré map  $P_\gamma$ .

**PROPOSITION 7.2.** *We have*

$$\det H_{2r} = L^{2r(1-n)} \prod_{j=1}^{n-1} (2 - 2 \cos r\alpha_j).$$

We will use Proposition 7.3 in the following subsection to prove Proposition 7.2.

**7.1.1. Poincaré map and Hessian of the length functional.** The linear Poincaré map  $P_\gamma$  of  $\gamma$  is the derivative at  $\gamma(0)$  of the first return map to a transversal to  $\Phi^t$  at  $\gamma(0)$ . By a non-degenerate periodic reflecting ray  $\gamma$ , we mean one whose linear Poincaré map  $P_\gamma$  has no eigenvalue equal to one. For the definitions and background, we refer to [PS, KT].

There is an important relation between the spectrum of the Poincaré map  $P_\gamma$  of a periodic  $m$ -link reflecting ray and the Hessian  $H_m$  of the length functional at the corresponding critical point of  $L : (\partial\Omega)^m \rightarrow \mathbb{R}$ . For the following, see [KT] (Theorem 3).

**PROPOSITION 7.3.** *Let  $\gamma$  be a periodic  $m$ -link reflecting ray in plane domain  $\Omega$ . Then we have:*

$$\det(I - P_\gamma) = \det(H_m) \left( \prod_{p=1}^m H_{p,p+1} \right)^{-1},$$

where  $H_{p,p+1}$  is the  $(p, p+1)$ -th entry of  $H_m$ .

Proposition 7.3 is stated only for the plane domains. One can probably prove it for higher dimensions, but the formulae for the plane domains is enough for us to prove Prop.7.2

*Proof of Prop.7.2.* Let us first assume  $n = 2$ . Let  $\lambda_r, \lambda_r^{-1}$  be the eigenvalues of  $P_{\gamma r}$ , so that  $\det(I - P_{\gamma r}) = 2 - (\lambda_r + \lambda_r^{-1})$ . Since in our case  $H_{p,p+1} = -1/L$ , from Prop.7.3 it follows that

$$(44) \quad \det(I - P_{\gamma r}) = L^{2r} \det H_{2r}; \quad (\gamma \text{ 2-link.})$$

This is because if the eigenvalues of  $P_\gamma$  are  $\{e^{\pm i\alpha}\}$  (say, in the elliptic case) then those of  $P_{\gamma r}$  are  $\{e^{\pm ir\alpha}\}$ , hence the left side of (44) equals  $2 - 2\cos r\alpha$ . Now assume  $n \geq 2$  and assume  $\{e^{\pm i\alpha_1}, \dots, e^{\pm i\alpha_{n-1}}\}$  are the eigenvalues of  $P_\gamma$ . We just showed that if we define

$$H_{j,2r} = \frac{-1}{L} \left\{ \begin{array}{ccccc} a_{j,+} & 1 & 0 & \dots & 1 \\ 1 & a_{j,-} & 1 & \dots & 0 \\ 0 & 1 & a_{j,+} & 1 & 0 \\ \dots & \dots & \dots & \dots & \dots \\ 1 & 0 & 0 & \dots & a_{j,-} \end{array} \right\}_{2r \times 2r},$$

then  $\det H_{j,2r} = L^{-2r}(2 - 2\cos(r\alpha_j))$ . Now we notice because all the blocks of the matrix  $H_{2r}$  are diagonal matrices, therefore they commute and we can write

$$\det H_{2r} = \det(\text{Diag}(\det H_{1,2r}, \dots, \det H_{n-1,2r})) = L^{2r(1-n)} \prod_{j=1}^{n-1} (2 - 2\cos r\alpha_j).$$

We now consider the inverse Hessian  $\mathcal{H}_+ = H_{2r}^{-1}$ , which will be important in the calculation of wave invariants. We denote its matrix elements by  $h_+^{ij,pq}$  which corresponds to the  $(i, j)$ -th entry of the  $(p, q)$ -th block of the matrix  $\mathcal{H}_+$ . We also denote by  $\mathcal{H}_-$  the matrix in which the roles of  $A_+, A_-$  are interchanged; it is the Hessian of  $\mathcal{L}_-$ . We also notice since  $H_{2r}$  is a block matrix with each block a diagonal matrix so is its inverse  $\mathcal{H}_+$ . Hence the only non-zero entries of the inverse Hessian  $\mathcal{H}_+$  are of the form  $h_+^{ii,pq}$ .

**PROPOSITION 7.4.** *The diagonal matrix elements  $h_+^{ii,pp}$  are constant when the parity of  $p$  is fixed, and for every  $1 \leq i \leq n-1$  we have:*

$$\begin{aligned} p \text{ odd} &\implies h_\pm^{ii,pp} = h_\pm^{ii,11}, & p \text{ even} &\implies h_\pm^{ii,pp} = h_\pm^{ii,22} \\ h_+^{ii,11} &= h_-^{ii,22}, & h_+^{ii,22} &= h_-^{ii,11} \end{aligned} .$$

*Proof.* It is enough to show this for  $n = 2$ . This is because  $H_{2r}$  is a block matrix with commuting blocks and each block is diagonal. In fact based on our definition in (7.1.1) we have

$$(45) \quad h^{ii,pq} = (H_{2r})^{ii,pq} = (H_{i,2r})^{pq}.$$

Hence it is enough to prove the proposition for the entries of the inverse of the  $2r \times 2r$  matrix  $H_{i,2r}$ . This reduces the problem to  $n - 1 = 1$ .

Now let us introduce the cyclic shift operator on  $\mathbb{R}^{2r}$  given by  $Pe_j = e_{j+1}$ , where  $\{e_j\}$  is the standard basis, and where  $Pe_{2r} = e_1$ . It is then easy to check that  $P\mathcal{H}_+P^{-1} = \mathcal{H}_-$ , hence that  $P\mathcal{H}_+^{-1}P^{-1} = \mathcal{H}_-^{-1}$ . Since  $P$  is unitary, this says

$$h_-^{pq} = \langle \mathcal{H}_- e_p, e_q \rangle = \langle P\mathcal{H}_+^{-1}P^{-1}e_p, e_q \rangle = \langle \mathcal{H}_+^{-1}P^{-1}e_p, P^{-1}e_q \rangle = h_+^{p-1,q-1}.$$

It follows that the matrix  $\mathcal{H}_\pm$  is invariant under even powers of the shift operator, which shifts the indices  $p \rightarrow p + 2k$  ( $k = 1, \dots, r$ ). Hence, diagonal matrix elements of like parity are equal. □

**7.1.2. Inverse Hessian at  $(\mathbb{Z}/2\mathbb{Z})^n$ -symmetric bouncing ball orbits.** We first observe that in the case of  $(\mathbb{Z}/2\mathbb{Z})^n$  symmetric domains, the  $(2r)(n-1) \times (2r)(n-1)$  Hessian of Proposition 7.1 simplifies to:

$$(46) \quad H_{2r} = \frac{-1}{L} \begin{pmatrix} A & I & 0 & \dots & I \\ I & A & I & \dots & 0 \\ 0 & I & A & I & 0 \\ 0 & 0 & I & A & I \dots \\ \dots & \dots & \dots & \dots & \dots \\ I & 0 & 0 & \dots & A \end{pmatrix}$$

which is a  $2r \times 2r$  block matrix, each block of size  $(n-1) \times (n-1)$ . Here  $A = \text{Diag}(a_j)$  and  $I$  is the rank  $n-1$  identity matrix. The diagonal entries  $a_j$  are given by

$$(47) \quad a_j = 2 \cos \alpha_j/2 = -2(1+L\nu_j) \quad (\text{elliptic case}), \quad a_j = 2 \cosh \alpha_j/2 = -2(1+L\nu_j) \quad (\text{hyperbolic case}).$$

We can express the inverse Hessian matrix elements  $h_{2r}^{ij,pq}$  in terms of Chebychev polynomials  $T_m$ , resp.  $U_m$ , of the first, resp. second, kind. The Chebychev polynomials are defined by:

$$T_m(\cos \theta) = \cos m\theta, \quad U_m(\cos \theta) = \frac{\sin(m+1)\theta}{\sin \theta}.$$

**PROPOSITION 7.5.** [Z3] *Suppose that  $H_{2r}$  is given by (46). Then the matrix elements of  $H_{2r}^{-1}$  are given by*

$$h_{2r}^{ii,pq} = \frac{1}{2[1-T_{2r}(-a_i/2)]} [U_{2r-q+p-1}(-a_i/2) + U_{q-p-1}(-a_i/2)], \quad (1 \leq p \leq q \leq 2r; \quad 1 \leq i \leq n-1)$$

$$h_{2r}^{ij,pq} = 0, \quad i \neq j.$$



*Proof.* This formula was proved in [Z3] for the case  $n = 2$ . For the general case  $n \geq 2$ , we just use the equation (45) and reduce it to  $n = 2$ .  $\square$

We note that  $h^{ij,pq} = h^{ij,qp}$  so this formula determines all of the matrix elements. It follows in the elliptic case that

$$(48) \quad h_{2r}^{ii,pq} = \begin{cases} \frac{(-1)^{p-q}}{2(1-\cos r\alpha_i)} \left( \frac{\sin(2r-q+p)\alpha_i/2}{\sin \alpha_i/2} + \frac{\sin(q-p)\alpha_i/2}{\sin \alpha_i/2} \right) & (1 \leq p \leq q \leq 2r) \\ \frac{(-1)^{p-q}}{2(1-\cos r\alpha_i)} \left( \frac{\sin(2r-p+q)\alpha_i/2}{\sin \alpha_i/2} + \frac{\sin(p-q)\alpha_i/2}{\sin \alpha_i/2} \right) & (1 \leq q \leq p \leq 2r) \end{cases}$$

There are obvious analogues in the hyperbolic and mixed cases.

The case of interest to us is

$$(49) \quad h_{2r}^{ii,11} = \frac{1}{2(1-\cos r\alpha_i)} \frac{\sin r\alpha_i}{\sin \alpha_i/2} = \frac{1}{\sin \alpha_i/2} \cot \frac{r\alpha_i}{2}.$$

**7.2. Properties of the phase function  $\mathcal{L}_+$  and the amplitude  $a_{2r}^{pr,+}$ .** Since  $\mathcal{L}_+$  and  $a_{2r}^{pr,+}$  are functions of  $2r(n-1)$  variables  $x_p^j$  where  $1 \leq p \leq 2r$  and  $1 \leq j \leq n-1$ , to simplify our notations we denote:

Let  $[\gamma] = (\gamma_p^j)$ ,  $1 \leq p \leq 2r$ ,  $1 \leq j \leq n-1$  be a  $2r \times (n-1)$  matrix of indices. We let  $m = |[\gamma]| = \sum_{p,j} \gamma_p^j$ . Then we define

$$D_{[\gamma]}^m = \frac{\partial^m}{(\partial x_1^1)^{\gamma_1^1} \dots (\partial x_{2r}^{n-1})^{\gamma_{2r}^{n-1}}}.$$

We will use  $\vec{\gamma}_p$  for the  $p$ -th row of  $[\gamma]$ , and sometimes if  $\vec{\gamma}_p, \vec{\gamma}_q, \dots$  are the only non-zero rows of  $[\gamma]$  we write  $D_{\vec{\gamma}_p, \vec{\gamma}_q, \dots}^m$  for  $D_{[\gamma]}^m$  to emphasize that  $\vec{\gamma}_p, \vec{\gamma}_q, \dots$  are the only non-zero rows of  $[\gamma]$ . The calculation of the highest derivative terms of the Balian-Bloch wave invariants uses only the following properties of the phase and principal amplitude which may be derived directly from the formulae in Theorem 6.3.

The following Lemma is the higher dimensional generalization of Lemma 4.5 of [Z3]. It is proved in the same way, and the proof is therefore omitted.

**LEMMA 7.6.** *The phase and principal amplitude of the principal oscillatory integrals  $I_{2r,\rho}^{0,\pm}$  have the following properties:*

1. *In its dependence on the boundary defining functions  $f_{\pm}$ , the amplitude  $a_{2r}^{pr,+}$  has the form  $\alpha_r(k, x', f_{\pm}, f'_{\pm})$ .*
2. *As above, in its dependence on  $x'$*

$$a_{2r}^{pr,+}(k, x'_1, \dots, x'_{2r}) = \mathcal{L}_+(x'_1, \dots, x'_{2r}) A_{2r}^{pr,+}(k, x'_1, \dots, x'_{2r}) + \frac{1}{i} \frac{\partial}{\partial k} A_{2r}^{pr,+}(k, x'_1, \dots, x'_{2r}),$$

where

$$A_{2r}^{pr,+}(k, x'_1, \dots, x'_{2r}) = \Pi_{p=1}^{2r} A_p(x'_p, x'_{p+1}), \quad 2r+1 \equiv 1. \quad (\text{see (39)})$$

3. *At the critical point, the principal amplitude has the asymptotics*

$$a_{2r}^{pr,+}(k, 0) \sim (2rL)L^{-(n-1)r} \mathcal{A}(r) + O(k^{-1}),$$

where  $\mathcal{A}(r)$  depends only on  $r$  and not on  $\Omega$ .

4.  $\frac{a_{2r}^{pr,+}(k,0)e^{i(k+i\tau)\mathcal{L}_+(0)+i\pi/4\text{sgn Hess}\mathcal{L}_+(0)}}{\sqrt{\det \text{Hess}\mathcal{L}_+}} \sim (2rL) \mathcal{A}(r) \mathcal{D}_{D,\gamma^r}(k)(1 + O(k^{-1}))$  (cf.4).
5.  $\nabla a_{2r}^{pr,+}(k, x'_1, \dots, x'_{2r})|_{x'=0} = 0$ .
6.  $D_{\vec{\gamma}_p}^{2j+1}\mathcal{L}_+|_{x'=0} \equiv 2w_+(p)D_{\vec{\gamma}_p}^{2j+1}f_{w_+(p)}(0) \pmod{R_{2r}(\mathcal{J}^{2j}f_+(0), \mathcal{J}^{2j}f_-(0))}$ .
7.  $D_{\vec{\gamma}_p}^{2j+2}\mathcal{L}_+|_{x=0} \equiv 2w_+(p)D_{\vec{\gamma}_p}^{2j+2}f_{w_+(p)}(0) \pmod{R_{2r}(\mathcal{J}^{2j}f_+(0), \mathcal{J}^{2j}f_-(0))}$ .
8. If  $[\gamma]$  has more than one non-zero row, say  $\vec{\gamma}_p, \vec{\gamma}_q, \dots$ , then

$$D_{\vec{\gamma}_p, \vec{\gamma}_q, \dots}^{2j+1}\mathcal{L}_+(0) \equiv 0 \pmod{R_{2r}(\mathcal{J}^{2j}f_+(0), \mathcal{J}^{2j}f_-(0))},$$

and

$$D_{\vec{\gamma}_p, \vec{\gamma}_q, \dots}^{2j+2}\mathcal{L}_+(0) \equiv 0 \pmod{R_{2r}(\mathcal{J}^{2j}f_+(0), \mathcal{J}^{2j}f_-(0))}.$$

Above,  $\equiv$  means equality modulo lower order derivatives of  $f$ .

**7.3. Stationary phase diagrammatics.** We briefly review the stationary phase expansion from the diagrammatic point of view. For more details we refer to [A, E, Z3].

The stationary phase expansion gives an asymptotic expansion for an oscillatory integral

$$Z(k) = \int_{\mathbb{R}^n} a(x)e^{ikS(x)}dx$$

where  $a \in C_0^\infty(\mathbb{R}^n)$  and where  $S$  has a unique non-degenerate critical point in  $\text{supp}(a)$  at  $x = 0$ . Let us write  $H$  for the Hessian of  $S$  at 0. The stationary phase expansion is:

$$Z(k) = \left(\frac{2\pi}{k}\right)^{n/2} \frac{e^{i\pi \text{sgn}(H)/4}}{\sqrt{|\det H|}} e^{ikS(0)} Z_A(k),$$

where

$$Z_A(k) = \sum_{j=0}^{\infty} k^{-j} \sum_{(\Gamma, \ell) \in \mathcal{G}_{V,I}, I-V=j} \frac{I_\ell(\Gamma)}{S(\Gamma)},$$

where  $\mathcal{G}_{V,I}$  consists of labelled graphs  $(\Gamma, \ell)$  with  $V$  closed vertices of valency  $\geq 3$  (each corresponding to the phase), with one open vertex (corresponding to the amplitude), and with  $I$  edges. The function  $\ell$  ‘labels’ each end of each edge of  $\Gamma$  with an index  $j \in \{1, \dots, n\}$ .

Above,  $S(\Gamma)$  denotes the order of the automorphism group of  $\Gamma$ , and  $I_\ell(\Gamma)$  denotes the ‘Feynman amplitude’ associated to  $(\Gamma, \ell)$ . By definition,  $I_\ell(\Gamma)$  is obtained by the following rule: To each edge with end labels  $j, k$  one assigns a factor of  $ih^{jk}$  where  $H^{-1} = (h^{jk})$ . To each closed vertex one assigns a factor of  $i \frac{\partial^\nu S(0)}{\partial x^{i_1} \dots \partial x^{i_\nu}}$  where  $\nu$  is the valency of the vertex and  $i_1, \dots, i_\nu$  at the index labels of the edge ends incident on the vertex. To the open vertex, one assigns the factor  $\frac{\partial^\nu a(0)}{\partial x^{i_1} \dots \partial x^{i_\nu}}$ , where  $\nu$  is its valence. Then  $I_\ell(\Gamma)$  is the product of all these factors. To the empty graph one assigns the amplitude 1. In summing over  $(\Gamma, \ell)$  with a fixed graph  $\Gamma$ , one sums the product of all the factors as the indices run over  $\{1, \dots, n\}$ .

We note that the power of  $k$  in a given term with  $V$  vertices and  $I$  edges equals  $k^{-\chi_{\Gamma'}}$ , where  $\chi_{\Gamma'} = V - I$  equals the Euler characteristic of the graph  $\Gamma'$  defined to be  $\Gamma$  minus the open vertex. We note that there are only finitely many graphs for each  $\chi$  because the valency condition forces  $I \geq 3/2V$ . Thus,  $V \leq 2j, I \leq 3j$ .

**7.4. The stationary phase calculations of  $I_{2r,\rho}^{0,+}$ : The data  $D_{2\vec{\gamma}}^{2j+2} f_{\pm}(0)$ .** In this section we will repeatedly use different parts of Lemma 7.6 without quoting them.

We first claim that in the stationary phase expansion of  $I_{2r,\rho}^{0,+}$ , the data  $D_{2\vec{\gamma}}^{2j+2} f_{\pm}(0)$  appears first in the  $k^{-j}$  term. This is because any labelled graph  $(\Gamma, \ell)$  for which  $I_{\ell}(\Gamma)$  contains the factor  $D_{2\vec{\gamma}}^{2j+2} f_{\pm}(0)$  must have a closed vertex of valency  $\geq 2j + 2$ , or the open vertex must have valency  $\geq 2j + 1$ . The minimal absolute Euler characteristic  $|\chi(\Gamma')|$  in the first case is  $j$ . Since the Euler characteristic is calculated after the open vertex is removed, the minimal absolute Euler characteristic in the second case is  $j + 1$  (there must be at least  $j + 1$  edges.) Hence the latter graphs do not have minimal absolute Euler characteristic. More precisely, we have:

**PROPOSITION 7.7.** *In the stationary phase expansion of  $I_{2r,\rho}^{0,+}$ , the only labelled graph  $(\Gamma, \ell)$  with  $\chi(\Gamma') = V - I = -j$  and  $I_{\ell}(\Gamma)$  containing  $D_{2\vec{\gamma}}^{2j+2} f_{\pm}(0)$  is given by:*

- $\Gamma_{1,j+1} \in \mathcal{G}_{1,j+1}$  (i.e.  $V = 1, I = j + 1$ ). The graph  $\Gamma_{1,j+1}$  has no open vertex, one closed vertex and  $j + 1$  loops at the closed vertex.
- The only labels producing the desired data are those  $\ell_p$ , with  $1 \leq p \leq 2r$  fixed, which labels all endpoints of all edges as  $(i, p)$  where  $1 \leq i \leq n - 1$ . (Notice the label  $(i, p)$  corresponds to the variable  $x_p^i$ .)

In addition, the sum of the Feynman amplitudes corresponding to the labelled graphs  $(\Gamma_{1,j+1}, \ell_p)$  above, for a fixed  $p$ , is

$$\sum_{\ell_p} I_{\ell_p}(\Gamma_{1,j+1}) \equiv (4rL)L^{-(n-1)r} \mathcal{A}(r) i^{j+2} \sum_{|\vec{\gamma}_p|=j+1} \frac{(j+1)!}{\vec{\gamma}_p!} (h_+^{1\vec{1},pp})^{\vec{\gamma}_p} w_+(p) D_{2\vec{\gamma}_p}^{2j+2} f_{w_+(p)}(0)$$

where we neglect terms with  $\leq 2j + 1$  derivatives.

*Proof.* We argued diagrammatically that the power  $k^{-j}$  is the greatest power of  $k$  in which  $D_{2\vec{\gamma}}^{2j+2} f_{\pm}(0)$  appears. We also showed that a labeled graph with Euler characteristic  $-j$  which produces  $D_{2\vec{\gamma}}^{2j+2} f_{\pm}(0)$  must have a closed vertex of valency  $\geq 2j + 2$ . Now it is clear that such graph must have only one closed vertex and  $j + 1$  loops. This proves the first part of the proposition. The second part follows easily from Lemma 7.6.

Now let us determine  $\sum_{\ell_p} I_{\ell_p}(\Gamma_{1,j+1})$  for the labelled graphs  $(\Gamma_{1,j+1}, \ell_p)$  above. We have

$$(50) \quad \sum_{\ell_p} I_{\ell_p}(\Gamma_{1,j+1}) \equiv (2rL)L^{-(n-1)r} \mathcal{A}(r) i^{j+2} \sum_{\gamma_p^1 + \dots + \gamma_p^{n-1} = j+1} \frac{(j+1)!}{\gamma_p^1! \dots \gamma_p^{n-1}!} (h_+^{1\vec{1},pp})^{\gamma_p^1} \dots (h_+^{(n-1)(n-1),pp})^{\gamma_p^{n-1}} D_{2\vec{\gamma}_p}^{2j+2} \mathcal{L}_+(0).$$

So by Lemma 7.6 and using short-hand notations for multi-indices we get

$$\sum_{\ell_p} I_{\ell_p}(\Gamma_{1,j+1}) \equiv (4rL)L^{-(n-1)r} \mathcal{A}(r) i^{j+2} \sum_{|\vec{\gamma}_p|=j+1} \frac{(j+1)!}{\vec{\gamma}_p!} (h_+^{1\vec{1},pp})^{\vec{\gamma}_p} w_+(p) D_{2\vec{\gamma}_p}^{2j+2} f_{w_+(p)}(0)$$

□

**7.5. Proof of Theorem 2.** Now we are ready to prove Theorem 2. The discussion above shows that modulus derivatives of order  $\leq 2j + 1$

$$B_{\gamma^r,j} = (2rL)^{-1} L^{(n-1)r} \sum_{p=1}^{2r} \frac{\sum_{\ell_p} I_{\ell_p}(\Gamma_{1,j+1})}{S(\Gamma_{1,j+1})}.$$

We notice that  $S(\Gamma_{1,j+1}) = |\text{Aut}(\Gamma_{1,j+1})| = 2^{j+1}(j+1)!$ . We then break up the sums over  $p$  of even/odd parity and use Proposition 7.4 to replace the odd parity Hessian elements by  $h_+^{11}$  and the even ones by  $h_+^{22}$ . Taking into account that  $w_+(p) = 1(-1)$  if  $p$  is even (odd), we conclude that ( by the formula in Proposition 7.7)

$$B_{\gamma^r,j} = \frac{B_{\gamma^r,0}}{(2i)^{j+1}} \sum_{|\gamma|=j+1} \frac{r}{\bar{\gamma}!} \left\{ (\overrightarrow{h_{+,2r}^{11}})^{\bar{\gamma}} D_{2\bar{\gamma}}^{2j+2} f_+(0) - (\overrightarrow{h_{-,2r}^{11}})^{\bar{\gamma}} D_{2\bar{\gamma}}^{2j+2} f_-(0) \right\}.$$

So far we have proved all parts of Theorem 2 except the last part which finds a formula for the wave invariants in the case of symmetries.

7.5.1. *Balian-Bloch invariants at bouncing ball orbits of  $(\mathbb{Z}/2\mathbb{Z})^n$  symmetric domains.* Now if we assume the  $(\mathbb{Z}/2\mathbb{Z})^n$  symmetry assumptions, namely  $f_+ = f = -f_-$  and  $f$  being even in all variables, then using (49) the formula above simplifies to

$$(51) \quad B_{\gamma^r,j} = \frac{B_{\gamma^r,0}}{(2i)^{j+1}} \sum_{|\bar{\gamma}|=j+1} \frac{r}{\bar{\gamma}!} \left( \frac{1}{\sin \frac{\alpha}{2}} \cot \frac{r\bar{\alpha}}{2} \right)^{\bar{\gamma}} D_{2\bar{\gamma}}^{2j+2} f(0) \\ + \{\text{polynomial of Taylor coefficients of order } \leq 2j\}.$$

This finishes the proof of Theorem 2. Q.E.D.

7.6. **Recovering the Taylor Coefficients and the Proof of Theorem 1.** First of all we prove the following lemma

**LEMMA 7.8.** *If  $\{\alpha_1, \dots, \alpha_{n-1}\}$  are linearly independent over  $\mathbb{Q}$  then the functions*

$$\left( \cot \frac{r\bar{\alpha}}{2} \right)^{\bar{\gamma}}$$

*are linearly independent over  $\mathbb{C}$  as functions of  $r \in \mathbb{N}$ .*

*Proof.* Suppose that there exist coefficients  $c_{\bar{\gamma}}$  such that

$$\sum_{\bar{\gamma}} c_{\bar{\gamma}} \left( \cot \frac{r\bar{\alpha}}{2} \right)^{\bar{\gamma}} = 0, \quad \forall r \in \mathbb{N}.$$

Consider the function

$$\psi(z_1, \dots, z_{n-1}) := \sum_{\bar{\gamma}} c_{\bar{\gamma}} (\cot \bar{z})^{\bar{\gamma}}.$$

This function is meromorphic and periodic of period  $2\pi$  in each variable  $z_j$ , so it may be regarded as a meromorphic function on  $(\mathbb{C}/\mathbb{Z})^{n-1}$ . It vanishes when  $z_j = r\alpha_j/2$  modulo  $2\pi$  for all  $r = 1, 2, 3, \dots$ . But such points are dense in the real submanifold  $(\mathbb{R}/\mathbb{Z})^{n-1}$  and hence the function vanishes identically on  $(\mathbb{C}/\mathbb{Z})^{n-1}$ . This is a contradiction since the functions  $\prod_{j=1}^{n-1} w_j^{\gamma_j}$  are independent functions and by the change of variables  $w_j = \cot z_j$  the functions  $\prod_{j=1}^{n-1} (\cot z_j)^{\gamma_j}$  must also be independent.  $\square$

Now assume  $\Omega \subset \mathbb{R}^n$  is a domain in the class  $\mathcal{D}_L$  defined in (3). Take a non-degenerate bounding ball orbit  $\gamma$  of length  $2L$  which satisfies all the properties listed in (3). We would like to use mathematical induction and recover the Taylor coefficients of the function  $f$  where  $f$  and  $-f$  are the local defining functions of  $\partial\Omega$  near the top and bottom of the bouncing ball

orbit respectively. First, it is possible to recover all the  $\alpha_j$ ,  $1 \leq j \leq n-1$ , under a permutation [Fr]. This is because  $|\det(I - P_{\gamma r})|$  is a spectral invariant (the 0-th wave invariant). But we know that  $|\det(I - P_{\gamma r})| = \prod_{j=1}^{n-1} (2 - 2 \cos(r\alpha_j))$ . Hence  $\prod_{j=1}^{n-1} (\sin^2(r\alpha_j/2))$  is a spectral invariant for all  $r \in \mathbb{N}$ . It is easy to see that this condition determines  $\alpha_j$  under a permutation. We fix this permutation and we argue inductively to recover all the Taylor coefficients. Since  $f$  is even in all the variables, the odd order Taylor coefficients are zero. Now assume  $D_{2^{\vec{\gamma}}}^{2|\vec{\gamma}|} f(0)$  are given for all  $|\vec{\gamma}| \leq j$ . Hence the remainder polynomial term of (51) is given. Now by the above lemma, since all the functions  $(\cot \frac{r\vec{\alpha}}{2})^{\vec{\gamma}}$  are linearly independent, we can recover the Taylor coefficients  $D_{2^{\vec{\gamma}}}^{2j+2} f(0)$ . This concludes the proof of Theorem 1.

The analogous arguments will follow in the hyperbolic or mixed hyperbolic-elliptic cases.

## REFERENCES

- [Al] I. Alexandrova, Semi-classical wavefront set and Fourier integral operators. *Canad. J. Math.* 60 (2008), no. 2, 241-263.
- [AG] D. Alonso and P. Gaspard,  $\hbar$  expansion for the periodic orbit quantization of chaotic systems. *Chaos* 3 (1993), no. 4, 601-612.
- [AM] K. G. Andersson and R. B. Melrose, The propagation of singularities along gliding rays. *Invent. Math.* 41 (1977), no. 3, 97-232.
- [A] S. Axelrod, Overview and warmup example for perturbation theory with instantons. *Geometry and physics (Aarhus, 1995)*, 321-338, *Lecture Notes in Pure and Appl. Math.*, 184, Dekker, New York, 1997.
- [BBa] V.M.Babic and V.S. Buldyrev: *Short-Wavelength Diffraction Theory*, Springer Series on Wave Phenomena 4, Springer-Verlag, New York (1991).
- [BB1] R. Balian and C. Bloch, Distribution of eigenfrequencies for the wave equation in a finite domain I: three-dimensional problem with smooth boundary surface, *Ann. Phys.* 60 (1970), 401-447.
- [BB2] R. Balian and C. Bloch, Distribution of eigenfrequencies for the wave equation in a finite domain. III. Eigenfrequency density oscillations. *Ann. Physics* 69 (1972), 76-160.
- [CP] F. Cardoso and G. Popov, Quasimodes with exponentially small errors associated with elliptic periodic rays. *Asymptot. Anal.* 30 (2002), no. 3-4, 217-247.
- [Ch] J. Chazarain, Construction de la parametrix du probleme mixte hyperbolique pour l'equation des ondes. *C. R. Acad. Sci. Paris Sr. A-B* 276 (1973), A1213-A1215.
- [Chr] T. Christiansen, Sojourn times, manifolds with infinite cylindrical ends, and an inverse problem for planar waveguides. *J. Anal. Math.* 107 (2009), 79-106.
- [CV] Y. Colin de Verdière, Sur les longuers des trajectoires periodiques d'un billard, In: P.Dazord and N. Desolneux-Moulis (eds.) *Geometrie Symplectique et de Contact: Autour du Theoreme de Poincare-Birkhoff. Travaux en Cours, Sem. Sud-Rhodanien de Geometrie III* Paris: Herman (1984), 122-139.
- [CVG] Y. Colin De Verdière and V. Guillemin, A semi-classical inverse problem I: Taylor expansions (arXiv:0802.1605).
- [E] P. Etingof, *Lecture Notes on Quantum Field Theory* (MIT OpenCourseWare).
- [EZ] L.C. Evans and M. Zworski, *Lectures on semiclassical analysis*, Lecture notes, available at <http://math.berkeley.edu/~zworski/semiclassical.pdf>.
- [Fr] D. Fried, Cyclic resultants of reciprocal polynomials. *Holomorphic dynamics (Mexico, 1986)*, 124-128, *Lecture Notes in Math.*, 1345, Springer, Berlin, 1988.
- [Gh] M. Ghomi, Shortest periodic billiard trajectories in convex bodies. *Geom. Funct. Anal.* 14 (2004), no. 2, 295-302.
- [GS] C. S. Gordon and Z. I. Szabo, Isospectral deformations of negatively curved Riemannian manifolds with boundary which are not locally isometric. *Duke Math. J.* 113 (2002), no. 2, 355-383.
- [GW] C. S. Gordon and D. L. Webb, Isospectral convex domains in Euclidean space. *Math. Res. Lett.* 1 (1994), no. 5, 539-545

- [GWW] C. Gordon, D. Webb, and S. Wolpert, Isospectral plane domains and surfaces via Riemannian orbifolds. *Invent. Math.* 110 (1992), no. 1, 1–22
- [G] V. Guillemin, Wave trace invariants, *Duke Math. J.* 83 (1996), 287–352.
- [G2] V. Guillemin, Wave trace invariants and a theorem of Zelditch, *Int. Math. Res. Not.* 12 (1993), 303–308.
- [GU] V. Guillemin and A. Uribe, Some inverse spectral results for semi-classical Schrödinger operators. *Math. Res. Lett.* 14 (2007), no. 4, 623–632.
- [GM] V. Guillemin and R.B. Melrose, The Poisson summation formula for manifolds with boundary, *Adv. in Math.* 32 (1979), 204 - 232.
- [GuS] V. Guillemin and S. Sternberg, *Geometric asymptotics*. Mathematical Surveys, No. 14. American Mathematical Society, Providence, R.I., 1977.
- [HZ] A. Hassell and S. Zelditch, Quantum ergodicity of boundary values of eigenfunctions, *Comm. Math. Phys.* 248 (2004), no. 1, 119–168 (preprint arXiv: math.SP/0211140).
- [H] H. Hezari, Inverse spectral problems for Schrödinger operators, *Comm. Math. Phys.* 288 (2009), no. 3, 1061–1088 (arXiv:0801.3283).
- [HeZ] H. Hezari and S. Zelditch, Analytic spectral rigidity of plane domains with one symmetry (in preparation).
- [Hö] L. Hörmander, *The Analysis of Linear Partial Differential Operators, Volumes I–IV*, Springer-Verlag Berlin Heidelberg, 1983.
- [ISZ] A. Iantchenko, J. Sjöstrand and M. Zworski, Birkhoff normal forms in semi-classical inverse problems, *Math. Res. Lett.* 9 (2002), no. 2-3, 337–362 (arXiv.org/abs/math.SP/0201191).
- [KAC] M. Kac, Can one hear the shape of a drum? *Amer. Math. Monthly* 73 1966 no. 4, part II, 1–23.
- [KT] V.V. Kozlov and D. V. Treshchev, *Billiards: A Genetic Introduction to the Dynamics of Systems with Impacts*, Translations of Math. Monographs 89, AMS publications, Providence, R.I. (1991).
- [L] V.F. Lazutkin, Construction of an asymptotic series of eigenfunctions of the “bouncing ball” type, *Proc. Steklov Inst. Math.* 95 (1968), 125- 140.
- [LT] V.F. Lazutkin and D.Ya. Terman, Number of quasimodes of ‘bouncing ball’ type, *J. Soviet Math.* (1984), 373-379.
- [PS] V.M. Petkov and L.N. Stoyanov, *Geometry of Reflecting Rays and Inverse Spectral Problems*, John Wiley and Sons, N.Y. (1992).
- [SZ] J. Sjöstrand and M. Zworski, Quantum monodromy and semi-classical trace formulae, *J. Math. Pures Appl.* (9) 81 (2002), no. 1, 1–33.
- [T] M.E. Taylor, *Partial Differential Equations, I- II*, Appl. Math. Sci. 115 -116, Springer-Verlag (1996).
- [U] H. Urakawa, Bounded domains which are isospectral but not congruent. *Ann. Sci. École Norm. Sup.* (4) 15 (1982), no. 3, 441-456.
- [Z1] S. Zelditch, Wave invariants at elliptic closed geodesics. *Geom. Funct. Anal.* 7 (1997), no. 1, 145-213.
- [Z2] S. Zelditch, Spectral determination of analytic bi-axisymmetric plane domains, *Geom. Funct. Anal.* 10 (2000), no. 3, 628-677.
- [Z3] S. Zelditch, Inverse spectral problem for analytic domains II: domains with one symmetry, *Ann. of Math.* (2) 170 (2009), no. 1, 205–269 ( arXiv:math/0111078).
- [Z4] S. Zelditch, Inverse spectral problem for analytic domains. I. Balian-Bloch trace formula. *Comm. Math. Phys.* 248 (2004), no. 2, 357-407.
- [Z5] S. Zelditch, Inverse resonance problem for  $\mathbb{Z}_2$ -symmetric analytic obstacles in the plane. *Geometric methods in inverse problems and PDE control*, 289-321, IMA Vol. Math. Appl., 137, Springer, New York, 2004.

DEPARTMENT OF MATHEMATICS, JOHNS HOPKINS UNIVERSITY, BALTIMORE, MD 21218, USA

*E-mail address:* zelditch@math.jhu.edu, hhezari@math.jhu.edu