

# COUNTING GENERIC MEASURES FOR A SUBSHIFT OF LINEAR GROWTH

VAN CYR AND BRYNA KRA

ABSTRACT. In 1984 Boshernitzan proved an upper bound on the number of ergodic measures for a minimal subshift of linear block growth and asked if it could be lowered without further assumptions on the shift. We answer this question, showing that Boshernitzan's bound is sharp. We further prove that the same bound holds for the, a priori, larger set of nonatomic generic measures, and that this bound remains valid even if one drops the assumption of minimality. Applying these results to interval exchange transformations, we give an upper bound on the number of nonatomic generic measures of a minimal IET, answering a question recently posed by Chaika and Masur.

## 1. INTRODUCTION

Let  $(X, \sigma)$  be a subshift, meaning that  $X \subset \mathcal{A}^{\mathbb{Z}}$ , where  $\mathcal{A}$  is a finite alphabet, and  $X$  is a closed set that is invariant under the left shift  $\sigma: \mathcal{A}^{\mathbb{Z}} \rightarrow \mathcal{A}^{\mathbb{Z}}$ . A classic problem is to find conditions that imply  $(X, \sigma)$  is uniquely ergodic or, more generally, has a finite number of ergodic measures. In the 1980's, Boshernitzan [1] showed that the complexity of the subshift can be used to obtain such a result. More precisely, if  $P_X(n)$  is the number of words of length  $n$  which occur in any  $x \in X$ , he showed that if  $(X, \sigma)$  is minimal and  $\limsup_{n \rightarrow \infty} P_X(n)/n < 3$ , then it is uniquely ergodic (see also related results in [3]). More generally, Boshernitzan showed that if

$$(1) \quad \liminf_{n \rightarrow \infty} \frac{P_X(n)}{n} < k,$$

then there are at most  $k - 1$  ergodic measures. Some motivation for studying this problem is generalizing the well-known bound on the number of ergodic measures for an interval exchange transformation (IET), that had been previously proven, independently, by Katok and Veech. Boshernitzan's Theorem applies to a much broader class of dynamical systems than the interval exchange transformations, but the bound he obtains is weaker than that of Katok and Veech in the case of an IET. Boshernitzan asked in [1], and then again in [2], whether his bound could be lowered in this more general setting. One of our main results answers Boshernitzan's question: for the class of minimal subshifts whose complexity function satisfies (1), Boshernitzan's bound is a sharp bound for the number of nonatomic ergodic measures. Our technique also shows that the bound is more general than originally stated: the same bound remains valid (and sharp) even without the assumption of minimality and even if one seeks to bound the (a priori, larger) set of nonatomic generic measures.

---

2010 *Mathematics Subject Classification.* 37B10 (primary), 37A25, 68R15.

*Key words and phrases.* subshift, automorphism, block complexity.

The second author was partially supported by NSF grant 1500670.

The particular case of minimal interval exchange transformations has been well studied (for example Katok [10], Keane [12], and Veech [14]). A minimal  $k$ -interval exchange transformation ( $k$ -IET) has a natural symbolic cover, its natural coding, and this subshift satisfies the hypothesis of Boshernitzan's Theorem. As an application, this shows that a minimal  $k$ -IET (see Section 4 for the definition) has at most  $k - 1$  ergodic measures. The optimal bound of  $\lfloor k/2 \rfloor$  was proven, independently, by Katok [10] and Veech [14]. In a recent paper, Chaika and Masur [4] studied the broader class of generic measures for an IET and asked whether there are bounds on the number of such measures. An interesting facet of this problem is that although several quite different proofs of the bound given by Katok and Veech for the number of ergodic measures exist in the literature, they all use ergodicity in an essential way.

If  $X$  is a compact metric space,  $\mathcal{B}$  the Borel  $\sigma$ -algebra,  $\mu$  a Borel probability measure on  $\mathcal{B}$ , and  $T: X \rightarrow X$  is a measurable map preserving the measure  $\mu$ , a point  $x \in X$  is a *generic point for the measure  $\mu$*  if for every continuous function  $f: X \rightarrow \mathbb{R}$ ,

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=0}^{N-1} f(T^n x) = \int f d\mu.$$

The measure  $\mu$  is *generic* if it has a generic point. Thus, by the Pointwise Ergodic Theorem, if the measure  $\mu$  is ergodic almost every point is generic. However, a generic measure need not be ergodic. Chaika and Masur [4] constructed a 6-interval exchange transformation that has a generic, but not ergodic, measure. They asked if there is a bound on the number of generic measures for a  $k$ -IET. We show:

**Theorem 1.1.** *If  $(X, \sigma)$  is a subshift and there exists  $k \in \mathbb{N}$  such that*

$$\liminf_{n \rightarrow \infty} \frac{P_X(n)}{n} < k,$$

*then  $(X, \sigma)$  has at most  $k - 1$  distinct, nonatomic, generic measures.*

In particular, this applies to interval exchange transformations by passing to the natural cover. Theorem 1.1 generalizes Boshernitzan's Theorem [1] in two ways: there is no assumption of minimality and our bound holds for the more general class of generic measures. We also give an analogous bound for  $\limsup$  (note the technical assumption is vacuous for minimal subshifts that are not uniquely ergodic).

**Theorem 1.2.** *Suppose  $(X, \sigma)$  is a subshift and there exists  $k \in \mathbb{N}$  such that*

$$\limsup_{n \rightarrow \infty} \frac{P_X(n)}{n} < k.$$

*If  $(X, \sigma)$  has a generic measure  $\mu$  and a generic point  $x_\mu$  for which the orbit closure*

$$\overline{\{\sigma^k x_\mu : k \in \mathbb{Z}\}}$$

*is not uniquely ergodic, then  $(X, \sigma)$  has at most  $k - 2$  distinct, nonatomic, generic measures.*

Recently Damron and Fickenscher [5] proved a related result, showing that any minimal shift  $(X, \sigma)$  whose complexity function satisfies  $P_X(n) = kn + c$  for some constant  $c$ ,  $k \geq 4$ , and all  $n$  sufficiently large has at most  $k - 2$  ergodic measures.

Moreover, we show that these theorems are sharp, even if  $X$  is assumed to be minimal and the measures are required to be ergodic.

**Theorem 1.3.** *Suppose  $d > 1$  is an integer. There exists a minimal subshift  $(X, \sigma)$  which has exactly  $d$  ergodic measures, zero nonergodic generic measures, and which satisfies*

$$\begin{aligned} \liminf_{n \rightarrow \infty} \frac{P_X(n)}{n} &= d; \\ \limsup_{n \rightarrow \infty} \frac{P_X(n)}{n} &= d + 1. \end{aligned}$$

We include several other examples in Section 5, showing other senses in which Theorems 1.1 and 1.2 can be said to be sharp.

As an application of Theorem 1.1, we answer Chaika and Masur’s question:

**Theorem 1.4.** *For  $k > 2$ , a minimal  $k$ -interval exchange transformation has at most  $k - 2$  generic measures.*

For  $k = 2$ , a minimal 2-interval exchange is an ergodic rotation, which is uniquely ergodic. For  $k = 3$  and 4, Theorem 1.4 is sharp upper bound, but we do not know if it is sharp for  $k \geq 5$ . In particular, we do not know if we can improve the symbolic result of Theorem 1.1 for systems that arise as the natural coding of an interval exchange transformation. Similarly, we would like to know if the conclusion of Theorem 1.4 can be strengthened to replace  $k - 2$  by  $\lfloor k/2 \rfloor$ . We also do not know if there can be a second generic measure in the example of Chaika and Masur, nor if a 6-interval exchange with three ergodic measures can also have a generic (and obviously nonergodic) measure.

## 2. BACKGROUND AND NOTATION

If  $\mathcal{A}$  is a finite alphabet, a *word*  $w$  in the alphabet is a concatenation of letters in  $\mathcal{A}$  and the length  $|w|$  of the word is the number (finite or infinite) of letters. A word  $w = w_1 \dots w_\ell$  occurs in a word  $u = u_1 \dots u_k$  if there is some  $m \in \{1, \dots, k - \ell\}$  such that  $w_1 = u_m, \dots, w_\ell = u_{m+\ell}$ , and we refer to  $w$  as a *subword* of  $u$ . The analogous definitions hold for a finite word  $w$  occurring as a subword of an infinite word  $u$ .

A *language*  $\mathcal{L}$  is a set of (finite) words such that if  $w \in \mathcal{L}$ , then any subword is also contained in  $\mathcal{L}$ . The language *determined by a word* (finite or infinite) is the collection of all finite subwords of the word. We let  $\mathcal{L}_n$  denote all the words in the language  $\mathcal{L}$  of length  $n$ . If  $w \in \mathcal{L}$ , we write  $[w]$  for the *cylinder set determined by  $w$* , meaning that

$$[w] = \{u \in \mathcal{L} : \text{the first } |w| \text{ symbols of } u \text{ agree with } w\}.$$

We assume that the alphabet  $\mathcal{A}$  is endowed with the discrete topology and if  $x \in \mathcal{A}^{\mathbb{Z}}$ , we use  $x(n)$  to denote the value of  $x$  at  $n \in \mathbb{Z}$ . The space  $\mathcal{A}^{\mathbb{Z}}$  is a compact metric space when endowed with the product topology (and a compatible metric).

A *subshift*  $(X, \sigma)$  is a closed subset  $X \subset \mathcal{A}^{\mathbb{Z}}$  that is invariant under the left shift  $\sigma: \mathcal{A}^{\mathbb{Z}} \rightarrow \mathcal{A}^{\mathbb{Z}}$  defined by  $(\sigma x)(n) = x(n + 1)$ . If  $\mathcal{L}$  is the language of the system  $X$ , meaning the set of all finite subwords that arise for any  $x \in X$ , we write  $\mathcal{L} = \mathcal{L}(X)$  and we write  $\mathcal{L}_n = \mathcal{L}_n(X)$  for the words of length  $n$ . We define the *complexity function*  $P_X: X \rightarrow \mathbb{N}$  by

$$P_X(n) = |\mathcal{L}_n(X)|.$$

For a word  $w \in \mathcal{L}(X)$ , we write  $\mathbf{1}_{[w]}$  for the indicator function of the word  $w$ . We say that  $x = (x(n))_{n \in \mathbb{Z}} \in X$  is *periodic* if there exists  $m \neq 0$  such that

$x(m+n) = x(n)$  for all  $n \in \mathbb{Z}$  and otherwise it is *aperiodic*. The point  $x$  is *eventually periodic* if there exists  $m \neq 0$  and  $N \in \mathbb{N}$  such that  $x(m+n) = x(n)$  for all  $n \geq N$ .

For a system  $(X, \sigma)$ , the *orbit* of  $x \in X$  is defined to be  $\{\sigma^n x : n \in \mathbb{Z}\}$  and the system is *minimal* if the orbit closure  $\overline{\{\sigma^n x : n \in \mathbb{Z}\}} = X$  for any  $x \in X$ .

For  $N, m \in \mathbb{N}$ , define  $w(N, m) \in \mathcal{L}_N(X)$  by

$$(2) \quad w(N, m) := (x(m), x(m+1), \dots, x(m+N-1))$$

to be the word of length  $N$  that occurs in  $x$  starting at location  $m$ . We make use of the following theorem (though stated differently) of Epifanio, Koskas, and Mignosi [7]:

**Theorem 2.1** ([7, Theorem 2.2]). *Assume  $x \in \mathcal{A}^{\mathbb{N}}$  is not eventually periodic and fix  $M, N_0 \in \mathbb{N}$ . Suppose that for some  $N \geq N_0$ , there exist  $M \leq m_1 < m_2 \leq N$  such that  $w_x(N, m_1) = w_x(N, m_2)$ . Then there exists  $K \geq m_1$  such that*

- (i) (*Distinct Words Condition*): *for all  $K \leq k_1 < k_2 \leq K + N - N_0$  we have  $w_x(N, k_1) \neq w_x(N, k_2)$ ;*
- (ii) (*Prefix First Occurrence Condition*): *for all  $K \leq k < K + N - N_0$  there exists  $M \leq l_k \leq N$  such that  $w_x(N_0, k) = w_x(N_0, l_k)$ .*

For completeness, we include the proof, but it is merely a translation of the proof in [7] using our hypotheses and emphasizing the stronger conclusion.

*Proof.* Suppose  $w_x(N, m_1) = w_x(N, m_2)$ . Then the word  $w_x(N + m_2 - m_1, m_1)$  is periodic of period  $m_2 - m_1$ . Since  $x$  is not eventually periodic, there exists  $N' \geq N + m_2 - m_1$  such that  $w_x(N', m_1)$  is periodic of period  $m_2 - m_1$ , while  $w_x(N' + 1, m_1)$  is not. Let  $1 \leq p \leq m_2 - m_1$  be the minimal period of  $w_x(N', m_1)$  and define  $K := m_1 + N' - N - p \geq m_1$ . By minimality of  $p$  and the fact that  $N \geq p$ , if  $K \leq i < j \leq K + p - 1$  then  $w_x(N, i) \neq w_x(N, j)$  (and all such words are periodic of period  $p$ ).

For contradiction, suppose there exist  $K \leq i < j \leq K + N - N_0$  such that  $w_x(N, i) = w_x(N, j)$ . Since  $i, j$  cannot both be smaller than  $K + p$ , it follows that  $j \geq K + p$ . The word  $w_x(N + (j - i), i)$  is periodic of period  $j - i$  and its prefix of length  $p + j - i$  is periodic of period  $p$ . By the Fine-Wilf Theorem [9], it follows that this prefix is periodic of period  $\gcd(j - i, p)$ . Since this prefix has length at least  $p$ , it follows that  $w_x(N + (j - i), i)$  is periodic of period  $\gcd(j - i, p)$  and, in particular, is periodic of period  $p$ . Moreover,  $K \leq i \leq K + N - N_0$  and so  $w_x(N + (j - i), i)$  begins at least  $p$  spaces before  $N' + 1$  and ends at location

$$i + N + (j - i) = N + j \geq N + (K + p) = m_1 + N' + 1.$$

Thus the periodicity of  $w_x(N + (j - i), i)$  contradicts the fact that  $w_x(N', m_1)$  is not periodic of period  $p$ , which implies that  $w_x(N, i) \neq w_x(N, j)$  for any  $M \leq i < j \leq K + n - N_0$ .

Since  $w_x(N', m_1)$  is periodic of period  $p \leq n$  and the length  $N_0$  prefix of  $w_i(N, i)$  is a subword of  $w_x(N', m_1)$ , the second statement follows.  $\square$

### 3. MAIN RESULTS

Theorems 1.1 and 1.2 follow from the following estimate:

**Theorem 3.1.** *Let  $(X, \sigma)$  be a subshift which has at least  $d \geq 1$  distinct, nonatomic, generic measures. Then*

$$\liminf_{n \rightarrow \infty} \frac{P_X(n)}{n} \geq d.$$

*If, in addition,  $(X, \sigma)$  has a generic measure  $\mu$  and a generic point  $x_\mu$  whose orbit closure*

$$\overline{\{\sigma^k x_\mu : k \in \mathbb{N}\}}$$

*is not uniquely ergodic, then*

$$\limsup_{n \rightarrow \infty} \frac{P_X(n)}{n} \geq d + 1.$$

*Proof.* We show that for arbitrarily small  $\delta > 0$ , we have

$$(3) \quad \liminf_{n \rightarrow \infty} \frac{P_X(n)}{n} > d - 2d\delta$$

and, under the additional hypothesis of a generic measure and associated generic point whose orbit closure is not uniquely ergodic,

$$(4) \quad \limsup_{n \rightarrow \infty} \frac{P_X(n)}{n} > d + 1 - 2d\delta.$$

The theorem follows immediately from these estimates.

Fix  $\delta > 0$ , and for convenience assume that  $1/\delta \in \mathbb{N}$ . Suppose  $\mu_1, \dots, \mu_d$  are distinct, nonatomic, generic measures for  $(X, \sigma)$  and choose  $x_1, \dots, x_d \in X$  such that for each  $1 \leq i \leq d$ ,  $x_i$  is generic for  $\mu_i$ . Observe that if  $x$  is eventually periodic and is generic for some measure  $\mu$ , then  $\mu$  must be the atomic measure supported on the (eventual) period of  $x$ . Therefore, since  $\mu_i$  is nonatomic,  $x_i$  is not eventually periodic for all  $i$ . By definition of  $x_i$ , for all  $w \in \mathcal{L}(X)$  we have

$$(5) \quad \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{k=0}^{N-1} 1_{[w]}(T^k x_i) = \mu_i([w]).$$

For  $1 \leq j_1 < j_2 \leq d$ , choose words  $w_{(j_1, j_2)} \in \mathcal{L}(X)$  such that  $\mu_{j_1}([w_{(j_1, j_2)}]) \neq \mu_{j_2}([w_{(j_1, j_2)}])$ . Set

$$(6) \quad \varepsilon := \min\{|\mu_{j_1}([w_{(j_1, j_2)}]) - \mu_{j_2}([w_{(j_1, j_2)}])| : 1 \leq j_1 < j_2 \leq d\}$$

and set

$$(7) \quad B := \frac{\delta}{16 - 4\delta}.$$

By (5), for each  $1 \leq i \leq d$  there exists  $N_i \in \mathbb{N}$  such that for all  $N \geq N_i$  and all  $1 \leq j_1 < j_2 \leq d$ , we have

$$(8) \quad \left| \frac{1}{N} \sum_{k=0}^{N-1} 1_{[w_{(j_1, j_2)}]}(T^k x_i) - \mu_i([w_{(j_1, j_2)}]) \right| < B \cdot \varepsilon.$$

Set

$$(9) \quad M := \max_{1 \leq i \leq d} N_i.$$

Analogous to notation (2), for  $1 \leq i \leq d$  and for  $N, m \in \mathbb{N}$ , define  $u_i(N, m) \in \mathcal{L}_N(X)$  by

$$u_i(N, m) := (x_i(m), x_i(m+1), x_i(m+2), \dots, x_i(m+N-1))$$

to be the word of length  $N$  that occurs in  $x_i$  starting at location  $m$ .

If  $u, w \in \mathcal{L}(X)$  and  $|u| \geq |w|$ , define the frequency with which  $w$  occurs as a subword in  $u$  to be

$$(10) \quad F(u, w) := \frac{1}{|u| - |w| + 1} \sum_{k=0}^{|u|-|w|} \mathbf{1}_{[w]}(T^k x),$$

where  $x \in [u]$ . Note that this frequency does not depend on the choice of  $x \in [u]$ , as it only depends on the first  $|u|$  coordinates of  $x$ . Suppose

$$N \geq \frac{1}{\delta} \cdot (M + \max\{|w_{(j_1, j_2)}| : 1 \leq j_1 < j_2 \leq d\})$$

is fixed and define  $L_N := \lfloor (2-\delta)N \rfloor$  and  $\ell_N := \lfloor \delta N \rfloor$ . By definition,  $\ell_N - |w_{(j_1, j_2)}| \geq M$  for all  $w_{(j_1, j_2)}$ . If  $1 \leq i \leq d$ ,  $1 \leq j_1 < j_2 \leq d$ , and  $M \leq L \leq L_N$ , then the frequency with which the word  $w_{(j_1, j_2)}$  occurs in the subword of  $x_i$  with length  $\ell_N$  and starting from location  $L$  is given by (recall that  $u_i(\ell_N, L)$  is the word of length  $\ell_N$  that starts at location  $L$  in  $x_i$ )

$$\begin{aligned} & F(u_i(\ell_N, L), w_{(j_1, j_2)}) \\ &= \frac{1}{\ell_N - |w_{(j_1, j_2)}| + 1} \sum_{k=0}^{\ell_N - |w_{(j_1, j_2)}|} \mathbf{1}_{[w_{(j_1, j_2)}]}(T^k(T^L x_i)) \\ &= \frac{1}{\ell_N - |w_{(j_1, j_2)}| + 1} \sum_{k=L}^{L + \ell_N - |w_{(j_1, j_2)}|} \mathbf{1}_{[w_{(j_1, j_2)}]}(T^k x_i) \\ &= \frac{1}{\ell_N - |w_{(j_1, j_2)}| + 1} \left( \sum_{k=0}^{L + \ell_N - |w_{(j_1, j_2)}|} \mathbf{1}_{[w_{(j_1, j_2)}]}(T^k x_i) - \sum_{k=0}^{L-1} \mathbf{1}_{[w_{(j_1, j_2)}]}(T^k x_i) \right) \\ &= \frac{L + \ell_N - |w_{(j_1, j_2)}| + 1}{\ell_N - |w_{(j_1, j_2)}| + 1} \cdot \frac{1}{L + \ell_N - |w_{(j_1, j_2)}| + 1} \sum_{k=0}^{L + \ell_N - |w_{(j_1, j_2)}|} \mathbf{1}_{[w_{(j_1, j_2)}]}(T^k x_i) \\ &\quad - \frac{L}{\ell_N - |w_{(j_1, j_2)}| + 1} \cdot \frac{1}{L} \sum_{k=0}^{L-1} \mathbf{1}_{[w_{(j_1, j_2)}]}(T^k x_i). \end{aligned}$$

But by (8),

$$\left| \frac{1}{L + \ell_N - |w_{(j_1, j_2)}| + 1} \sum_{k=0}^{L + \ell_N - |w_{(j_1, j_2)}|} \mathbf{1}_{[w_{(j_1, j_2)}]}(T^k x_i) - \mu_i([w_{(j_1, j_2)}]) \right| < B \cdot \varepsilon$$

and since  $L \geq M$ , we have

$$\left| \frac{1}{L} \sum_{k=0}^{L-1} \mathbf{1}_{[w_{(j_1, j_2)}]}(T^k x_i) - \mu_i([w_{(j_1, j_2)}]) \right| < B \cdot \varepsilon.$$

Therefore

$$\begin{aligned}
& |F(u_i(\ell_N, L), w_{(j_1, j_2)}) - \mu_i([w_{(j_1, j_2)}])| \\
& \leq \frac{L + \ell_N - |w_{(j_1, j_2)}| + 1}{\ell_N - |w_{(j_1, j_2)}| + 1} \cdot B \cdot \varepsilon + \frac{L}{\ell_N - |w_{(j_1, j_2)}| + 1} \cdot B \cdot \varepsilon \\
& = \frac{2L + \ell_N - |w_{(j_1, j_2)}| + 1}{\ell_N - |w_{(j_1, j_2)}| + 1} \cdot B \cdot \varepsilon \\
& \leq \frac{2\lfloor(2 - \delta)N\rfloor + \lfloor\delta N\rfloor - |w_{(j_1, j_2)}| + 1}{\lfloor\delta N\rfloor - |w_{(j_1, j_2)}| + 1} \cdot B \cdot \varepsilon.
\end{aligned}$$

By Definition (7) that  $B = \frac{\delta}{16-4\delta}$ , for all sufficiently large  $N$  this inequality implies

$$(11) \quad |F(u_i(\ell_N, L), w_{(j_1, j_2)}) - \mu_i([w_{(j_1, j_2)}])| < \frac{\varepsilon}{2}.$$

By (6), for all sufficiently large  $N$  and all  $L_1, L_2 \in \{M, M+1, \dots, \lfloor(2-\delta)N\rfloor\}$  we have that if  $1 \leq i_1 < i_2 \leq d$ , then the frequency with which  $w_{(i_1, i_2)}$  occurs in  $u_{i_1}(\ell_N, L_1)$  is different than its frequency in  $u_{i_2}(\ell_N, L_2)$ . Therefore  $u_{i_1}(\ell_N, L_1) \neq u_{i_2}(\ell_N, L_2)$ . For  $1 \leq i \leq d$  define

$$\mathcal{W}_i(N) := \{u_i(\ell_N, L) : M \leq L \leq \lfloor(2-\delta)N\rfloor\} \subseteq \mathcal{L}_{\ell_N}(X).$$

We have shown that for all sufficiently large  $N$ , if  $1 \leq i_1 < i_2 \leq d$ , then

$$(12) \quad \mathcal{W}_{i_1}(N) \cap \mathcal{W}_{i_2}(N) = \emptyset.$$

Fix  $i$  with  $1 \leq i \leq d$  and fix  $N$  sufficiently large such that (12) holds. If the words

$$(13) \quad u_i(N, M), u_i(N, M+1), u_i(N, M+2), \dots, u_i(N, \lfloor(1-\delta)N\rfloor)$$

are all distinct, then the set

$$(14) \quad \mathcal{S}_i := \{w \in \mathcal{L}_N(X) : \text{all subwords of } w \text{ of length } \ell_N \text{ are elements of } \mathcal{W}_i(N)\}$$

contains at least  $\lfloor(1-\delta)N\rfloor - M$  elements. If, on the other hand, the words in (13) are not all distinct, then there exist  $M \leq L_1 < L_2 \leq \lfloor(1-\delta)N\rfloor$  such that  $u_i(N, L_1) = u_i(N, L_2)$ . In this case, by Theorem 2.1 there exists  $K \in \mathbb{N}$  such that

- (i) (*Distinct Words Condition*): for all  $K \leq k_1 < k_2 \leq K + N - \ell_N$  we have  $u_i(N, k_1) \neq u_i(N, k_2)$ ;
- (ii) (*Prefix First Occurrence Condition*): for all  $K \leq k \leq K + N - \ell_N$  there exists  $\ell_N \leq l_k \leq N$  such that  $u_i(\ell_N, k) = u_i(\ell_N, l_k)$ .

Thus in this case, the set

$$(15) \quad \mathcal{T}_i := \{w \in \mathcal{L}_N(X) : \text{the leftmost subword } w \text{ of length } \ell_N \text{ lies in } \mathcal{W}_i(N)\}$$

contains at least  $N - \ell_N$  elements.

By (12),  $\mathcal{S}_{i_1} \cap \mathcal{S}_{i_2} = \emptyset$  whenever  $i_1 \neq i_2$  (and both sets are defined). A similar statement holds when comparing any  $\mathcal{S}_{i_1}$  to  $\mathcal{T}_{i_2}$  for any  $i_2$ , or when comparing  $\mathcal{T}_{i_1}$  to  $\mathcal{T}_{i_2}$ . Thus for each  $1 \leq i \leq d$ , we have associated either the set  $\mathcal{S}_i$  or the set  $\mathcal{T}_i$  and

$$P_X(N) \geq d \cdot \min\{N - \ell_N, \lfloor(1-\delta)N\rfloor - M\} = d \cdot \min\{N - \lfloor\delta N\rfloor, \lfloor(1-\delta)N\rfloor - M\}.$$

Therefore,

$$\frac{P_X(N)}{N} \geq \frac{d \cdot \min\{N - \lfloor\delta N\rfloor, \lfloor(1-\delta)N\rfloor - M\}}{N},$$

which is larger than  $d - 2d\delta$  for all sufficiently large  $N$ , thus establishing (3).

To prove (4), suppose that there exists  $1 \leq i \leq d$  such that the orbit closure of  $x_i$  is not uniquely ergodic. Then for any fixed (and sufficiently large)  $N \in \mathbb{N}$ , there exist infinitely many  $L \in \mathbb{N}$  such that  $u_i(\ell_N, L) \notin \mathcal{W}_i(N)$ . Fix  $N \in \mathbb{N}$ .

If the words

$$u_i(N, M), u_i(N, M+1), u_i(N, M+2), \dots, u_i(N, \lfloor (1-\delta)N \rfloor)$$

are all distinct, then we define  $\mathcal{S}_i$  as in (14). In this case, choose the smallest  $L \geq M$  for which  $u_i(\ell_N, L) \notin \mathcal{W}_i(N)$ ; clearly  $L > L_N$ . Then each of the words

$$u_i(N, L - N + \ell_N), u_i(N, L - N + \ell_N + 1), \dots, u_i(N, L - \ell_N)$$

has the property that its leftmost subword of length  $\ell_N$  is an element of  $\mathcal{W}_i(N)$ , these words are pairwise distinct (in  $u_i(N, L - N + \ell_N + j)$  the leftmost occurrence of a subword of length  $\ell_N$  that is not in  $\mathcal{W}_i(N)$  begins at location  $L - \ell_N - j$ ). These  $N - \ell_N$  words of length  $N$  do not lie in  $\mathcal{S}_i$ , and are not contained in any  $\mathcal{S}_j$  or  $\mathcal{T}_j$  for any  $j \neq i$  (as defined in (15)), since their leftmost subword of length  $\ell_N$  is in  $\mathcal{W}_i$ . Therefore by the bound on the size of  $\mathcal{S}_i$  following (14),

$$\begin{aligned} P_X(N) &\geq d \cdot \min\{N - \ell_N, \lfloor (1-\delta)N \rfloor - M\} \\ &= d \cdot \min\{N - \lfloor \delta N \rfloor, \lfloor (1-\delta)N \rfloor - M\} + (N - \ell_N) \end{aligned}$$

and so in this case,

$$\frac{P_X(N)}{N} \geq \frac{d \cdot \min\{N - \lfloor \delta N \rfloor, \lfloor (1-\delta)N \rfloor - M\}}{N} + \frac{N - \lfloor \delta N \rfloor}{N}.$$

If  $N$  is sufficiently large, this is larger than  $d + 1 - 2d\delta$ .

Thus we are left with showing that there are infinitely many  $N \in \mathbb{N}$  for which the words

$$(16) \quad u_i(N, M), u_i(N, M+1), u_i(N, M+2), \dots, u_i(N, \lfloor (1-\delta)N \rfloor)$$

are all distinct. Fix some  $N \in \mathbb{N}$  and assume that these words are not all distinct. As before, let  $L_1, L_2 \in \{M, M+1, \dots, \lfloor (1-\delta)N \rfloor\}$  with  $L_1 < L_2$  be such that  $u_i(N, L_1) = u_i(N, L_2)$ . Let  $p$  be the minimal period of the word  $u_i(N + L_2 - L_1, L_1)$  and let  $K$  be the largest integer for which  $u_i(K, L_1)$  is periodic with period  $p$  (note that  $K$  is finite since  $x_i$  is not eventually periodic). Then the words

$$(17) \quad u_i(K, M), u_i(K, M+1), \dots, u_i(K, \lfloor (1-\delta)K \rfloor)$$

are all distinct: if  $j > L_1 - M$  then the word  $u_i(K, M + j)$  begins with a word that is periodic of period  $p$  and has length exactly  $K - L_1 - j$  (so no two words of this form can coincide), and if  $j \leq L_1 - M$  then  $u_i(K, M + j)$  either begins with a word of length  $K - L_1 + j$  that is periodic of period  $p$ , or has a prefix of length at most  $L_1$  followed by a word of length at least  $K - L_1 > N$  that is periodic of period  $p$  (which occurs in a different location for each such  $j$ ). Therefore, for each  $N \in \mathbb{N}$  there exists  $K \geq N$  such that the words in (17) are all distinct, and in particular there are infinitely many  $N$  such that the words in (16) are distinct. This establishes (4).  $\square$

As immediate corollaries of Theorem 3.1, we have the theorems stated in the introduction:

**Corollary** (Theorem 1.1). *If  $(X, \sigma)$  is a subshift and there exists  $k \in \mathbb{N}$  such that*

$$\liminf_{n \rightarrow \infty} \frac{P_X(n)}{n} < k,$$



then  $(X, \sigma)$  has at most  $k - 1$  distinct, nonatomic, generic measures.

**Corollary** (Theorem 1.2). *If  $(X, \sigma)$  is a subshift and there exists  $k \in \mathbb{N}$  such that*

$$\limsup_{n \rightarrow \infty} \frac{P_X(n)}{n} < k,$$

*and if  $(X, \sigma)$  has a generic measure  $\mu$  and a generic point  $x_\mu$  whose orbit closure is not uniquely ergodic, then  $(X, \sigma)$  has at most  $k - 2$  distinct, nonatomic, generic measures.*

In Section 5, we show that both of these corollaries are sharp. In particular, the linear growth rate in Theorem 1.1 is optimal, in the sense that a superlinear growth rate does not suffice for showing that the set of ergodic measures is finite, and the technical condition of Theorem 1.2 (and in Theorem 3.1) on the existence of a point whose orbit closure is not uniquely ergodic can not be dropped.

#### 4. THE NATURAL CODING OF AN IET

Let  $k \geq 1$  be an integer and  $\pi$  be a permutation of  $\{1, \dots, k\}$ . Let  $I = [0, \lambda)$  be an interval and choose  $0 = \lambda_0 < \lambda_1 < \dots < \lambda_k = \lambda$ . The *interval exchange transformation*  $T: [0, \lambda) \rightarrow [0, \lambda)$  is defined to be the map that is an isometry on each subinterval  $[\lambda_{i-1}, \lambda_i)$  for  $i = 1, \dots, k$  and rearranges the order of these subintervals according to the permutation  $\pi$ . Without loss of generality, we can assume that  $k$  is the smallest number of subintervals needed to define the transformation  $T$  (otherwise we can join two consecutive subintervals into a single one). We refer to this interval exchange transformation as a *k-IET* or just an *IET* when  $k$  is clear from the context.

Given an interval exchange transformation, there is a natural coding by an associated dynamical system. For  $x \in I$ , define  $\mathbf{x} = (x_n) \in \{1, \dots, k\}^{\mathbb{N}}$  by setting

$$x_n = i \text{ if and only if } T^n x \in [\lambda_{i-1}, \lambda_i).$$

The *language* of  $\mathbf{x}$  is the set of all finite words that appear and the *natural coding* of the interval exchange transformation is the symbolic system, endowed with the shift, that has the same language as  $\mathbf{x}$ . The *natural symbolic cover* of an interval exchange transformation is the subshift that codes every  $x \in I$ , meaning it is the symbolic system, endowed with the shift, whose language consists of all finite words that arise in the orbit of any  $x \in I$ . While the image of the interval  $[0, \lambda)$  under this coding is invariant under the shift, it is not necessarily closed, and so we take its orbit closure to produce a semiconjugacy from the coding to the interval exchange. For further details, see [11, (Chapter 15, Section 5)].

If a point does not lie in the orbit of one of the endpoints of the subintervals defining the IET, then it has a unique preimage under the semiconjugacy, and otherwise it has at most two preimages corresponding to the coding of iterates.

If  $T$  is a minimal interval exchange transformation, then any  $x \in I$  gives rise to the same language and it suffices to take the orbit of a single point. More generally, the symbolic coding is not topologically conjugate to  $T$ , as up to countably many points may have multiple preimages (though a point can only have finitely many preimages). However, since the points with non-unique preimage can only support an atomic measure, it is a measure theoretic isomorphism for any nonatomic generic measure on  $X$ .

Namely, we claim that a generic measure for an interval exchange transformation lifts to a generic measure in the symbolic cover. A basic open set in the symbolic cover is a cylinder set and thus corresponds to an interval or a finite union of intervals in  $[0, \lambda)$ . Thus it suffices to check the claim for a finite interval  $J \subseteq [0, \lambda)$ . Let  $x \in [0, \lambda)$  be a generic point for the measure  $\mu$ . Choose continuous functions  $f$  and  $g$  on  $[0, \lambda)$  such that  $0 \leq f \leq \mathbf{1}_J \leq g$  and  $\int g d\mu - \varepsilon/2 \leq \mu(J) \leq \int f d\mu + \varepsilon/2$ . Then

$$\left| \frac{1}{N} \sum_{n=0}^{N-1} f(T^n x) - \int f d\mu \right| < \varepsilon/2$$

for all sufficiently large  $N$ , and the same holds for  $g$ . Thus

$$\frac{1}{N} \sum_{n=0}^{N-1} \mathbf{1}_J(T^n x) \leq \frac{1}{N} \sum_{n=0}^{N-1} g(T^n x) \leq \varepsilon/2 + \int g d\mu \leq \varepsilon + \int f d\mu \leq \varepsilon + \frac{1}{N} \sum_{n=0}^{N-1} \mathbf{1}_J(T^n x).$$

Thus the difference

$$\left| \mu(J) - \frac{1}{N} \sum_{n=0}^{N-1} \mathbf{1}_J(T^n x) \right| < \varepsilon.$$

Since this holds for all  $\varepsilon > 0$ , for any open set  $J \subset [0, \lambda)$ , we have

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=0}^{N-1} \mathbf{1}_J(T^n x) = \mu(J).$$

Write  $\phi: (X, \sigma) \rightarrow ([0, \lambda), T)$  for the factor map from the symbolic coding  $(X, \sigma)$  to the interval exchange  $([0, \lambda), T)$ . Let  $\mathcal{L}(X)$  denote the language of the coding and let  $\mu$  be a generic measure on  $([0, \lambda), T)$  with generic point  $x$ . Let  $x^* \in \phi^{-1}(x)$ . Then for any word  $w \in \mathcal{L}(X)$ ,

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=0}^{N-1} \mathbf{1}_{[w]}(\sigma^n x^*) = \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=0}^{N-1} \mathbf{1}_{\phi([w])}(T^n x) = \mu(\phi([w])),$$

since  $\phi([w])$  is a finite union of intervals. Since  $\mu$  is a nonatomic, generic measure, the pullback  $\phi^*(\mu([w])) = \phi^*(\mu(\phi^{-1}(\phi([w])))$  is also nonatomic, as only countably many points in  $([0, \lambda), T)$  have multiple pre-images and each of these only has finitely many preimages. (In other words, the pushforward of the pullback of the measure is the measure itself.) Thus a generic measure for the interval exchange transformation corresponds to a generic measure in the symbolic coding.

It is well known that an IET has linear complexity (see for example [8]). We include a proof for completeness:

**Proposition 4.1.** *The natural coding of a minimal  $k$ -IET has complexity*

$$P(n) \leq (k-1)n + 1.$$

*If the  $k$ -IET satisfies the infinite distinct orbits condition (IDOC), then the complexity is exactly  $P(n) = (k-1)n + 1$ .*

*Proof.* We proceed by induction on  $n$ . For  $n=1$ , this is the alphabet  $k$  and the result is clear. Assume that  $P(n) \leq (k-1)n + 1$ . Fixing a particular word of length  $n$ , the cylinder set defined by this word distinguishes an interval in the exchange, and by considering the cylinder sets associated to each word of length  $n$ , we obtain a partition of the exchange. Thus we have associated a partition  $\mathcal{I}$  of the exchange to the  $(k-1)n + 1$  words of length  $n$ , and this partition has  $(k-1)n + 2$  endpoints.

Furthermore, these endpoints all arise as iterates of the endpoints of the original  $k + 1$  endpoints of the interval exchange. Each of the  $k + 1$  original endpoints lies in some  $T(I)$ , where  $T$  is the exchange map and  $I$  is one of the intervals in the partition  $\mathcal{I}$ . We note that if the exchange satisfies the IDOC condition, then the endpoints arise as distinct iterates, each of the original endpoints lies in the interior of some  $T(I)$ , but without this condition there may be overlap in the iterates and this is only an upper bound.

Thus we have  $M \leq k - 1$  intervals in  $(T(I))_{I \in \mathcal{I}}$  which cover all of the original endpoints. These  $M$  intervals may each cover more than one of the original endpoints, say  $m$  of them, and there are at most  $m + 1$  distinct ways to continue the orbit of a word of length  $n$ . Thus in total, we have  $(k - 1)n + 1 - M + (k - 1) + M$  continuations, which is exactly the bound  $P(n + 1) \leq (k - 1)(n + 1) + 1$ .

If the exchange satisfies the IDOC condition, then as the endpoints arise as distinct iterates, we have that the complexity is exactly  $P(n) = (k - 1)n + 1$ .  $\square$

Combining this with Theorem 3.1, we have the statement of Theorem 1.4:

**Corollary** (Theorem 1.4). *For  $k > 2$ , a minimal  $k$ -IET has at most  $k - 2$  generic measures.*

### 5. SHARPNESS

In this section show that the bound in Theorem 3.1 is sharp. We recall the statement of Theorem 1.3 for convenience.

**Theorem** (Theorem 1.3). *Let  $d > 1$  be fixed. There exists a minimal subshift  $(X, \sigma)$  such that*

$$\begin{aligned} \liminf_{n \rightarrow \infty} \frac{P_X(n)}{n} &= d, \\ \limsup_{n \rightarrow \infty} \frac{P_X(n)}{n} &= d + 1, \end{aligned}$$

and  $X$  has exactly  $d$  ergodic measures.

Before we delve into the details of the construction, we outline the basic ideas involved. The ideas of this argument were partly inspired by a construction of a minimal and not uniquely ergodic subshift by Quas on mathoverflow [13] (see also Denker, Grillenberger, and Sigmund [6]).

Fixing  $d > 1$  and the alphabet  $\mathcal{A} = \{1, \dots, d\}$ , we inductively construct  $d$  sequences of words  $\{w_1^j\}_{j=1}^\infty, \{w_2^j\}_{j=1}^\infty, \dots, \{w_d^j\}_{j=1}^\infty$  in  $\mathcal{L}(\mathcal{A}^\mathbb{Z})$ . Roughly speaking, the procedure we use constructs the words in these sequences in the following (somewhat unusual) order:  $w_1^1, w_2^1, \dots, w_d^1, w_1^2, w_2^2, \dots, w_d^2, w_1^3, w_2^3, \dots, w_d^3, \dots$ . That is, we first construct the first word in each of the sequences, then construct the second word in each of the sequences, and so on. The words are constructed such that:

- (i) If  $i_1, i_2 \in \mathcal{A}$  and  $j_1 < j_2$ , then  $w_{i_1}^{j_1}$  occurs as a subword of  $w_{i_2}^{j_2}$  syndetically<sup>1</sup>, with gap size bounded by a constant that depends only on  $j_1$ .
- (ii) For any  $i \in \mathcal{A}$  and  $j \in \mathbb{N}$ , the frequency with which the letter  $i$  occurs in  $w_i^j$  (as a percentage of the length of  $w_i^j$ ) is greater than an absolute constant which is greater than  $1/2$ .

---

<sup>1</sup>A word  $v$  occurs syndetically in a word  $w$  with gap  $g$  if every subword of  $w$  of length  $g$  contains a copy of  $v$  as a sub-subword.

By taking a limit along a subsequence of  $\{w_1^j\}_{j=1}^\infty$ , we produce a semi-infinite word  $w_1^\infty$ , and taking its orbit closure under the shift  $\sigma$  and passing to the natural two sided extension, we obtain a closed subshift  $X \subset \mathcal{A}^\mathbb{Z}$ . It follows from the construction that  $(X, \sigma)$  is minimal and that  $w_i^j \in \mathcal{L}(X)$  for all  $i \in \mathcal{A}$  and  $j \in \mathbb{N}$ . For fixed  $i \in \mathcal{A}$ , there are arbitrarily long words in  $\mathcal{L}(X)$  for which the frequency of the letter  $i$  is at least (a constant greater than)  $1/2$  and so the system  $(X, \sigma)$  has an ergodic measure assigning the cylinder set  $[i]$  measure greater than  $1/2$ . Thus  $(X, \sigma)$  has at least  $|\mathcal{A}| = d$  ergodic measures. By carefully choosing the lengths of the words, we further show that the system  $(X, \sigma)$  satisfies the desired upper and lower bounds on the complexity. Applying Theorem 3.1, it follows that  $(X, \sigma)$  has at most  $d$  ergodic measures, and so exactly  $d$  ergodic measures.

We now make these ideas precise:

*Proof of Theorem 1.3.* Set  $\mathcal{A} := \{1, 2, \dots, d\}$ . Choose  $\kappa_1, \kappa_2, \dots$  to be a sequence of real numbers in  $(0, 1)$  such that

$$\prod_{j=1}^{\infty} \kappa_j > 1/2$$

choose  $\delta_1, \delta_2, \dots$  to be a strictly decreasing sequence of real numbers in  $(0, 1)$  such that  $\lim_{j \rightarrow \infty} \delta_j = 0$ .

**Step 1.** (*Construction of the sequences  $\{w_1^j\}_{j=1}^\infty, \dots, \{w_d^j\}_{j=1}^\infty$* ): Define the word

$$w_1^1 := \underbrace{11 \cdots 1}_{\text{length } N_{(1,1)}^{[1]}} 234 \cdots d,$$

where  $N_{(1,1)}^{[1]} \in \mathbb{N}$  is chosen such that  $N_{(1,1)}^{[1]} > \kappa_1 |w_1^1|$ . Define the words  $s_1^1 := w_1^1$  and

$$p_1^1 := \underbrace{11 \cdots 1}_{\text{length } N_{(1,1)}^{[1]}} .$$

The word  $s_1^1$  represents the *suffix* of  $w_1^1$  that starts with the beginning of  $w_1^1$  and  $p_1^1$  denotes the *prefix* of  $w_1^1$  consisting of its initial block of 1's (the nomenclature of suffix and prefix is further clarified in the inductive construction, but note that the previous level suffix becomes the beginning of the next level prefix). We refer to the groupings by a length  $N_{j,k}^{[n]}$  as a *block* in the word (thus  $p_1^1$  has a single block), and the change from one block to the following block as a *transition* (the word  $w_2^1$  has one transition) and we refer to words that are not entirely within one block as *transition words*.

Next define the word

$$w_2^1 := \underbrace{s_1^1 w_1^1 w_1^1 \cdots w_1^1 p_1^1}_{\text{length } N_{(2,1)}^{[1]}} \underbrace{222 \cdots 2}_{\text{length } N_{(2,2)}^{[1]}} \underbrace{333 \cdots 3}_{\text{length } N_{(2,3)}^{[1]}} \cdots \underbrace{ddd \cdots d}_{\text{length } N_{(2,d)}^{[1]}} ,$$

where  $N_{(2,1)}^{[1]}, \dots, N_{(2,d)}^{[1]} \in \mathbb{N}$  are chosen such that

$$(18) \quad \begin{aligned} |w_1^1| &< (\delta_1)^2 \cdot N_{(2,1)}^{[1]} < (\delta_1)^4 \cdot N_{(2,2)}^{[1]} < (\delta_1)^6 \cdot N_{(2,d-1)}^{[1]} < (\delta_1)^8 \cdot N_{(2,d-2)}^{[1]} \\ &< \cdots < (\delta_1)^{2d-2} \cdot N_{(2,3)}^{[1]} < (\delta_1)^{2d} \cdot N_{(2,2)}^{[1]} \end{aligned}$$

and  $N_{(2,2)}^{[1]} > \kappa_1 |w_2^1|$ . The ordering of the lengths  $N_{(2,k)}^{[1]}$  is important, with the index  $k$  passing from 1 to  $d$  and then down to  $d-1$  and continuing cycling step by step down to 2. The choice of the lengths is used only in estimating the growth of  $P_X(n)$ ; the exact choices of the lengths and the estimates of (18) can be ignored for a first reading of Steps 1 and 2.

Further note that the initial portion of  $w_2^1$  is a concatenation of the word  $w_1^1$  with itself a large number of times, followed by its prefix word  $p_1^1$ . We include  $p_1^1$  so that the word  $w_1^1 p_1^1 2$  has already occurred as a subword of the concatenation  $w_1^1 w_1^1 \cdots w_1^1$ , allowing us to make the transition in  $w_2^1$  from the block of  $w_1^1$ 's to the block of 2's while keeping the possible new words as low as possible. We iterate this technique at each step of the construction.

Next we define the word  $w_3^1$ . To do this, we make use of two auxiliary words. Let

$$p_2^1 := \underbrace{s_1^1 w_1^1 w_1^1 \cdots w_1^1 p_1^1}_{\text{length } N_{(2,1)}^{[1]}} \underbrace{22 \cdots 2}_{\text{length } N_{(2,2)}^{[1]}}$$

be the prefix of  $w_2^1$  that includes everything through the block of 2's and let

$$s_2^1 := \underbrace{22 \cdots 2}_{\text{length } N_{(2,2)}^{[1]}} \underbrace{33 \cdots 3}_{\text{length } N_{(2,3)}^{[1]}} \cdots \underbrace{dd \cdots d}_{\text{length } N_{(2,d)}^{[1]}}$$

be the suffix of  $w_2^1$  that starts with the block of 2's. Finally, we define

$$w_3^1 := \underbrace{s_1^1 w_1^1 w_1^1 \cdots w_1^1 p_1^1}_{\text{length } N_{(3,1)}^{[1]}} \underbrace{s_2^1 w_2^1 w_2^1 \cdots w_2^1 p_2^1}_{\text{length } N_{(3,2)}^{[1]}} \underbrace{33 \cdots 3}_{\text{length } N_{(3,3)}^{[1]}} \cdots \underbrace{dd \cdots d}_{\text{length } N_{(3,d)}^{[1]}} ,$$

where  $N_{(3,1)}^{[1]}, N_{(3,2)}^{[1]}, \dots, N_{(3,d)}^{[1]} \in \mathbb{N}$  are chosen such that

$$(19) \quad \begin{aligned} |w_2^1| &< (\delta_1)^2 \cdot N_{(3,2)}^{[1]} < (\delta_1)^4 \cdot N_{(3,1)}^{[1]} < (\delta_1)^6 \cdot N_{(3,d)}^{[1]} < (\delta_1)^8 \cdot N_{(3,d-1)}^{[1]} \\ &< \cdots < (\delta_1)^{2d-2} \cdot N_{(3,4)}^{[1]} < (\delta_1)^{2d} \cdot N_{(3,3)}^{[1]} \end{aligned}$$

and  $N_{(3,3)}^{[1]} > \kappa_1 |w_3^1|$ . Again, the initial block of  $w_3^1$  is periodic (based on the word  $w_1^1$ ), with the period ending when its pattern dictates the next letter should be 2. The second block is periodic (based on the word  $w_2^1$ ) that begins with a block of 2's and ends when its pattern dictates the next collection of letters should be a block of 3's ( $w_3^1$  then continues with a block of 3's). As in the previous step, this gives two transitions in  $w_3^1$ : first from a periodic block of  $w_1^1$ 's to a periodic block of  $w_2^1$ 's and second from a periodic block of  $w_2^1$ 's to a block of 3's. The choice of the prefixes and suffixes guarantee that these transitions introduce the smallest possible number of new words (the analysis of this number of new words is made precise in Step 3, where we analyze the growth of the complexity  $P_X(n)$ ).

We continue inductively. Let  $i < d$  be fixed and suppose we have constructed words  $w_1^1, \dots, w_i^1, p_1^1, \dots, p_i^1, s_1^1, \dots, s_i^1$ . Define the word (note that many of the

words we define are too long to fit in a single line, and the line break is arbitrary)

$$w_{i+1}^1 := \underbrace{s_1^1 w_1^1 w_1^1 \cdots w_1^1 p_1^1}_{\text{length } N_{(i+1,1)}^{[1]}} \underbrace{s_2^1 w_2^1 w_2^1 \cdots w_2^1 p_2^1 \cdots}_{\text{length } N_{(i+1,2)}^{[1]}} \underbrace{s_i^1 w_i^1 w_i^1 \cdots w_i^1 p_i^1}_{\text{length } N_{(i+1,i)}^{[1]}} \underbrace{(i+1)(i+1) \cdots (i+1)}_{\text{length } N_{(i+1,i+1)}^{[1]}} \cdots \underbrace{dd \cdots d}_{\text{length } N_{(i+1,d)}^{[1]}},$$

where  $N_{(i+1,1)}^{[1]}, \dots, N_{(i+1,d)}^{[1]} \in \mathbb{N}$  are chosen such that

$$(20) \quad \begin{aligned} |w_i^1| &< (\delta_1)^2 \cdot N_{(i+1,i)}^{[1]} < (\delta_1)^4 \cdot N_{(i+1,i-1)}^{[1]} < (\delta_1)^6 \cdot N_{(i+1,i-2)}^{[1]} \\ &< (\delta_1)^8 \cdot N_{(i+1,i-3)}^{[1]} < \cdots < (\delta_1)^{2i} \cdot N_{(i+1,1)}^{[1]} < (\delta_1)^{2i+2} \cdot N_{(i+1,d)}^{[1]} \\ &< (\delta_1)^{2i+4} \cdot N_{(i+1,d-1)}^{[1]} < \cdots < (\delta_1)^{2d-2} \cdot N_{(i+1,i+2)}^{[1]} < (\delta_1)^{2d} \cdot N_{(i+1,i+1)}^{[1]} \end{aligned}$$

and  $N_{(i+1,i+1)}^{[1]} > \kappa_1 |w_{i+1}^1|$ . Again, the lengths are chosen to control the growth of the complexity, and the index  $k$  in  $N_{(i+1,k)}^{[1]}$  is taken in a cyclical order. Also define a prefix and a suffix of  $w_{i+1}^1$  by

$$\begin{aligned} p_{i+1}^1 &:= \underbrace{s_1^1 w_1^1 w_1^1 \cdots w_1^1 p_1^1}_{\text{length } N_{(i+1,1)}^{[1]}} \underbrace{s_2^1 w_2^1 w_2^1 \cdots w_2^1 p_2^1 \cdots}_{\text{length } N_{(i+1,2)}^{[1]}} \underbrace{s_i^1 w_i^1 w_i^1 \cdots w_i^1 p_i^1}_{\text{length } N_{(i+1,i)}^{[1]}} \underbrace{(i+1)(i+1) \cdots (i+1)}_{\text{length } N_{(i+1,i+1)}^{[1]}} \\ s_{i+1}^1 &:= \underbrace{(i+1)(i+1) \cdots (i+1)}_{\text{length } N_{(i+1,i+1)}^{[1]}} \underbrace{(i+2)(i+2) \cdots (i+2)}_{\text{length } N_{(i+1,i+2)}^{[1]}} \cdots \underbrace{dd \cdots d}_{\text{length } N_{(i+1,d)}^{[1]}} \end{aligned}$$

so that  $p_{i+1}^1$  is the prefix of  $w_{i+1}^1$  that includes everything through the block of  $(i+1)$ 's and  $s_{i+1}^1$  is the suffix of  $w_{i+1}^1$  that begins with the block of  $(i+1)$ 's. By induction, this defines words  $w_1^1, \dots, w_d^1, p_1^1, \dots, p_d^1, s_1^1, \dots, s_d^1$ .

For each  $i \in \mathcal{A}$ , it follows immediately from the construction that:

- (a) Every letter in  $\mathcal{A}$  appears in  $w_i^1$ ;
- (b) The frequency with which the letter  $i$  occurs in  $w_i^1$  is at least  $\kappa_1$ , by choice of the integer  $N_{(i+1,i+1)}^{[1]}$  and the relations given in (20).

We continue to define the sequences of words inductively. Assuming that we have already defined words  $w_1^j, w_2^j, \dots, w_d^j, p_1^j, \dots, p_d^j, s_1^j, \dots, s_d^j$ , we define

$$w_1^{j+1} := \underbrace{s_1^j w_1^j w_1^j \cdots w_1^j p_1^j}_{\text{length } N_{(1,1)}^{[j+1]}} \underbrace{s_2^j w_2^j w_2^j \cdots w_2^j p_2^j \cdots}_{\text{length } N_{(1,2)}^{[j+1]}} \underbrace{s_d^j w_d^j w_d^j \cdots w_d^j p_d^j}_{\text{length } N_{(1,d)}^{[j+1]}}$$

where

$$(21) \quad \begin{aligned} |w_d^j| &< (\delta_{j+1})^2 \cdot N_{(1,d)}^{[j+1]} < (\delta_{j+1})^4 \cdot N_{(1,d-1)}^{[j+1]} < (\delta_{j+1})^6 \cdot N_{(1,d-2)}^{[j+1]} \\ &< (\delta_{j+1})^8 \cdot N_{(1,d-3)}^{[j+1]} < \cdots < (\delta_{j+1})^{2d-2} \cdot N_{(1,2)}^{[j+1]} < (\delta_{j+1})^{2d} \cdot N_{(1,1)}^{[j+1]} \end{aligned}$$

and  $N_{(1,1)}^{[j+1]} > \kappa_{j+1} |w_1^{j+1}|$ . Define prefixes and suffixes by

$$\begin{aligned} p_1^{j+1} &:= \underbrace{s_1^j w_1^j w_1^j \cdots w_1^j p_1^j}_{\text{length } N_{(1,1)}^{[j]}} \\ s_1^{j+1} &:= w_1^{j+1}. \end{aligned}$$

We have analogs of properties (a) and (b) for the base case of the construction: each of the words  $w_1^j, w_2^j, \dots, w_d^j$  occurs as a subword of  $w_1^{j+1}$  and the frequency with which the letter 1 occurs in  $w_1^{j+1}$  is at least  $\prod_{k=1}^{j+1} \kappa_k$ , provided that the frequency with which it occurs in  $w_1^j$  was at least  $\prod_{k=1}^j \kappa_k$ .

Continuing inductively, for  $i < d$ , we define the word (note the change in superscript half way through)

$$w_{i+1}^{j+1} := \underbrace{s_1^{j+1} w_1^{j+1} w_1^{j+1} \dots w_1^{j+1} p_1^{j+1}}_{\text{length } N_{(i+1,1)}^{[j+1]}} \underbrace{s_2^{j+1} w_2^{j+1} \dots w_2^{j+1} p_2^{j+1} \dots}_{\text{length } N_{(i+1,2)}^{[j+1]}} \dots$$

$$\underbrace{s_i^{j+1} w_i^{j+1} \dots w_i^{j+1} p_i^{j+1}}_{\text{length } N_{(i+1,i)}^{[j+1]}} \underbrace{s_{i+1}^j w_{i+1}^j \dots w_{i+1}^j p_{i+1}^j}_{\text{length } N_{(i+1,i+1)}^{[j+1]}} \dots \underbrace{s_d^j w_d^j \dots w_d^j p_d^j}_{\text{length } N_{(i+1,d)}^{[j+1]}}$$

where

$$(22) \quad \begin{aligned} |w_i^{j+1}| &< (\delta_{j+1})^2 \cdot N_{(i+1,i)}^{[j+1]} < (\delta_{j+1})^4 \cdot N_{(i+1,i-1)}^{[j+1]} < (\delta_{j+1})^6 \cdot N_{(i+1,i-2)}^{[j+1]} \\ &< (\delta_{j+1})^8 \cdot N_{(i+1,i-3)}^{[j+1]} < \dots < (\delta_{j+1})^{2i} \cdot N_{(i+1,1)}^{[j+1]} \\ &< (\delta_{j+1})^{2i+2} \cdot N_{(i+1,d)}^{[j+1]} < (\delta_{j+1})^{2i+4} \cdot N_{(i+1,d-1)}^{[j+1]} < \dots \\ &< (\delta_{j+1})^{2d-2} \cdot N_{(i+1,i+2)}^{[j+1]} < (\delta_{j+1})^{2d} \cdot N_{(i+1,i+1)}^{[j+1]} \end{aligned}$$

and  $N_{(i+1,i+1)}^{[j+1]} > \kappa_{j+1} |w_{i+1}^{j+1}|$ , and define the prefixes and suffixes by

$$p_{i+1}^{j+1} := \underbrace{s_1^{j+1} w_1^{j+1} w_1^{j+1} \dots w_1^{j+1} p_1^{j+1}}_{\text{length } N_{(i+1,1)}^{[j+1]}} \underbrace{s_2^{j+1} w_2^{j+1} \dots w_2^{j+1} p_2^{j+1} \dots}_{\text{length } N_{(i+1,2)}^{[j+1]}} \dots$$

$$\underbrace{s_i^{j+1} w_i^{j+1} \dots w_i^{j+1} p_i^{j+1}}_{\text{length } N_{(i+1,i)}^{[j+1]}} \underbrace{s_{i+1}^j w_{i+1}^j \dots w_{i+1}^j p_{i+1}^j}_{\text{length } N_{(i+1,i+1)}^{[j+1]}}$$

and

$$s_{i+1}^{j+1} := \underbrace{s_{i+1}^j w_{i+1}^j \dots w_{i+1}^j p_{i+1}^j}_{\text{length } N_{(i+1,i+1)}^{[j+1]}} \underbrace{s_{i+2}^j w_{i+2}^j \dots w_{i+2}^j p_{i+2}^j \dots}_{\text{length } N_{(i+1,i+2)}^{[j+1]}} \dots \underbrace{s_d^j w_d^j \dots w_d^j p_d^j}_{\text{length } N_{(i+1,d)}^{[j+1]}}$$

Again, we point out that the words  $w_1^j, w_2^j, \dots, w_d^j$  occur as subwords of  $w_{i+1}^{j+1}$ , and the frequency with which the letter  $i+1$  occurs in  $w_{i+1}^{j+1}$  is at least  $\prod_{k=1}^{j+1} \kappa_k$ , provided that the frequency with which it occurs in  $w_{i+1}^j$  was at least  $\prod_{k=1}^j \kappa_k$ .

By induction, we obtain sequences  $\{w_1^j\}_{j=1}^\infty, \{w_2^j\}_{j=1}^\infty, \dots, \{w_d^j\}_{j=1}^\infty$  satisfying:

(a) For any  $j > 2$ , any  $1 \leq k < j - 2$ , and any  $i_1, i_2 \in \mathcal{A}$ , the word  $w_{i_1}^k$  occurs in each of the words  $w_1^{k+1}, w_2^{k+1}, \dots, w_d^{k+1}$  by construction. Each of the words

$$(23) \quad w_1^{k+2}, \dots, w_d^{k+2}, p_1^{k+2}, \dots, p_d^{k+2}, s_1^{k+2}, \dots, s_d^{k+2},$$

contains at least one of the words  $w_1^{k+1}, w_2^{k+1}, \dots, w_d^{k+1}$  and so  $w_{i_1}^k$  occurs in each of these words. Since  $w_{i_2}^j$  is a concatenation the words appearing in (23), it follows that  $w_{i_1}^k$  occurs in  $w_{i_2}^j$  syndetically, and the maximal gap length is at most

$$g_k := \max\{|w_l^{k+2}| : l \in \mathcal{A}\} \cup \{|p_l^{k+2}| : l \in \mathcal{A}\} \cup \{|s_l^{k+2}| : l \in \mathcal{A}\} = |w_d^{k+2}|;$$

- (b) For any  $i \in \mathcal{A}$  and any  $j \in \mathbb{N}$ , the frequency with which the letter  $i$  occurs as a subword of  $w_i^j$  is at least  $\prod_{k=1}^j \kappa_k \geq \prod_{k=1}^{\infty} \kappa_k > 1/2$ .

We further note that given the freedom with which the lengths are chosen, we can assume that  $N_{(i,k)}^{[j]}$  divides  $N_{(i,k)}^{[j+1]}$  for all  $i, k \in \mathcal{A}$  and all  $j \in \mathbb{N}$ . We make this assumption for the remainder of the proof.

**Step 2.** (*Construction and ergodic properties of the subshift  $(X, \sigma)$* ): Observe that  $w_1^j$  is the leftmost subword of  $w_1^{j+1}$  for all  $j \in \mathbb{N}$ , and so we can define a (one-sided) infinite word  $w_1^\infty$  by declaring that for all  $j$ , the leftmost subword of  $w_1^\infty$  of length  $|w_1^j|$  is  $w_1^j$ . Then for any  $i \in \mathcal{A}$  and any  $j \in \mathbb{N}$ , the word  $w_i^j$  occurs as a subword of  $w_1^\infty$  syndetically. Moreover, every subword of  $w_1^\infty$  occurs as a sub-subword of  $w_1^j$  for some  $j$ . Therefore all subwords of  $w_1^\infty$  occur syndetically.

Let  $X \subset \mathcal{A}^{\mathbb{Z}}$  be the set of all bi-infinite sequences whose language is comprised only of subwords of  $w_1^\infty$ , meaning it is the natural extension of the closure of  $w_1^\infty$  under  $\sigma$ . Since all words in  $\mathcal{L}(X)$  occur syndetically in every element of  $X$ ,  $(X, \sigma)$  is minimal. Moreover,  $w_i^j \in \mathcal{L}(X)$  for all  $i \in \mathcal{A}$  and  $j \in \mathbb{N}$ . Therefore, for fixed  $i \in \mathcal{A}$ , there are arbitrarily long words in  $\mathcal{L}(X)$  for which the frequency with which the letter  $i$  occurs is at least  $\prod_{k=1}^{\infty} \kappa_k > 1/2$ . For each such word  $w_i^n$ , if  $y_n \in [w_i^n]$  and

$$\mu_n := \frac{1}{n} \sum_{k=0}^{|w_i^n|-1} \delta_{\sigma^k(y_n)},$$

then any weak-\* limit point  $\nu$  of the sequence  $(\mu_n)$  is a  $\sigma$ -invariant probability measure supported on  $X$  such that  $\nu([i]) > 1/2$  (such a construction of  $\nu$  is, for example, a standard way to prove the Krylov-Bogolyubov Theorem). Consequently, as the ergodic measures are the extreme points in the (convex) set of invariant probability measures on  $X$ , there exists an ergodic measure  $\mu_i$  supported on  $X$  for which  $\mu_i([i]) > 1/2$ . It follows that  $\mu_i([j]) < 1/2$  for all  $j \neq i$  and so  $\mu_j \neq \mu_i$  for any  $j \neq i$ . Thus  $(X, \sigma)$  has at least  $d$  ergodic measures. If we can show that

$$\liminf_{n \rightarrow \infty} \frac{P_X(n)}{n} < d + 1,$$

then there are at most  $d$  ergodic measures by Theorem 3.1; hence exactly  $d$ .

Thus we are left with showing that:

$$\begin{aligned} \liminf_{n \rightarrow \infty} \frac{P_X(n)}{n} &= d, \\ \limsup_{n \rightarrow \infty} \frac{P_X(n)}{n} &= d + 1. \end{aligned}$$

**Step 3.** (*Analysis of the growth rate of  $P_X(n)$* ): Let  $n > |w_1^2|$  be a fixed integer. We estimate the number of words in  $\mathcal{L}_n(X)$ , which is, by definition,  $P_X(n)$ . By construction,

$$|w_1^1| < |w_2^1| < \dots < |w_d^1| < |w_1^2| < |w_2^2| < \dots < |w_d^2| < |w_1^3| < \dots$$

We make the convention that  $w_{d+1}^j := w_1^{j+1}$ ,  $w_{d+2}^j := w_2^{j+1}$ , and so on (with the analogous convention for  $N_{(i_1, i_2)}^{[j]}$  when  $i_2 > d$ ). Therefore, there exist  $i_1 \in \mathcal{A}$  and  $j_1 \in \mathbb{N}$  such that

$$|w_{i_1}^{j_1}| \leq n < |w_{i_1+1}^{j_1}|.$$



With this convention, observe that

$$\begin{aligned}
 w_{i_1+1}^{j_1+1} &:= \underbrace{s_1^{j_1+1} w_1^{j_1+1} \cdots w_1^{j_1+1} p_1^{j_1+1}}_{\text{length } N_{(i_1+1,1)}^{[j_1+1]}} \underbrace{s_2^{j_1+1} w_2^{j_1+1} \cdots w_2^{j_1+1} p_2^{j_1+1} \cdots}_{\text{length } N_{(i_1+1,2)}^{[j_1+1]}} \\
 &\quad \underbrace{s_{i_1}^{j_1+1} w_{i_1}^{j_1+1} \cdots w_{i_1}^{j_1+1} p_{i_1}^{j_1+1}}_{\text{length } N_{(i_1+1,i_1)}^{[j_1+1]}} \underbrace{s_{i_1+1}^{j_1} w_{i_1+1}^{j_1} \cdots w_{i_1+1}^{j_1} p_{i_1+1}^{j_1}}_{\text{length } N_{(i_1+1,i_1+1)}^{[j_1]}} \cdots \underbrace{s_d^{j_1} w_d^{j_1} \cdots w_d^{j_1} p_d^{j_1}}_{\text{length } N_{(i_1+1,d)}^{[j_1]}}
 \end{aligned}$$

where

$$(24) \quad n < |w_{i_1+1}^{j_1+1}| < |w_{i_1+2}^{j_1+1}| < \cdots < |w_d^{j_1+1}| < |w_1^{j_1+1}| < \cdots < |w_{i_1}^{j_1+1}|.$$

It follows from the construction that if  $i_2 \in \mathcal{A}$  and  $j_2 \in \mathbb{N}$  is such that  $|w_{i_2}^{j_2}| \geq |w_{i_1+1}^{j_1+1}|$ , then  $w_{i_2}^{j_2}$  can also be written as a concatenation of words from the sets

$$\{w_1^{j_1+1}, w_2^{j_1+1}, \dots, w_{i_1}^{j_1+1}, w_{i_1+1}^{j_1}, w_{i_1+2}^{j_1}, \dots, w_d^{j_1}\} = \{w_{i_1}^{j_1+1}, w_{i_1+1}^{j_1}, \dots, w_{i_1+d-1}^{j_1}\}$$

and

$$\{p_i^{j_1+1} : 1 \leq i \leq i_1\} \cup \{p_i^{j_1} : i_1 < i \leq d\} \cup \{s_i^{j_1+1} : 1 \leq i \leq i_1\} \cup \{s_i^{j_1} : i_1 < i \leq d\}.$$

Moreover, there are restrictions on the order in which these words may be concatenated in  $w_{i_2}^{j_2}$ :

- (i) If  $i_1 + 1 \leq i < i_1 + d$ , then the only words that may be concatenated with  $w_i^{j_1}$  are  $w_i^{j_1}$  itself and  $p_i^{j_1} s_{i+1}^{j_1} w_{i+1}^{j_1}$ ;
- (ii) The only words that may be concatenated to the right end of  $w_{i_1+d}^{j_1}$  (=  $w_{i_1}^{j_1+1}$ ) are  $w_{i_1}^{j_1+1}$  itself and  $p_{i_1}^{j_1+1} s_{i_1+1}^{j_1} w_{i_1+1}^{j_1}$ .

Therefore, by (24), the only words of length  $n$  that appear as subwords of  $w_{i_2}^{j_2}$  are those which appear as subwords of words from the set:

$$(25) \quad \left\{ w_i^{j_1} w_i^{j_1} : i_1 < i \leq i_1 + d \right\} \cup \left\{ w_i^{j_1} p_i^{j_1} s_{i+1}^{j_1} w_{i+1}^{j_1} : i_1 < i < i_1 + d \right\} \cup \left\{ w_{i_1+d}^{j_1} p_{i_1+d}^{j_1} s_{i_1+1}^{j_1} w_{i_1+1}^{j_1} \right\},$$

with superscripts following the convention that if the subscript is larger than  $d$ , increment the superscript by 1. Since all words in  $\mathcal{L}_n(X)$  occur as subwords of  $w_1^{j_2}$  for all sufficiently large  $j_2$ , we have that all words in  $\mathcal{L}_n(X)$  appear as subwords of the  $2d$  words in the set in (25).

We now analyze the words that appear in (25) by decomposing them into words of length comparable to  $n$ . By construction, if  $1 \leq m \leq d$  then  $w_{i_1+m}^{j_1}$  can be written as a concatenation of words from the set (recall the divisibility of the

lengths assumed at the end of Step 1)

(26)

$$\begin{aligned} & \{w_{i_1+1}^{j_1}\} \cup \bigcup_{k=2}^d \left\{ \underbrace{s_{i_1+k-1}^{j_1-1} w_{i_1+k-1}^{j_1-1} \cdots w_{i_1+k-1}^{j_1-1} p_{i_1+k-1}^{j_1-1}}_{N_{(i_1+k, i_1+k-1)}^{[j_1]}}, \underbrace{s_{i_1+k}^{j_1-1} w_{i_1+k}^{j_1-1} \cdots w_{i_1+k}^{j_1-1} p_{i_1+k}^{j_1-1}}_{N_{(i_1+k, i_1+k)}^{[j_1]}}, \right. \\ & \quad \left. \underbrace{s_{i_1+k+1}^{j_1-1} w_{i_1+k+1}^{j_1-1} \cdots w_{i_1+k+1}^{j_1-1} p_{i_1+k+1}^{j_1-1}}_{N_{(i_1+k, i_1+k+1)}^{[j_1]}}, \dots, \underbrace{s_{d-1}^{j_1-1} w_{d-1}^{j_1-1} \cdots w_{d-1}^{j_1-1} p_{d-1}^{j_1-1}}_{N_{(i_1+k, d-1)}^{[j_1]}}, \right. \\ & \quad \left. \underbrace{s_d^{j_1-1} w_d^{j_1-1} \cdots w_d^{j_1-1} p_d^{j_1-1}}_{N_{(i_1+k, d)}^{[j_1]}}, \underbrace{s_1^{j_1} w_1^{j_1} \cdots w_1^{j_1} p_1^{j_1}}_{N_{(i_1+k, 1)}^{[j_1]}}, \underbrace{s_2^{j_1} w_2^{j_1} \cdots w_2^{j_1} p_2^{j_1}}_{N_{(i_1+k, 2)}^{[j_1]}}, \dots, \underbrace{s_{i_1}^{j_1} w_{i_1}^{j_1} \cdots w_{i_1}^{j_1} p_{i_1}^{j_1}}_{N_{(i_1+k, i_1)}^{[j_1]}} \right\} \end{aligned}$$

obeying the analogous rules for concatenation ( $w_{i_1+1}^{j_1}$  may be concatenated with itself or any block that begins with  $s_{i_1+2}^{j_1}$ , and any word that ends with  $p_{i_1+k}^{j_1}$  may be concatenated with a word that begins  $s_{i_1+k+1}^{j_1}$ , again understood cyclically). In turn, the words of length  $n$  that occur in words produced from this set are, themselves, words that occur when words from the set

$$\begin{aligned} (27) \quad & \left\{ w_{i_1+1}^{j_1}, \underbrace{s_{i_1+2}^{j_1-1} w_{i_1+2}^{j_1-1} \cdots w_{i_1+2}^{j_1-1} p_{i_1+2}^{j_1-1}}_{N_{(i_1+2, i_1+2)}^{[j_1]}}, \dots, \right. \\ & \quad \left. \underbrace{s_d^{j_1-1} w_d^{j_1-1} \cdots w_d^{j_1-1} p_d^{j_1-1}}_{N_{(i_1+2, d)}^{[j_1]}}, \underbrace{s_1^{j_1} w_1^{j_1} \cdots w_1^{j_1} p_1^{j_1}}_{N_{(i_1+2, 1)}^{[j_1]}}, \dots, \underbrace{s_{i_1}^{j_1} w_{i_1}^{j_1} \cdots w_{i_1}^{j_1} p_{i_1}^{j_1}}_{N_{(i_1+2, i_1)}^{[j_1]}} \right\} \end{aligned}$$

are concatenated (with the analogous rules for concatenation). Moreover,

$$(28) \quad |w_{i_1+2}^{j_1-1}| < |w_{i_1+3}^{j_1-1}| < \cdots < |w_d^{j_1-1}| < |w_1^{j_1}| < \cdots < |w_{i_1}^{j_1}| \leq n < |w_{i_1+1}^{j_1}|$$

and

$$(29) \quad n < |w_{i_1+1}^{j_1}| < N_{(i_1+2, m)}^{[j_1]}$$

for all  $1 \leq m \leq d$  (again, if  $i_1 + 2 > d$  then increment the superscript of  $N_{(i_1+2, m)}^{[j_1]}$  by one and reduce the subscript by  $d$ ). In particular, every word in the set (25) can be obtained by concatenating words from the set (26) and any word of length  $n$  that occurs in a word in (25) can also be found in a word obtained by concatenating words from (27).

For  $i_1 + 2 \leq i < i_1 + d + 1$ , define

$$q_i := \cdots w_i^{j_1-1} w_i^{j_1-1} w_i^{j_1-1} p_i^{j_1-1} s_{i+1}^{j_1-1} w_{i+1}^{j_1-1} w_{i+1}^{j_1-1} w_{i+1}^{j_1-1} \cdots$$

to be the bi-infinite word whose restriction to the set of nonnegative indices is the infinite concatenation  $s_{i+1}^{j_1-1} w_{i+1}^{j_1-1} w_{i+1}^{j_1-1} w_{i+1}^{j_1-1} \cdots$ , and whose restriction to the set of negative indices is the infinite concatenation  $\cdots w_i^{j_1-1} w_i^{j_1-1} w_i^{j_1-1} p_{i+1}^{j_1}$  (note that the dependence of  $q_i$  on  $j_1$  suppressed in our notation). Similarly define

$$q_{i_1+d+1} := \cdots w_{i_1+1}^{j_1} w_{i_1+1}^{j_1} w_{i_1+1}^{j_1} p_{i_1+1}^{j_1} s_{i_1+2}^{j_1-1} w_{i_1+2}^{j_1-1} w_{i_1+2}^{j_1-1} w_{i_1+2}^{j_1-1} \cdots$$

The set of words length  $n$  that arise by concatenating words from the set (27) is precisely the set of words of length  $n$  that appear in  $q_{i_1+1}, q_{i_1+2}, \dots, q_{i_1+d}$ , by (29).

By the estimates in (19), (20), (21), and (22), we have that

$$|w_i^{j_1-1}| < \delta_{j_1} \cdot |w_{i_1}^{j_1}| \leq \delta_{j_1} \cdot n$$

for all  $i_1 + 1 < i < d + i_1$ . It follows that:

- (i) If  $i_1 + 1 < i < i_1 + d - 1$ , then the number of words of  $q_i$  of length  $n$  is at least  $n + 1$  (since  $q_i$  is aperiodic) and at most  $n + 2\delta_{j_1}n$ , as there are at most  $\delta_{j_1}n$  words in each block (periodic part) of  $q_i$  and at most  $n$  transition words obtained from words that overlap the origin;
- (ii) The number of words of  $q_{i_1+d-1}$  of length  $n$  is at least  $n + 1$  and at most  $n + \delta_{j_1}n + |w_{i_1}^{j_1}|$ ;
- (iii) The only new words of  $q_{i_1+d}$  are the  $n + 1$  transition words which appear in  $\underbrace{w_{i_1}^{j_1} \cdots w_{i_1}^{j_1} p_{i_1}^{j_1}}_{N_{(i_1+2, i_1)}^{[j_1]}} s_{i_1+1}^{j_1} w_{i_1+1}^{j_1}$  as well as words that appear in  $w_{i_1+1}^{j_1} w_{i_1+1}^{j_1}$ ;
- (iv) The only new words of  $q_{i_1+d+1}$  are the  $n + 1$  transition words which appear in  $w_{i_1+1}^{j_1} p_{i_1+1}^{j_1} \underbrace{s_{i_1+2}^{j_1-1} w_{i_1+2}^{j_1-1} \cdots w_{i_1+2}^{j_1-1}}_{N_{(i_1+2, i_1+2)}^{[j_1]}}$ .

Thus we are left with counting subwords of  $w_{i_1+1}^{j_1} w_{i_1+1}^{j_1}$  that have not already appeared.

Write

$$w_{i_1+1}^{j_1} := \underbrace{s_1^{j_1} w_1^{j_1} \cdots w_1^{j_1} p_1^{j_1}}_{\text{length } N_{(i_1+1, 1)}^{[j_1]}} \underbrace{s_2^{j_1} w_2^{j_1} \cdots w_2^{j_1} p_2^{j_1} \cdots}_{\text{length } N_{(i_1+1, 2)}^{[j_1]}} \underbrace{s_{i_1+1}^{j_1-1} w_{i_1+1}^{j_1-1} \cdots w_{i_1+1}^{j_1-1} p_{i_1+1}^{j_1-1} \cdots}_{\text{length } N_{(i_1+1, i_1+1)}^{[j_1]}} \underbrace{s_d^{j_1-1} w_d^{j_1-1} \cdots w_d^{j_1-1} p_d^{j_1-1}}_{\text{length } N_{(i_1+1, d)}^{[j_1]}}.$$

Then by (22) and the observation that  $|w_{i_1+1}^{j_1}| < N_{(i_1+1, i_1+1)}^{[j_1]}/\delta_{j_1}$  for all sufficiently large  $j_1$ , we have

$$\begin{aligned} |w_{i_1}^{j_1}| &< \delta_{j_2} \cdot N_{(i_1+1, i_1)}^{[j_1]} < N_{(i_1+1, i_1)}^{[j_1]}/\delta_{j_1} < \delta_{j_1} \cdot N_{(i_1+1, i_1-1)}^{[j_1]} < N_{(i_1+1, i_1-1)}^{[j_1]}/\delta_{j_1} \\ &< \delta_{j_1} \cdot N_{(i_1+1, i_1-2)}^{[j_1]} < N_{(i_1+1, i_1-2)}^{[j_1]}/\delta_{j_1} < \delta_{j_1} \cdot N_{(i_1+1, i_1-3)}^{[j_1]} < N_{(i_1+1, i_1-3)}^{[j_1]}/\delta_{j_1} \\ &< \cdots < \delta_{j_1} \cdot N_{(i_1+1, 1)}^{[j_1]} < N_{(i_1+1, 1)}^{[j_1]}/\delta_{j_1} < \delta_{j_1} \cdot N_{(i_1+1, d)}^{[j_1]} < N_{(i_1+1, d)}^{[j_1]}/\delta_{j_1} \\ &< \delta_{j_1} \cdot N_{(i_1+1, d-1)}^{[j_1]} < N_{(i_1+1, d-1)}^{[j_1]}/\delta_{j_1} < \cdots < \delta_{j_1} \cdot N_{(i_1+1, i_1+2)}^{[j_1]} \\ &< N_{(i_1+1, i_1+2)}^{[j_1]}/\delta_{j_1} < \delta_{j_1} \cdot N_{(i_1+1, i_1+1)}^{[j_1]} < |w_{i_1+1}^{j_1}| < N_{(i_1+1, i_1+1)}^{[j_1]}/\delta_{j_1}. \end{aligned}$$

Thus there are four possibilities:

- (i)  $|w_{i_1}^{j_1}| \leq n < N_{(i_1+1, i_1)}^{[j_1]}$ ;
- (ii)  $N_{(i_1+1, i_1)}^{[j_1]} \leq n < N_{(i_1+1, i_1-1)}^{[j_1]}$  (indices taken modulo  $d$ );
- (iii)  $N_{(i_1+1, i_1-1)}^{[j_1]} \leq n < N_{(i_1+1, i_1+1)}^{[j_1]}$  and there exists  $i_2 \in \mathcal{A} \setminus \{i_1, i_1 + 1\}$  such that

$$N_{(i_1+1, i_2)}^{[j_1]} \leq n < N_{(i_1+1, i_2-1)}^{[j_1]} \text{ (indices taken modulo } d\text{);}$$

- (iv)  $N_{(i_1+1, i_1+1)}^{[j_1]} \leq n < |w_{i_1+1}^{j_1}|$  (by condition (29)).

In case (i), there are no words of length  $n$  in  $w_{i_1+1}^{j_1} w_{i_1+1}^{j_1}$  that were not previously counted (all blocks in its decomposition are of length larger than  $n$ ). In this case we have the estimate

$$(30) \quad P_X(n) \leq (d-1)\delta_{j_1} n + |w_{i_1}^{j_1}| + dn.$$

In particular, since  $|w_{i_1}^{j_1}| < \delta_{j_1} N_{(i_1+1, i_1)}^{[j_1]}$ , we are in case (i) when  $n = \left\lfloor \frac{|w_{i_1}^{j_1}|}{\delta_{j_1}} \right\rfloor$  and so equation (30) holds. This implies that

$$P_X \left( \left\lfloor \frac{|w_{i_1}^{j_1}|}{\delta_{j_1}} \right\rfloor \right) \leq (d + d\delta_{j_1}) \cdot \left\lfloor \frac{|w_{i_1}^{j_1}|}{\delta_{j_1}} \right\rfloor.$$

This situation arises infinitely often (once for each  $\delta_j$ ) and since  $\delta_j \xrightarrow{j \rightarrow \infty} 0$ ,

$$\liminf_{n \rightarrow \infty} \frac{P_X(n)}{n} \leq d.$$

Combining this with the fact that  $(X, \sigma)$  has at least  $d$  distinct nonatomic ergodic measures and applying Theorem 3.1, we have that

$$\liminf_{n \rightarrow \infty} \frac{P_X(n)}{n} = d.$$

In particular, this implies that there are exactly  $d$  ergodic measures.

In case (ii), we have

$$N_{(i_1+1, i_1)}^{[j_1]} \leq n < N_{(i_1+1, i_1-1)}^{[j_1]}$$

and  $n < N_{(i_1+1, k)}^{[j_1]}$  for all  $k \in \mathcal{A} \setminus \{i_1\}$ . In this case, the only new words of length  $n$  that arise in  $w_{i_1+1}^{j_1} w_{i_1+1}^{j_1}$  are the  $n - N_{(i_1+1, i_1)}^{[j_1]}$  transition words that arise in

$$\underbrace{s_{i_1-1}^{j_1} w_{i_1-1}^{j_1} \cdots w_{i_1-1}^{j_1} p_{i_1-1}^{j_1}}_{\text{length } N_{(i_1+1, i_1-1)}^{[j_1]}} \underbrace{s_{i_1}^{j_1} w_{i_1}^{j_1} \cdots w_{i_1}^{j_1} p_{i_1}^{j_1}}_{\text{length } N_{(i_1+1, i_1)}^{[j_1]}} \underbrace{s_{i_1+1}^{j_1-1} w_{i_1+1}^{j_1-1} \cdots w_{i_1+1}^{j_1-1} p_{i_1+1}^{j_1-1}}_{\text{length } N_{(i_1+1, i_1+1)}^{[j_1]}}$$

where a word is a transition word if it completely contains the middle block (all other blocks have length larger than  $n$  and so contribute no new words). Thus, in case (ii),

$$(31) \quad P_X(n) \leq dn + 2d\delta_{j_1} n + \left( n - N_{(i_1+1, i_1)}^{[j_1]} \right) \leq (d+1)n + 2d\delta_{j_1} n.$$

In case (iii),

$$(32) \quad n \geq N_{(i_1+1, i_2)}^{[j_1]} > \delta_{j_1} \cdot N_{(i_1+1, i_2+1)}^{[j_1]} > \delta_{j_1} \cdot N_{(i_1+1, i_2+2)}^{[j_1]} > \cdots > \delta_{j_1} \cdot N_{(i_1+1, i_1)}^{[j_1]}$$

and

$$n < N_{(i_1+1, i_2-1)}^{[j_1]} < N_{(i_1+1, i_2-2)}^{[j_1]} < \cdots < N_{(i_1+1, i_1+1)}^{[j_1]}$$

by (19), (20), (21), and (22). Therefore the only new words of length  $n$  that arise in  $w_{i_1+1}^{j_1} w_{i_1+1}^{j_1}$  are the transition words that arise in

$$\underbrace{s_{i_2-1}^{j_1} w_{i_2-1}^{j_1} \cdots w_{i_2-1}^{j_1} p_{i_2-1}^{j_1}}_{\text{length } N_{(i_1+1, i_2-1)}^{[j_1]}} \underbrace{s_{i_2}^{j_1} w_{i_2}^{j_1} \cdots w_{i_2}^{j_1} p_{i_2}^{j_1}}_{\text{length } N_{(i_1+1, i_2)}^{[j_1]}} \cdots \underbrace{s_{i_1}^{j_1} w_{i_1}^{j_1} \cdots w_{i_1}^{j_1} p_{i_1}^{j_1}}_{\text{length } N_{(i_1+1, i_1)}^{[j_1]}} \underbrace{s_{i_1+1}^{j_1-1} w_{i_1+1}^{j_1-1} \cdots w_{i_1+1}^{j_1-1} p_{i_1+1}^{j_1-1}}_{\text{length } N_{(i_1+1, i_1+1)}^{[j_1]}}.$$

That is, this word decomposes into blocks such that the first and last of which have length larger than  $n$ ; the transition words are those that fully contain one of the blocks of length smaller than  $n$ . There are at most

$$\left(n - N_{(i_1+1, i_2)}^{[j_1]}\right) + N_{(i_1+1, i_2+1)}^{[j_1]} + N_{(i_1+1, i_2+2)}^{[j_1]} + \cdots + N_{(i_1+1, i_1-1)}^{[j_1]} + N_{(i_1+1, i_1)}^{[j_1]}$$

such blocks. By (32), this is at most  $n - N_{(i_1+1, i_2)}^{[j_1]} + d\delta_{j_1}n$ . So in case (iii),

$$(33) \quad P_X(n) \leq dn + 2d\delta_{j_1}n + n - N_{(i_1+1, i_2)}^{[j_1]} + d\delta_{j_1}n \leq (d+1)n + 3d\delta_{j_1}n.$$

Finally, in case (iv), the only new words are the transition words that occur in

$$\underbrace{s_{i_1+1}^{j_1-1} w_{i_1+1}^{j_1-1} \cdots w_{i_1+1}^{j_1-1} p_{i_1+1}^{j_1-1}}_{\text{length } N_{(i_1+1, i_1+1)}^{[j_1]}} \underbrace{s_{i_1+2}^{j_1-1} w_{i_1+2}^{j_1-1} \cdots w_{i_1+2}^{j_1-1} p_{i_1+2}^{j_1-1} \cdots}_{\text{length } N_{(i_1+1, i_1+2)}^{[j_1]}}$$

$$\underbrace{s_{i_1}^{j_1} w_{i_1}^{j_1} \cdots w_{i_1}^{j_1} p_{i_1}^{j_1}}_{\text{length } N_{(i_1+1, i_1)}^{[j_1]}} \underbrace{s_{i_1+1}^{j_1-1} w_{i_1+1}^{j_1-1} \cdots w_{i_1+1}^{j_1-1} p_{i_1+1}^{j_1-1}}_{\text{length } N_{(i_1+1, i_1+1)}^{[j_1]}} \underbrace{s_{i_1+2}^{j_1-1} w_{i_1+2}^{j_1-1} \cdots w_{i_1+2}^{j_1-1} p_{i_1+2}^{j_1-1} \cdots}_{\text{length } N_{(i_1+1, i_1+2)}^{[j_1]}}$$

$$\underbrace{s_{i_1}^{j_1} w_{i_1}^{j_1} \cdots w_{i_1}^{j_1} p_{i_1}^{j_1}}_{\text{length } N_{(i_1+1, i_1)}^{[j_1]}}$$

where now a word is a transition word if it completely contains any of the blocks of length smaller than  $N_{(i_1+1, i_1+1)}^{[j_1]}$ . However, by (19), (20), (21), and (22), we have

$$\delta_{j_1}n \geq \delta_{j_1}N_{(i_1+1, i_1+1)}^{[j_1]} > N_{(i_1+1, k)}^{[j_1]}$$

for all  $k \in \mathcal{A} \setminus \{i_1 + 1\}$ , and so there are at most  $n + d\delta_{j_1}n$  such words. Thus for case (iv), we have

$$(34) \quad P_X(n) \leq dn + 2d\delta_{j_1}n + n + d\delta_{j_1}n = (d+1)n + 3d\delta_{j_1}n.$$

It follows from (30), (31), (33), and (34) that

$$\limsup_{n \rightarrow \infty} \frac{P_X(n)}{n} \leq d+1$$

and therefore is equal to  $d+1$  by Theorem 3.1.  $\square$

We end with several constructions showing various senses in which our results cannot be improved. We first review some standard facts about Sturmian shifts. A *Sturmian shift*  $(Y, \sigma)$  is a minimal subshift of  $\{0, 1\}^{\mathbb{Z}}$  whose complexity function satisfies  $P_Y(n) = n + 1$  for all  $n \in \mathbb{N}$ . Any Sturmian shift is uniquely ergodic, and for any  $\alpha \in (0, 1) \setminus \mathbb{Q}$ , there exists a Sturmian shift  $(Y_\alpha, \sigma)$  whose unique invariant probability measure  $\mu$  satisfies  $\mu([0]) = \alpha$ . In particular, there are uncountably many distinct Sturmian shifts.

We first show that the technical condition (that there exists a generic measure  $\mu$  and a generic point  $x_\mu$  such that the orbit closure of  $x_\mu$  is not uniquely ergodic) cannot be dropped from the second statement in Theorem 3.1:

**Proposition 5.1.** *For  $d \geq 1$ , there exists a subshift  $(X, \sigma)$  which has precisely  $d$  ergodic measures, zero nonergodic generic measures, and whose complexity function satisfies  $P_X(n) = dn + d$  for all  $n \in \mathbb{N}$ . This subshift has the property that every  $x \in X$  is generic for some ergodic measure and the orbit closure of any point is uniquely ergodic.*

*Proof.* Fix  $d \in \mathbb{N}$  and fix a Sturmian shift  $(Y, \sigma)$  on the alphabet  $\{0, 1\}$ . Let  $\mathcal{A} := \{0_1, 1_1, 0_2, 1_2, \dots, 0_d, 1_d\}$  and for  $1 \leq i \leq d$  let  $Y_i \subset \mathcal{A}^{\mathbb{Z}}$  be the image of  $(Y, \sigma)$  under the 1-block code that sends  $0 \mapsto 0_i$  and  $1 \mapsto 1_i$ . Let

$$X := \bigcup_{i=1}^d Y_i \subset \mathcal{A}^{\mathbb{Z}}$$

and observe that  $X$  is closed and  $\sigma$ -invariant. Moreover, we have  $P_X(n) = dn + d$  for all  $n \in \mathbb{N}$ . Each subshift  $Y_i \subset X$  supports a unique ergodic measure and so there are at least  $d$  ergodic measures for  $(X, \sigma)$ . Conversely, for each  $x \in X$  there exists  $1 \leq i \leq d$  such that  $x \in Y_i$ . Since  $Y_i$  is uniquely ergodic,  $x$  is generic for the (unique) ergodic measure supported on  $Y_i$ . Thus there can be no other measures that have a generic point.  $\square$

Finally we show that the assumption of linear growth in Theorem 1.1 is optimal, in the sense that there is no analog of Theorem 1.1 with an assumption of a super-linear growth rate and conclusion that the set of ergodic measures is finite for all subshifts whose complexity function grows at most at that rate.

**Proposition 5.2.** *Let  $(p_n)_{n=1}^{\infty}$  be a sequence of real numbers such that*

$$\liminf_{n \rightarrow \infty} \frac{p_n}{n} = \infty.$$

*Then there exists a subshift  $(X, \sigma)$  which has infinitely many nonatomic ergodic measures and is such that for all but finitely many  $n$ , we have  $P_X(n) \leq p_n$ .*

*Proof.* For each  $n \in \mathbb{N}$ , there exists a set  $\mathcal{F}_n \subset \{0, 1\}^n$  such that  $|\mathcal{F}_n| = n + 1$  and for uncountably many  $\alpha \in (0, 1)$ , we have  $\mathcal{L}_n(Y_\alpha) = \mathcal{F}_n$ . For  $N \leq n$ , let  $\mathcal{X}_N(\mathcal{F}_n)$  be the set of words of length  $N$  that arise as a subword of a word in  $\mathcal{F}_n$ . Clearly if  $\mathcal{L}_n(Y_\alpha) = \mathcal{F}_n$  then  $\mathcal{L}_N(Y_\alpha) = \mathcal{X}_N(\mathcal{F}_n)$ . Let  $\mathcal{G}_1 \subset \{0, 1\}$  be such that for infinitely many  $n \in \mathbb{N}$  we have  $\mathcal{G}_1 = \mathcal{X}_1(\mathcal{F}_n)$ . Inductively, we assume that we have defined  $\mathcal{G}_i \subset \{0, 1\}^i$  for all  $1 \leq i < j$  such that

- (i) For all  $1 \leq j_1 < j_2 < j$  we have  $\mathcal{G}_{j_1} = \mathcal{X}_{j_1}(\mathcal{G}_{j_2})$ ;
- (ii) There are infinitely many  $n$  for which  $\mathcal{G}_{j-1} = \mathcal{X}_{j-1}(\mathcal{F}_n)$ .

We then choose  $\mathcal{G}_j \subset \{0, 1\}^j$  such that among those  $n$  for which  $\mathcal{G}_{j-1} = \mathcal{X}_{j-1}(\mathcal{F}_n)$ , there are infinitely many  $n$  for which  $\mathcal{G}_j = \mathcal{X}_j(\mathcal{F}_n)$ . In this way, we obtain an infinite sequence  $\mathcal{G}_1, \mathcal{G}_2, \dots$  such that if  $1 \leq j_1 < j_2$ , then  $\mathcal{G}_{j_1} = \mathcal{X}_{j_1}(\mathcal{G}_{j_2})$  and there are uncountably many  $\alpha \in (0, 1)$  for which  $\mathcal{L}_{j_2}(Y_\alpha) = \mathcal{G}_{j_2}$ .

For each  $n \in \mathbb{N}$ , set

$$A_n := \{\alpha \in (0, 1) : \mathcal{L}_n(Y_\alpha) = \mathcal{G}_n\}.$$

Then by construction,  $A_n$  is uncountable for all  $n \in \mathbb{N}$ ,

$$A_1 \supseteq A_2 \supseteq A_3 \supseteq \dots$$

and for infinitely many  $n \in \mathbb{N}$  we have  $A_n \neq A_{n+1}$ . (If not, there exist distinct  $\alpha_1, \alpha_2 \in \cap A_n$ , and so  $\mathcal{L}_n(Y_{\alpha_1}) = \mathcal{L}_n(Y_{\alpha_2})$  for all  $n$ , contradicting the fact that the frequency with which the letter 0 occurs as a subword of any word in  $\mathcal{L}_n(Y_{\alpha_i})$  tends to  $\alpha_i$  for  $i = 1, 2$ .)

We now construct the subshift. Find  $N_1 \in \mathbb{N}$  such that for all  $n \geq N_1$  we have  $p_n > 2n + 2$ . Choose the smallest  $M_1 \geq N_1$  for which  $A_{M_1+1} \neq A_{M_1}$  and let  $\alpha_1 \in A_{M_1} \setminus A_{M_1+1}$ . Set  $X_1 := Y_{\alpha_1}$  and observe that  $P_{X_1}(n) = n + 1$  for all  $n$ . Now find  $N_2 \in \mathbb{N}$  such that for all  $n \geq N_2$  we have  $p_n > 3n + 3$ . Find the

smallest  $M_2 \geq N_2$  for which  $A_{M_2+1} \neq A_{M_2}$  and let  $\alpha_2 \in A_{M_2} \setminus A_{M_2+1}$ . Set  $X_2 := Y_{\alpha_1} \cup Y_{\alpha_2}$ . Then, by construction,  $\alpha_2 \in A_{M_1}$  and so  $\mathcal{L}_{M_1}(Y_{\alpha_2}) = \mathcal{L}_{M_1}(Y_{\alpha_1})$  but since  $\alpha_2 \in A_{M_2} \subset A_{M_1+1}$  we know that  $\mathcal{L}_{M_1+1}(Y_{\alpha_2}) \neq \mathcal{L}_{M_1+1}(Y_{\alpha_1})$ . Consequently  $P_{X_2}(n) = n + 1$  for all  $n \leq M_1$  and  $n + 1 < P_{X_2}(n) \leq 2n + 2$  for all  $n > M_1$ . Now recursively suppose we have chosen integers  $M_1 < M_2 < \dots < M_i$  such that for each  $1 \leq k \leq i$  we have  $p_n > (k + 1)n + (k + 1)$  for all  $n \geq M_k$  and moreover that  $A_{M_k+1} \neq A_{M_k}$ . Suppose further that we have chosen  $\alpha_1, \dots, \alpha_i$  such that  $\alpha_k \in A_{M_k} \setminus A_{M_k+1}$  for each  $k$ . Finally suppose  $X_i := Y_{\alpha_1} \cup \dots \cup Y_{\alpha_i}$ . Then, by construction, we have  $P_{X_i}(n) \leq kn + k$  for all  $n \leq M_k$ . Find  $N_{i+1} \in \mathbb{N}$  such that  $p_n > (i + 2)n + (i + 2)$  for all  $n \geq N_{i+1}$  and let  $M_{i+1}$  be the smallest integer larger than  $N_{i+1}$  for which  $A_{M_{i+1}} \neq A_{M_{i+1}+1}$ . Let  $\alpha_{i+1} \in A_{M_{i+1}} \setminus A_{M_{i+1}+1}$ . Define  $X_{i+1} := Y_{\alpha_1} \cup \dots \cup Y_{\alpha_i} \cup Y_{\alpha_{i+1}}$ . Since  $\alpha_{i+1} \in A_{M_{i+1}} \subset A_{M_i}$  we know that  $\mathcal{L}_k(X_i) = \mathcal{L}_k(X_{i+1})$  for all  $k \leq M_i$  and since  $\alpha_{i+1} \in A_{M_{i+1}}$  we know that  $\mathcal{L}_{M_{i+1}}(X_i) \neq \mathcal{L}_{M_{i+1}}(X_{i+1})$ . Consequently, we have  $P_{X_{i+1}}(n) = P_{X_i}(n)$  for all  $n \leq M_i$  and  $P_{X_{i+1}}(n) \leq (i + 1)n + (i + 1)$  for all  $n > M_i$ . By construction,  $p_n > (i + 1)n + (i + 1)$  for all  $n \geq M_i$ . Therefore our recursive construction continues for all  $i \in \mathbb{N}$ .

Thus we obtain a sequence of subshifts

$$X_1 \subset X_2 \subset X_3 \subset \dots$$

such that for all  $i \in \mathbb{N}$  and all  $n \geq N_1$ , we have  $P_{X_i}(n) < p_n$ . Setting

$$X := \overline{\bigcup_{i=1}^{\infty} X_i},$$

we have that  $\mathcal{L}_n(X) = \bigcup_{i=1}^{\infty} \mathcal{L}_n(X_i)$  for all  $n \in \mathbb{N}$ . Therefore, for all  $n \geq N_1$ , the complexity satisfies  $P_X(n) < p_n$ . On the other hand, for all  $i \in \mathbb{N}$  we have  $Y_{\alpha_i} \subset X$  and there is an ergodic probability supported on  $Y_{\alpha_i}$ . Since  $Y_{\alpha_i} \neq Y_{\alpha_j}$  for all  $i \neq j$  by construction,  $X$  has infinitely many ergodic measures.  $\square$

## REFERENCES

- [1] M. BOSHERNITZAN. A unique ergodicity of minimal symbolic flows with linear block growth. *Journal d'Analyse Math.* **44**, no. 1 (1984) 77–96.
- [2] M. BOSHERNITZAN. A condition for unique ergodicity of minimal symbolic flows. *Ergodic Theory Dynam. Systems* **12**, no. 3 (1992) 425–428.
- [3] M. BOSHERNITZAN. A condition for minimal interval exchange maps to be uniquely ergodic. *Duke Math. J.* **52** (1985), no. 3, 723–752.
- [4] J. CHAIKA & H. MASUR. There exists an interval exchange with a non-ergodic generic measure. *J. Mod. Dyn.* **9** (2015), 289–304.
- [5] M. DAMRON & J. FICKENSCHER. On the number of ergodic measures for minimal shifts with eventually constant complexity growth. To appear, *Ergodic Theory Dynam. Systems*. arXiv:1508.05952.
- [6] M. DENKER, C. GRILLENBERGER, & K. SIGMUND. Ergodic theory on compact spaces. Lecture Notes in Mathematics, Vol. 527. Springer-Verlag, Berlin-New York, 1976.
- [7] C. EPIFANIO, M. KOSKAS, & F. MIGNOSI. On a conjecture on bidimensional words. *Theor. Comp. Science* **299** (2003), 123–150.
- [8] S. FERENCZI & L. ZAMBONI. Languages of  $k$ -interval exchange transformations. *Bull. Lond. Math. Soc.* **40** no. 4, (2008) 705–714.
- [9] N. FINE & H. WILF. Uniqueness theorem for periodic functions. *Proc. Amer. Math. Soc.* **16** (1965), 109–114.
- [10] A. KATOK. Invariant measures of flows on orientable surfaces. *Dokl. Akad. Nauk SSSR* **211** (1973) 775–778.

- [11] A. KATOK & B. HASSELBLATT. Introduction to the modern theory of dynamical systems. Encyclopedia of Mathematics and its Applications, 54. Cambridge University Press, Cambridge, 1995
- [12] M. KEANE. Interval exchange transformations. *Math. Z.* **141** (1975), 25–31.
- [13] A. QUAS. Math Overflow. <http://mathoverflow.net/questions/152470/getting-unique-ergodicity-from-minimality/152524#152524>
- [14] W. VEECH. Interval exchange transformations. *J. Analyse Math.* **33** (1978) 222-272.

BUCKNELL UNIVERSITY, LEWISBURG, PA 17837 USA  
*E-mail address:* `van.cyr@bucknell.edu`

NORTHWESTERN UNIVERSITY, EVANSTON, IL 60208 USA  
*E-mail address:* `kra@math.northwestern.edu`