

## EASY PUTNAM PROBLEMS

(Last updated: December 11, 2017)

REMARK. The problems in the Putnam Competition are usually very hard, but practically every session contains at least one problem very easy to solve—it still may need some sort of ingenious idea, but the solution is very simple. This is a list of “easy” problems that have appeared in the Putnam Competition in past years—Miguel A. Lerma

**2017-A1.** Let  $S$  be the smallest set of positive integers such that

- (a) 2 is in  $S$ ,
- (b)  $n$  is in  $S$  whenever  $n^2$  is in  $S$ , and
- (c)  $(n + 5)^2$  is in  $S$  whenever  $n$  is in  $S$ .

Which positive integers are not in  $S$ ?

(The set  $S$  is “smallest” in the sense that  $S$  is contained in any other such set.)

**2017-B1.** Let  $L_1$  and  $L_2$  be distinct lines in the plane. Prove that  $L_1$  and  $L_2$  intersect if and only if, for every real number  $\lambda \neq 0$  and every point  $P$  not on  $L_1$  or  $L_2$ , there exist points  $A_1$  on  $L_1$  and  $A_2$  on  $L_2$  such that  $\overrightarrow{PA_2} = \lambda \overrightarrow{PA_1}$ .

**2016-A1.** Find the smallest positive integer  $j$  such that for every polynomial  $p(x)$  with integer coefficients and for every  $k$ , the integer

$$p^{(j)}(k) = \left. \frac{d^j}{dx^j} p(x) \right|_{x=k}$$

(the  $j$ -th derivative of  $p(x)$  at  $k$ ) is divisible by 2016.

**2016-B1.** Let  $x_0, x_1, x_2, \dots$  be the sequence such that  $x_0 = 1$  and for  $n \geq 0$ ,

$$x_{n+1} = \ln(e^{x_n} - x_n)$$

(as usual, the function  $\ln$  is the natural logarithm). Show that the infinite series

$$x_0 + x_1 + x_2 + \dots$$

converges and find its sum.

**2016-B3.** Suppose that  $S$  is a finite set of points in the plane such that the area of triangle  $\triangle ABC$  is at most 1 whenever  $A, B$ , and  $C$  are in  $S$ . Show that there exists a triangle of area 4 such that (together with its interior) covers the set  $S$ .

**2015-A1.** Let  $A$  and  $B$  be points on the same branch of the hyperbola  $xy = 1$ . Suppose that  $P$  is a point lying between  $A$  and  $B$  on this hyperbola, such that the area of the triangle  $APB$  is as large as possible. Show that the region bounded by the hyperbola and the

chord  $AP$  has the same area as the region bounded by the hyperbola and the chord  $PB$ .

**2015-B1.** Let  $f$  be a three times differentiable function (defined on  $\mathbb{R}$  and real-valued) such that  $f$  has at least five distinct real zeros. prove that  $f + 6f' + 12f'' + 8f'''$  has at least two distinct real zeros.

**2015-B4.** Let  $T$  be the set of all triples  $(a, b, c)$  of positive integers for which there exist triangles with side lengths  $a, b, c$ . Express

$$\sum_{(a,b,c) \in T} \frac{2^a}{3^b 5^c}$$

as a rational number in lowest terms.

**2014-A1.** Prove that every nonzero coefficient of the Taylor series of

$$(1 - x + x^2)e^x$$

about  $x = 0$  is a rational number whose numerator (in lowest terms) is either 1 or a prime number.

**2014-B1.** A *base 10 over-expansion* of a positive integer  $N$  is an expression of the form

$$N = d_k 10^k + d_{k-1} 10^{k-1} + \cdots + d_0 10^0$$

with  $d_k \neq 0$  and  $d_i \in \{0, 1, 2, \dots, 10\}$  for all  $i$ . For instance, the integer  $N = 10$  has two base 10 over-expansions:  $10 = 10 \cdot 10^0$  and the usual base 10 expansion  $10 = 1 \cdot 10^1 + 0 \cdot 10^0$ . Which positive integers have a unique base 10 over-expansion?

**2013-A1.** Recall that a regular icosahedron is a convex polyhedron having 12 vertices and 20 faces; the faces are congruent equilateral triangles. On each face of a regular icosahedron is written a nonnegative integer such that the sum of all 20 integers is 39. Show that there are two faces that share a vertex and have the same integer written on them.

**2013-B1.** For positive integers  $n$ , let the numbers  $c(n)$  be determined by the rules  $c(1) = 1$ ,  $c(2n) = c(n)$ , and  $c(2n+1) = (-1)^n c(n)$ . Find the value of

$$\sum_{n=1}^{2013} c(n)c(n+2).$$

**2012-A1.** Let  $d_1, d_2, \dots, d_{12}$  be real numbers in the interval  $(1, 12)$ . Show that there exist distinct indices  $i, j, k$  such that  $d_i, d_j, d_k$  are the side lengths of an acute triangle.

**2012-B1.** Let  $S$  be the class of functions from  $[0, \infty)$  to  $[0, \infty)$  that satisfies:

- (i) The functions  $f_1(x) = e^x - 1$  and  $f_2(x) = \ln(x+1)$  are in  $S$ .
- (ii) If  $f(x)$  and  $g(x)$  are in  $S$ , then functions  $f(x) + g(x)$  and  $f(g(x))$  are in  $S$ ;
- (iii) If  $f(x)$  and  $g(x)$  are in  $S$  and  $f(x) \geq g(x)$  for all  $x \geq 0$ , then the function  $f(x) - g(x)$  is in  $S$ .

Prove that if  $f(x)$  and  $g(x)$  are in  $S$ , then the function  $f(x)g(x)$  is also in  $S$ .

**2011-B1.** Let  $h$  and  $k$  be positive integers. Prove that for every  $\varepsilon > 0$ , there are positive integers  $m$  and  $n$  such that

$$\varepsilon < |h\sqrt{m} - k\sqrt{n}| < 2\varepsilon.$$

**2011-A1.** Given a positive integer  $n$ , what is the largest  $k$  such that the numbers  $1, 2, \dots, n$  can be put into  $k$  boxes so that the sum of the numbers in each box is the same? [When  $n = 8$ , the example  $\{1, 2, 3, 6\}, \{4, 8\}, \{5, 7\}$  shows that the largest  $k$  is *at least* 3.]

**2010-B1.** Is there an infinite sequence of real numbers  $a_1, a_2, a_3, \dots$  such that

$$a_1^m + a_2^m + a_3^m + \dots = m$$

for every positive integer  $m$ ?

**2010-B2.** Given that  $A, B$ , and  $C$  are noncollinear points in the plane with integer coordinates such that the distances  $AB, AC$ , and  $BC$  are integers, what is the smallest possible value of  $AB$ ?

**2009-A1.** Let  $f$  be a real-valued function on the plane such that for every square  $ABCD$  in the plane,  $f(A) + f(B) + f(C) + f(D) = 0$ . Does it follow that  $f(P) = 0$  for all points  $P$  in the plane?

**2009-B1.** Show that every positive rational number can be written as a quotient of products of factorials of (not necessarily distinct) primes. For example,

$$\frac{10}{9} = \frac{2! \cdot 5!}{3! \cdot 3! \cdot 3!}.$$

**2008-A1.** Let  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$  be a function such that  $f(x, y) + f(y, z) + f(z, x) = 0$  for all real numbers  $x, y$ , and  $z$ . Prove that there exists a function  $g : \mathbb{R} \rightarrow \mathbb{R}$  such that  $f(x, y) = g(x) - g(y)$  for all real numbers  $x$  and  $y$ .

**2008-A2.** Alan and Barbara play a game in which they take turns filling entries of an initially empty  $2008 \times 2008$  array. Alan plays first. At each turn, a player chooses a real number and places it in a vacant entry. The game ends when all the entries are filled. Alan wins if the determinant of the resulting matrix is nonzero; Barbara wins if it is zero. Which player has a winning strategy?

**2008-B1.** What is the maximum number of rational points that can lie on a circle in  $\mathbb{R}^2$  whose center is not a rational point? (A *rational point* is a point both of whose coordinates are rational numbers.)

**2007-A1.** Find all values of  $\alpha$  for which the curves  $y = \alpha x^2 + \alpha x + \frac{1}{24}$  and  $x = \alpha y^2 + \alpha y + \frac{1}{24}$  are tangent to each other.

**2007-B1.** Let  $f$  be a polynomial with positive integer coefficients. Prove that if  $n$  is a positive integer, then  $f(n)$  divides  $f(f(n) + 1)$  if and only if  $n = 1$ . [Note: one must assume  $f$  is nonconstant.]

**2006-A1.** Find the volume of the region of points  $(x, y, z)$  such that

$$(x^2 + y^2 + z^2 + 8)^2 \leq 36(x^2 + y^2).$$

**2006-B2.** Prove that, for every set  $X = \{x_1, x_2, \dots, x_n\}$  of  $n$  real numbers, there exists a non-empty subset  $S$  of  $X$  and an integer  $m$  such that

$$\left| m + \sum_{s \in S} s \right| \leq \frac{1}{n+1}.$$

**2005-A1.** Show that every positive integer is a sum of one or more numbers of the form  $2^r 3^s$ , where  $r$  and  $s$  are nonnegative integers and no summand divides another. (For example,  $23 = 9 + 8 + 6$ .)

**2005-B1.** Find a nonzero polynomial  $P(x, y)$  such that  $P(\lfloor a \rfloor, \lfloor 2a \rfloor) = 0$  for all real numbers  $a$ . (Note:  $\lfloor \nu \rfloor$  is the greatest integer less than or equal to  $\nu$ .)

**2004-A1.** Basketball star Shanille O'Keal's team statistician keeps track of the number,  $S(N)$ , of successful free throws she has made in her first  $N$  attempts of the season. Early in the season,  $S(N)$  was less than 80% of  $N$ , but by the end of the season,  $S(N)$  was more than 80% of  $N$ . Was there necessarily a moment in between when  $S(N)$  was exactly 80% of  $N$ ?

**2004-B2.** Let  $m$  and  $n$  be positive integers. Show that

$$\frac{(m+n)!}{(m+n)^{m+n}} < \frac{m! n!}{m^m n^n}.$$

**2003-A1.** Let  $n$  be a fixed positive integer. How many ways are there to write  $n$  as a sum of positive integers,  $n = a_1 + a_2 + \dots + a_k$ , with  $k$  an arbitrary positive integer and  $a_1 \leq a_2 \leq \dots \leq a_k \leq a_1 + 1$ ? For example, with  $n = 4$  there are four ways: 4, 2+2, 1+1+2, 1+1+1+1.

**2002-A1.** Let  $k$  be a fixed positive integer. The  $n$ -th derivative of  $\frac{1}{x^k-1}$  has the form  $\frac{P_n(x)}{(x^k-1)^{n+1}}$  where  $P_n(x)$  is a polynomial. Find  $P_n(1)$ .

**2002-A2.** Given any five points on a sphere, show that some four of them must lie on a closed hemisphere.

**2001-A1.** Consider a set  $S$  and a binary operation  $*$ , i.e., for each  $a, b \in S$ ,  $a * b \in S$ . Assume  $(a * b) * a = b$  for all  $a, b \in S$ . Prove that  $a * (b * a) = b$  for all  $a, b \in S$ .

**2000-A2.** Prove that there exist infinitely many integers  $n$  such that  $n, n+1, n+2$  are each the sum of the squares of two integers. [Example:  $0 = 0^2 + 0^2$ ,  $1 = 0^2 + 1^2$ ,  $2 = 1^2 + 1^2$ .]

**1999-A1.** Find polynomials  $f(x)$ ,  $g(x)$ , and  $h(x)$ , if they exist, such that for all  $x$ ,

$$|f(x)| - |g(x)| + h(x) = \begin{cases} -1 & \text{if } x < -1 \\ 3x + 2 & \text{if } -1 \leq x \leq 0 \\ -2x + 2 & \text{if } x > 0. \end{cases}$$

**1998-A1.** A right circular cone has base of radius 1 and height 3. A cube is inscribed in the cone so that one face of the cube is contained in the base of the cone. What is the side-length of the cube?

**1997-A5.** Let  $N_n$  denote the number of ordered  $n$ -tuples of positive integers  $(a_1, a_2, \dots, a_n)$  such that  $1/a_1 + 1/a_2 + \dots + 1/a_n = 1$ . Determine whether  $N_{10}$  is even or odd.

**1988-B1.** A *composite* (positive integer) is a product  $ab$  with  $a$  and  $b$  not necessarily distinct integers in  $\{2, 3, 4, \dots\}$ . Show that every composite is expressible as  $xy + xz + yz + 1$ , with  $x$ ,  $y$ , and  $z$  positive integers.